

## **Evidence Report:**

# **Risk of Impaired Control of Spacecraft/Associated Systems and Decreased Mobility Due to Vestibular/Sensorimotor Alterations Associated with Space flight**

Approved for Public Release: June 6, 2016

## **Human Research Program Human Health Countermeasures Element**

National Aeronautics and Space Administration  
Lyndon B. Johnson Space Center  
Houston, Texas

**CURRENT CONTRIBUTING AUTHORS:**

Jacob J. Bloomberg	NASA Johnson Space Center, Houston, TX
Millard F. Reschke	NASA Johnson Space Center, Houston, TX
Gilles R. Clément	Wyle Science Technology & Engineering Group, Houston, TX
Ajitkumar P. Mulavara	Universities Space Research Association, Houston, TX
Laura C. Taylor	Wyle Science Technology & Engineering Group, Houston, TX

**PREVIOUS CONTRIBUTING AUTHORS:**

William H. Paloski	NASA Johnson Space Center, Houston, TX
Charles M. Oman	Massachusetts Institute of Technology, Cambridge, MA
Scott J. Wood	Azusa Pacific University, Azusa, CA
Deborah L. Harm	NASA Johnson Space Center, Houston, TX
Brian T. Peters	Wyle Science Technology & Engineering Group, Houston, TX
James P. Locke	NASA Johnson Space Center, Houston, TX
Leland S. Stone	NASA Ames Research Center, Moffett Field, CA

## TABLE OF CONTENTS

I.	PRD RISK TITLE: RISK OF IMPAIRED CONTROL OF SPACECRAFT/ASSOCIATED SYSTEMS AND DECREASED MOBILITY DUE TO VESTIBULAR/SENSORIMOTOR ALTERATIONS ASSOCIATED WITH SPACE FLIGHT.....	7
II.	EXECUTIVE SUMMARY.....	7
III.	INTRODUCTION.....	8
IV.	EVIDENCE.....	8
	1. Evidence Obtained from Space Flight Operations.....	9
	1.1. Crew Verbal Reports.....	9
	1.2. Shuttle Entry and Landing Spatial Disorientation.....	12
	1.3. Apollo Lunar Landing Spatial Disorientation.....	14
	1.4. Apollo Landing Geographic Disorientation.....	15
	1.5. Rendezvous and Docking.....	16
	1.6. Teleoperator Tasks.....	17
	1.7. Driving Performance.....	17
	1.8. Implications for Exploration Class Vehicle Design.....	18
	1.9. Walking on the Moon.....	18
	2. Evidence Obtained from Scientific Investigations In Flight.....	20
	2.1. Decrements in Sensation .....	21
	2.1.1. Visual Acuity .....	21
	2.1.2. Proprioception.....	22
	2.1.3. Motion Perception.....	23
	2.2. Decrements in Gaze Control.....	24
	2.2.1. Smooth Pursuit Eye Movements.....	24
	2.2.2. Vestibulo-Ocular Reflex.....	24
	2.2.3. Eye-Head Coordination and Target Acquisition.....	25
	2.2.4. Optokinetic Nystagmus.....	26
	2.3. Decrements in Eye-Hand Coordination Performance.....	26
	2.3.1. Control of Aimed Arm Movements.....	27
	2.3.2. Reaching and Grasping.....	27
	2.3.3. Manual Tracking.....	27
	2.3.4. Force Discrimination and Control.....	28
	2.3.5. Fine Motor Control.....	28
	2.3.6. Dual Tasking and Manual Performance.....	29
	2.4. Decrements in Spatial Orientation Perception.....	29
	2.4.1. Spatial Disorientation in Aviation.....	29
	2.4.2. Geographic Disorientation and Terrain Awareness.....	30
	2.4.3. Spatial Disorientation During Space Flight.....	31

2.4.4. Visual Reorientation Illusion .....	32
2.4.5. Tilt-Translation and Tilt-Gain Illusions.....	33
2.4.6. Physiological Basis for Spatial Disorientation.....	35
2.5. Decrements in Cognitive Function.....	35
2.5.1. Mental Rotation.....	36
2.5.2. Distance and Size Perception.....	37
2.5.3. Time Perception.....	38
2.6. Space Motion Sickness.....	39
2.6.1. History.....	39
2.6.2. Symptoms.....	39
2.6.3. Incidence.....	40
2.6.4. Provocative Stimuli.....	41
2.6.5. Theories and Hypotheses.....	41
2.6.6. Predicting Susceptibility.....	42
2.7. Impacts of Vestibular Changes to Orthostatic Intolerance.....	42
2.8. Peripheral and Central Neural Changes Associated with Space flight.....	43
3. Evidence Obtained from Scientific Investigations During Reentry and After Landing....	45
3.1. Decrements in Postural Equilibrium.....	45
3.2. Decrements in Locomotion.....	47
3.2.1. Locomotor Control and Central Activation.....	47
3.2.2. Head-Trunk Coordination.....	49
3.2.3. Dynamic Visual Acuity.....	51
3.3. Decrements in Jump Performance.....	53
3.4. Perceptual Changes.....	54
3.4.1. Crew Reports.....	54
3.4.2. Proprioceptive Changes.....	55
3.4.3. Perceived Tilt and Translation.....	55
3.4.4. Otolith Asymmetry .....	57
3.5. Gaze Changes .....	58
3.5.1. Vestibulo-Ocular Reflex.....	58
3.5.2. Ocular Counterrolling.....	59
3.5.3. Eye Movements during Coriolis and Cross-Coupled Acceleration.....	59
3.5.4. Eye Movements during Linear Acceleration.....	60
3.6. Decrements in Functional Performance.....	61
3.6.1. Functional Mobility.....	61
3.6.2. Functional Tasks.....	62
3.6.3. Field Tests.....	65
3.6.4. Navigation.....	65
3.7. Decrements in Manual Control .....	65
3.8. Impacts of Vestibular Changes to Orthostatic Intolerance.....	65

V. COUNTERMEASURES.....	67
1. Pharmacological Countermeasures.....	67
1.1. Drug Classes.....	67
1.2. In-flight Medication.....	67
2. Training and Rehabilitation.....	68
2.1. Vestibular Training.....	68
2.2. Preflight Adaptation Training.....	69
2.3. Sensorimotor Adaptability Training .....	70
2.4. Biofeedback Training.....	72
2.5. In-flight Exercise.....	72
2.6. Head Movements During and After Reentry.....	73
2.7. Balance Rehabilitation Therapy.....	73
3. Mechanical Devices.....	74
3.1. Dynamic Foot Stimulation.....	74
3.2. Load Suits.....	76
3.3. Stroboscopic Goggles to Mitigate Motion Sickness.....	76
3.4. Tactile Spatial Awareness System.....	77
3.5. Electrical Devices.....	77
3.5.1. Galvanic Vestibular Stimulation.....	77
3.5.2. Vestibular Stochastic Resonance.....	78
4. Artificial Gravity.....	79
VI. COMPUTER-BASED MODELING AND SIMULATION.....	83
1. Models of Vestibular Function and Spatial Orientation.....	83
2. Models of Manual Control Performance.....	84
VII. RISK IN CONTEXT OF EXPLORATION MISSION OPERATIONAL SCENARIOS.....	87
1. Sensorimotor Standard.....	87
2. Risks During Piloted Landings.....	88
3. Risks During Vehicle Egress and Extravehicular Activities.....	88
4. Risks During Rover Operations.....	89
5. Risks During Rendezvous/Docking and Remote Manipulator System Operations.....	90
6. Risks While Operating Other Complex Systems.....	90
7. Risks During Near-Term Missions.....	90
VIII. GAPS.....	91
IX. CONCLUSION.....	92
X. REFERENCES.....	93
XI. TEAM.....	126

XII. LIST OF ACRONYMS.....	126
XIII. APPENDIX A: SHUTTLE LANDING PERFORMANCE .....	129
XIV. APPENDIX B: APOLLO LUNAR LANDING PERFORMANCE.....	144
XV. APPENDIX C: INCIDENTS DUE TO SITUATIONAL AWARENESS DURING EVA.....	149

## I. RISK TITLE: RISK OF IMPAIRED CONTROL OF SPACECRAFT/ASSOCIATED SYSTEMS AND DECREASED MOBILITY DUE TO VESTIBULAR/SENSORIMOTOR ALTERATIONS ASSOCIATED WITH SPACE FLIGHT

The National Aeronautics and Space Administration's (NASA) Human Research Program (HRP) has identified a number of potentially significant biomedical risks that might limit the agency's plans for future space exploration, including missions back to the Moon and on to Mars. Among these risks is the ***Risk of Impaired Control of Spacecraft/Associated Systems and Decreased Mobility Due to Vestibular/Sensorimotor Alterations Associated with Space flight.*** This risk is described as follows: *"Given that there is an alteration in vestibular/sensorimotor function during and immediately following gravitational transitions manifested as changes in eye-head-hand control, postural and/or locomotor ability, gaze function, and perception, there is a possibility that crew will experience impaired control of the spacecraft during landing or decreased mobility following a landing on a planetary surface (Earth or other) after long-duration space flight. These changes have not specifically been correlated with real time performance decrements. The risk of impairment is greatest during and soon after G-transitions when performance decrements may have high operational impact (landing, immediate egress following landing). The possible alterations in sensorimotor performance are of interest for Mars missions due to the prolonged microgravity exposure during transit followed by landing tasks. This risk must be better defined and documented and vestibular/sensorimotor changes must be correlated with performance issues."*

## II. EXECUTIVE SUMMARY

We examine the various dimensions of the ***Risk of Impaired Control of Spacecraft/Associated Systems and Decreased Mobility Due to Vestibular/Sensorimotor Alterations Associated with Space flight*** by reviewing the research and operational evidence demonstrating sensorimotor performance decrements during and after space flight. These sensorimotor performance decrements might affect vehicle and complex system control, including decreased visual acuity, eye-hand coordination, spatial and geographic orientation perception, and cognitive function. Sensorimotor performance decrements might also affect the ability to egress and walk away from the vehicle in case of an emergency or an extravehicular activity on a planetary surface. We also review the countermeasures that have been tested, including medication, prevention techniques and training exercises, physical rehabilitation, and mechanical devices. Furthermore, we identify the current knowledge and mitigation gaps that must be filled through further research and/or data mining efforts before the risk can be fully mitigated. We conclude that the true operational risks associated with the impacts of adaptive sensorimotor changes on mobility and crew abilities to control vehicles and other complex systems will only be estimable after the gaps have been filled and we have been able to accurately assess integrated performance in off-nominal operational settings (Shelhamer 2015).

### III. INTRODUCTION

Control of vehicles and other complex systems is a high-level integrative function of the central nervous system (CNS). It requires well-functioning subsystem performance, including good visual acuity, eye-hand coordination, spatial and geographic orientation perception, and cognitive function. Evidence from space flight research demonstrates that the function of each of these subsystems is altered by removing gravity, a fundamental orientation reference, which is sensed by vestibular, proprioceptive, and haptic receptors and used by the CNS for spatial orientation, posture, navigation, and coordination of movements. The available evidence also shows that the degree of alteration of each subsystem depends on a number of crew- and mission-related factors.

There is only limited operational evidence that these alterations cause functional impacts on mission-critical vehicle (or complex system) control capabilities. Furthermore, while much of the operational performance data collected during space flight has not been available for independent analysis, those that have been reviewed are somewhat equivocal owing to uncontrolled (and/or unmeasured) environmental and/or engineering factors. Whether this can be improved by further analysis of previously inaccessible operational data or by development of new operational research protocols remains to be seen. The true operational risks will be estimable only after we have filled the knowledge gaps and when we can accurately assess integrated performance in off-nominal operational settings (Paloski *et al.* 2008).

Thus, our current understanding of the ***Risk of Impaired Control of Spacecraft/Associated Systems and Decreased Mobility Due to Vestibular/Sensorimotor Alterations Associated with Space flight*** is limited primarily to extrapolation of scientific research findings, and, since there are limited ground-based analogs of the sensorimotor and vestibular changes associated with space flight, observation of their functional impacts is limited to studies performed in the space flight environment. Fortunately, many sensorimotor and vestibular experiments have been performed during and/or after space flight missions since 1959 (Reschke *et al.* 2007). While not all of these experiments were directly relevant to the question of vehicle/complex system control, most provide insight into changes in aspects of sensorimotor control that might bear on the physiological subsystems underlying this high-level integrated function.

### IV. EVIDENCE

This section begins with a summary of evidence obtained from observations of crew performance decrements during operational situations. While largely circumstantial and clearly multi-factorial (likely resulting from a combination of physiological, behavioral, environmental, and engineering factors), this evidence provides a basis for concerns regarding the operational impacts of sensorimotor adaptation to space flight, as well as justification for continued investigation into the relative roles of the various factors affecting crew performance.

Following the operational evidence section, summaries of evidence obtained from space flight scientific investigations are provided in separate sections for in-flight and post-flight evidence. Within each section, the levels of evidence should be clear from context and/or

references. Where necessary supporting studies from space flight analog environments (e.g., parabolic flight), space flight analog populations (e.g., vestibular deficient patients), or other critically relevant ground-based investigations are included.

## **1. EVIDENCE OBTAINED FROM SPACE FLIGHT OPERATIONS**

An accurate assessment of the risks posed by the impacts of physiological and psychological adaptations to space flight on control of vehicles and other complex systems must account for the potentially offsetting influences of training/recency and engineering aids to task performance. Thus, it behooves us to review performance data obtained from space flight crews engaged in true mission operations. Evidence of operational performance decrements during space flight missions has been obtained from several sources; however, to our knowledge no well-designed scientific studies have been performed on critical operational task performance, so interpretation is frequently confounded by small numbers of observations, inconsistent data collection techniques, and/or uncontrolled engineering and environmental factors. Much of the relevant, extant operational data has been previously inaccessible to (or uninterpretable by) life sciences researchers. Recent programmatic changes have putatively improved access to both data and experts to help with interpretation.

### **1.1. Crew Verbal Reports**

A number of (unpublished) crew verbal reports were obtained early after flight by some of the authors of this review. While difficult to combine, owing at least in part to the lack of standardized questions and structured interview techniques, these reports are informative in that they provide insight into the individual crewmember perceptions. As an example, the following transcript obtained by Dr. Reschke captures impressions from a Shuttle commander obtained immediately (< 4 hours) after flight. The first part of the discussion focused on target acquisition tasks the commander performed for Dr. Reschke during the flight and his difficulties with nausea, disorientation, posture, locomotion, etc. after the flight (*italicized text* indicates the crewmember's responses to the Dr. Reschke's questions).

Did you try to limit your head movements? *Oh yes, definitely.* When you were trying to acquire the targets only...did you notice any difficulty in spotting the targets? *Oh yeah, oh yeah.* Did it seem as though the target was moving or was it you? *I felt that it was me. I just couldn't get my head to stop when I wanted it to.* So it was a head control problem? *Yeah, yeah in addition to the discomfort problem it caused.* So when you first got out of your seat today, can you describe what that felt like? *Oh gosh, I felt so heavy, and, uh, if I even got slightly off axis, you know leaned to the right or to the left like this, I felt like everything was starting to tumble.* When you came down the stairs did you feel unstable? *Oh yeah, I had somebody hold onto my arm.* Did you feel like your legs had muscle weakness, or ... was it mainly in your head? *It was mainly in my head.*

Every crewmember interviewed on landing day (>200 crewmembers) has reported some degree of disorientation/perceptual illusion, often accompanied by nausea (or other symptoms of motion sickness), and frequently accompanied by malcoordination, particularly during

locomotion. Of particular relevance to the ability to perform landing tasks, common tilt-translation illusions (see below) include an overestimation of tilt magnitude or misperception of the type of motion. Most also reported having experienced similar symptoms early in flight. However, except in the most severely affected, there seems to be no correlation between the severity of the symptoms following ascent and those following descent. The severity and persistence of post-flight symptoms varies widely among crewmembers, but both tend to decrease with increasing numbers of space flight missions. Both severity and persistence increase with mission duration. Symptoms generally subsided within hours to days following 1-2 week Shuttle missions but persisted for a week or more following 3-6 month Mir Station and International Space Station (ISS) missions.

The degree to which these psychophysical effects might affect piloting skills is difficult to judge, as recent (relative to launch), intensive training may have offset any impact on Shuttle landings, especially under nominal engineering and environmental conditions, and long duration Mir and ISS crewmembers to date have only piloted ballistic entry spacecraft, which parachute in, allowing no human control inputs during the last 15 min before landing.

The second part of the discussion focus on the perceived ability of the crewmembers to egress the vehicle immediately after landing, in case they could not be assisted by the ground team due to an emergency. (PLT=pilot, MS=mission specialist, CMD=commander, PS=payload specialist) The crewmembers' response to Dr. Reschke's question: "***Could you have performed an emergency egress at wheels stop?***" are shown in italic.

**3908** (11m-PLT) *Yes. I don't really know how the disorientation would have affected me...coming down the ladder and getting out of the front seat there's a lot of turning. I think if I would have gone slow, I could have done it.*

**3227** (2s-PLT) *No. It would have been difficult. I could have gotten out, but not in any hurry. No way. [What if you had to go out from the top?] I don't know if I could have made it. I could try.*

**4171** (3s-MS) *Yes.*

**3745** (13m-PLT) *Yes. It's a very personal thing. If I moved slowly and deliberately I could have gotten out. The fast motions cause you to crash over.*

**6064** (2m-PLT) *I'm not sure. It would have been a real workout. We had our strength, and we would have banged around, and we would have been slower, but it's a continuum, not yes or no, it depends. [What about though the hatch on top?] I think so, it would have been a real chore. Very, very demanding. Somebody ought to try it. And sooner than 40 minutes.*

**3029** (1m-MS) *No. Post-landing emergency egress is questionable. If you've got to get out in a hurry-- know I would have trouble. I tried to stand up, and immediately sat back down again. As I stood up, the lockers translated down, it didn't feel right. If I had to get out of the Orbiter during the first 5 or 10 minutes it would have been tough. It*

wouldn't have been pretty. If I had to do anything that required any real coordination... **[What about through the hatch on top?]** Very difficult. No way. People are fooling themselves, the whole business of throwing a rope out and lowering down the side--no way. Some guys jump up and are ready to go. Some people more readily adapt.

**5697** (16m-MS) Yes.

**2058** (SC) No. I was unsteady, had ataxic gait, couldn't correct for my mistakes.

**2042** (15m-CMD) No. I think we would have major problems. **[What about coming out of the top?]** Impossible.

**6855** (1s - MS) No. I think there would be a pile of people in the hatch or at the bottom of the hatch. Vestibularly, it would worry me. Getting to the slide, I would have fallen at least once.

**6245** (12s - MS) Yes or No. I think if your life is threatened, you could run, but you're going to fall doing it. It would be one of these that you're going to get up and fall and get up and run. Sure, when there's a fire behind you, you're going to find your balance pretty fast. **[What about going down the slide?]** You would go flat on your face.

**5157** (13s - CMD) No. The pilot and commander would have a real problem getting out of their seat. We don't have anything to stand on. Can't stand up straight. Looked for help from Life Sciences--might be good to have something to pull myself out. The pilot and commander would have a hell of a time if it was life essential to get out.

**8555** (4s - PS) Yes or No. Depends on where you'd have to exit--out the hatch, ok. Out the top, it would be hard. Have to go slower.

**4393** (14m-CMD) Yes. It would have been slow, would not have been pretty. I would have tumbled down the chute, but I could have gotten out. **[What about going down the slide?]** It would have been tough, because of the suit.

**6453** (4m - PLT) Yes. Yeah, Yeah, Yeah I could have gotten out. **[What about going down the slide?]** Yeah.


**5624** (3m - MS) Yes. Yeah, hard to say, because if you were in an emergency, the adrenaline rush just sort of takes over. And it wasn't until I stopped being dizzy that I noticed I was really nauseated. So you tend to think, when you get done what you need to do and then your body has time to betray you.

**5599** (12m - MS) Yes or No. Probably couldn't have run, but I could have walked. Getting the slide out was my job. I could have done it not fast, but slow and methodically.

## 1.2. Shuttle Entry and Landing Spatial Disorientation

A complete description of the space shuttle landing process, as well as plots of all shuttle landings touchdown point, sink rate, crossrange position, altitude, groundspeed, pitch attitude, and equivalent airspeed are given in [Appendix A](#).

Despite intensive training for all Shuttle commanders and pilots, some Shuttle landings were outside of the desired performance specifications, perhaps, in part, because of spatial disorientation. Shuttle entry and landing spatial disorientation (SD) differs from aviation SD, at least in terms of prevalence. Most instrument rated aircraft pilots have experienced SD, but episodes occur relatively infrequently in ordinary flying. In contrast, stimuli capable of producing SD were present during every Shuttle landing. At issue is whether the astronaut commander could successfully fly through the SD. Tilt-translation illusions and other sensorimotor disturbances (see below) did not occur in astronauts practicing approaches in the Shuttle Training Aircraft (STA), so their first actual experiences with these illusions occurred during their first actual return from space. Crews were forewarned about them, but they did not know how to predict the direction and magnitude of the effect, so a first-time flier did not know in advance which way to compensate. This was generally handled operationally by requiring commanders to have previous space flight experience (as pilots). Fortunately, for all Shuttle flights flown, there were no accidents specifically attributed to SD. However, several lines of circumstantial evidence suggest that the margin for error may have been less than generally recognized.



*Figure 1. Cumulative distribution functions allowing comparison between landing performances (vertical velocity at touchdown) before flight in the Shuttle Training Aircraft (STA) and those at the end of mission in the Space Shuttle (STS). (Paloski et al. 2008).*

All Shuttle landings were successful, but landing performance was more variable than desired. The timing and shape of the commander's control input during the flare depended critically on correct perception of speed, altitude, attitude, and sink rate. The flare maneuver, in turn, determines the readily measurable landing performance metrics, such as touchdown sink

rate, speed, and distance (Ashkenas *et al.* 1983). Of all the landings between STS-1 and STS-108, the Shuttle crossed the runway threshold abnormally low 20 times. Seven landings touched down abnormally long or short, and 13 had high touchdown sink rates, with three exceeding the 5 ft/sec structural limit. Moore *et al.* (2008a) reported that touchdown speeds during the first 100 Shuttle landings varied widely, with 20% outside of acceptable limits and six equaling or exceeding the maximum speed of 217 knots/hr (main landing gear tires are rated at 225 knots/hr maximum speed). Paloski *et al.* (2008) also noted that the fastest landing on record (224 knots/hr) was linked to the commander's momentary spatial disorientation, as was the second fastest (220 knots/hr). Normally, commanders perform better than this when flying the STA and the flight simulators. A different analysis of Shuttle landing compared piloting performances in terms of sink rate at touchdown. Figure 1 shows preflight performances flying the STA and subsequent post-flight performances in the Shuttle by commanders of all missions from STS-43 to STS-108. The average STA and STS touchdown sink rates were similar, and almost all STA touchdown sink rates fell in the desirable range; however, the STS touchdown sink rate distribution exhibits greater variability, with more than 10% exceeding the desired sink rate at touchdown.

Of particular note was the landing of the eight-day STS-3 mission in 1982. The commander, who was flying visually, took over manual control of the vehicle 30 seconds before landing at White Sands, NM. The vehicle was decelerating at 0.25 g. Starting at flare, when the commander attempted to lower the nose of the Shuttle, the vehicle exhibited a pilot induced oscillation (PIO) of three full cycles with increasing amplitude that continued through touchdown. Post-flight analysis showed no engineering anomaly in the flight control system. The commander was a highly experienced test pilot, very familiar with conventional PIO and with the 0.25 g deceleration of landing. However, it is possible that he under-perceived his pitch attitude because of tilt-translation ambiguities and caused the PIO by making larger control stick movements than necessary to compensate for the misperception. This could have been further exacerbated by inappropriate manual control inputs to the stick caused by miscalibration of eye-hand coordination. In a follow-up interview, however, the commander denied having any issue with PIO, or misinterpreting pitch attitude. His recollection was that the nose came down earlier than expected as the Shuttle began to slow down. He said the stick was not responsive when he first attempted to pitch the nose back up, but then it seemed to over-respond and pitched up more than he expected. Because he was then concerned about a potential problem with the stick, he brought the nose down and left it down. The commander's recollection appears to be consistent with the landing video, but not with data from the control stick that showed five large amplitude reversals in the pitch plane command after main gear touchdown. While difficult to reconstruct so long after the event, this may be noteworthy as an unrecognized case of spatial disorientation in a highly experienced pilot.

Increasing pilot awareness of the PIO problem, modifying software to reduce control authority automatically when oscillatory control outputs were detected, adding a heads-up display (HUD) pitch attitude read-out, and restricting landings to low cross-wind and good visibility conditions probably prevented PIO recurrence. However, it is clear that control phase and gain margins during the landing maneuver were routinely near limits of stability, and that

pilots making their first Shuttle landing must overcome disorienting perceptions not encountered during preflight training in the STA.

Flight surgeons examined returning Shuttle crewmembers for evidence of neurological dysfunction within several hours of landing. Crewmembers were scored for subjective symptoms, coordination, and functional motor performance. [McCluskey \*et al.\* \(2001\)](#) analyzed data from nine missions, and noted trends, such as a correlation between touch down sink rate and post-flight difficulty performing a sit-to-stand maneuver without using the arms. Scores indicating neurovestibular dysfunction generally correlated with poorer flying performances, including a lower approach and landing shorter, faster, and harder.

### **1.3. Apollo Lunar Landing Spatial Disorientation**

A review of all Apollo lunar landings performance, visibility, and surface awareness is provided in [Appendix B](#).

The Apollo Lunar Module (LM) had a digital autopilot that on later missions was capable of fully automatic landings. While the Apollo crews used the autopilot through most of the descent, all elected to fly the landing phase manually, using angular rate and linear velocity control sticks to adjust the vehicle trajectory while visually selecting the landing point. Landing sites and times were chosen so that the sun angle provided good visibility, but the crews had problems recognizing landmarks and estimating distances because of ambiguities in the size of terrain features. The vehicles had no electronic map or landing profile displays. The commander flew visually, designating the landing spot using a window reticle, while the second astronaut verbally annunciated vehicle states and status. Unfortunately, the landing area was generally not visible to the crew until the LM pitched to nearly upright at an altitude of about 7000 feet and distance of about 5 miles from touchdown with only 1-2 minutes of fuel remaining. Spatial disorientation was a concern during landing because visibility was reduced by the window design (views downward and to the right were blocked) and by lunar dust blowback that impaired surface and attitude visibility. For example, the Apollo 11 and 12 crews reported difficulty in nulling horizontal rates during landing because of blowing dust, and the Apollo 12 and 15 crews reported virtually no outside visibility in the final moments of landing. Visibility was improved in later missions by new hovering maneuvering procedures that reduced blowing dust.

Horizontal linear accelerations could not be avoided during the gradual descent to the landing zone or during hover maneuvers just before touchdown. Since lunar gravity is only 1/6 that on Earth, lunar landers had to pitch or roll through angles six times larger than on Earth to achieve a given horizontal acceleration using the engine thrust vector. The directional changes in gravito-inertial force these tilts created would have been larger than those on Earth, arguably making tilt-translation ambiguity illusions more likely. The Apollo crews trained for their missions in a 1/6 g Lunar Landing Training Vehicle, which did not simulate the vestibular effects of 1/6 g. Prior to their missions the only 1/6 g vestibular stimulation they received was during limited parabolic flight training. The Apollo crews did not acknowledge any spatial disorientation events during landing. They did later admit feeling a little “wobbly” when they

emerged to walk onto the lunar surface, but reported that coordination improved steadily during first few hours of lunar ambulation.

#### 1.4. Apollo Landing Geographic Disorientation

The Apollo LM utilized inertial navigation, updated by occasional star sights, radar orbital data from Earth, and radar altimetry during descent. Nonetheless, there was uncertainty in the accuracy of their computed position as they descended into the landing zone. Since crews could not look straight down, the final approach trajectory to the landing area had to use low angles (16-25°) so crew could see ahead. Mission planners only knew the landing zone terrain to 10 m resolution, so the crews had to confirm visually the LM trajectory and then sight the computer's anticipated touchdown point using a front window reticle. Given the fractal nature of lunar craters, identification of surface features was challenging. Humans interpret surface shape from shading based on a "light comes from above" assumption. This can create a "Moon crater" illusion (Ramachandran 1988) in which distant concave features, such as lunar craters, can be perceived as convex objects, such as hills, when viewed looking "down sun." The crews had to choose a suitably flat landing area, as judged by surface albedo and the absence of shadows indicating small craters or fissures. Landings were planned with sun elevations of 5-23°, so shadows were of moderate length, and with the crew facing down sun at a slight angle, so that shadows would be visible. The human eye can resolve 1.5 ft detail at a distance of about 4000 ft. As more surface details became visible, the commander typically redesignated the landing point (often several times), and eventually took over and flew manually, usually to a point somewhat beyond the final computer redesignated spot. He judged horizontal velocity looking out the window or using a cockpit Doppler radar display, and he used the LM shadow as a gauge, while listening to callouts of altitude, altitude rate, horizontal velocities, and fuel status. Since surface slope is impossible to judge visually looking straight down, the commander chose the final landing spot looking horizontally, and then flew over it and began final descent. At 50-100 feet, dust often obscured the outside view, and the vertical descent to touchdown sometimes had to be made relying primarily on instruments. The descent engine was cut off just before touchdown, to avoid explosion or damage should it contact the surface. The landing gear design assumed a maximum surface elevation difference of two feet within the landing gear footprint, and a maximum 12° terrain slope (Rogers 1972). Finding a flat landing spot was highly desirable, since vehicle tilts on the surface complicated surface operations and subsequent takeoff.

All six Apollo landings were ultimately successful. However, the Apollo 15 crew experienced geographic disorientation. When they pitched over, they could not identify the craters they were expecting, and the commander had to choose a landing spot in an unplanned area. Maintaining full awareness of the terrain immediately beneath the lander was usually impossible during the final phase of landing, and in one case the LM engine was damaged on touchdown (Jones & Glover 2014; Mindell 2008). The Apollo 12 commander encountered heavy dust blowback and said, *"I couldn't tell what was underneath me. I knew it was a generally good area and I was just going to have to bite the bullet and land, because I couldn't tell whether there was a crater down there or not."* He later added, *"It turned out there were more craters there than we realized, either because we didn't look before the dust started or because the dust obscured them."* The following mission, Apollo 14, landed safely, but on a 7° slope. Apollo

15 experienced severe dust blowback that contributed to making the hardest landing of the program (6.8 ft/sec), with the vehicle straddling the rim of a 5 ft deep crater, buckling the bell of the descent engine, and causing an 8° vehicle tilt. Apollo 16 and 17 experienced less dust obscuration and landed closer to level.

It seems likely that similar problems will be encountered when crews land vehicles on Mars. Improved navigation aids could help to avoid geographic disorientation, and increased reliance on auto-land capabilities could help maintain the landing performance within equipment specifications. However, improved training techniques, including realistic simulation of visual-vestibular inputs, will likely be required should commanders choose to use manual landing modes. The challenge of manual landing is likely to be much greater for Mars landings, owing primarily to the increased transit time in microgravity. A combination of more profound adaptation to microgravity and decreased training recency will likely increase substantially the risks associated with manual landing on Mars. [Note that using continuous artificial gravity, created by rotating all or part of the vehicle during transit, might well mitigate this risk (as well as many of the other biomedical risks), but the impact of prolonged exposure to a rotating environment on piloting a spacecraft would need to be investigated before committing to such a solution.]

### **1.5. Rendezvous and Docking**

A top priority in the U.S. space program is assuring crew and vehicle safety. This priority gained significant focus in June 1997 following the collision of the Progress 234 resupply ship with the Mir space station during a manual docking practice session. There were two separate attempts to dock the Progress with the Mir that day. In the first attempt, docking was aborted after the radar used for range calculations apparently interfered with a camera view of the Progress. In the second, near fatal attempt, mission managers decided to turn the radar off and leave the camera on. For this arrangement to work the Mir commander asked his two crewmates to look for the Progress approach through a porthole, and once sighted, to provide range information with handheld range instruments. Trouble began when neither the camera view nor the visual spotters could locate the Progress as it closed on the station. When spotters moved between modules to obtain a better view, they lost their frame of reference, and were uncertain which direction to look. Once spotted, the Progress' speed was above an acceptable rate, and it was very close to the Mir. Braking rockets on the Progress, fired by the Mir commander, failed to slow the velocity of the approaching spacecraft. No range information or other position data were available to assist the commander. To complicate matters, one of the other crewmembers may have bumped into the commander as he attempted to make last second inputs to the approaching Progress via joystick. The resulting collision tore a portion of the solar panel on the Mir, punched a hole in the Spektr module, and caused a decompression of the station.

Loss of situational awareness, spatial disorientation, and sensorimotor problems, including difficulties with vision, head-hand-eye coordination, and an inability to judge distance and velocity with limited feedback likely contributed to this outcome. Target acquisition studies have shown dramatic changes in the speed at which target visualization can be achieved,

delaying response time by as much as a 1000 ms (Kolev & Reschke 2014). Eye-hand response could take as long as another full second. A delay of two seconds is a lifetime when a spacecraft is closing, and not responding to joystick commands intended to decrease forward velocity. Members of the Russian Institute of Biomedical Problems (IBMP) believe that the collision between Mir and Progress was caused by poor situational awareness, spatial disorientation, and sensorimotor problems (I.B. Kozlovskaya, personal communication). After the fact, Ellis (2000) performed a rigorous, quantitative analysis of the available visual and non-visual information and suggested a number of potential sensorimotor and cognitive/psychophysical contributions to the crash. To avoid human factors contributions to future crashes such rigorous analyses should be performed well before attempting any three-dimensional visual-motor control task.

## 1.6. Teleoperator Tasks

The International Space Station (ISS) teleoperation system was heavily used during construction, and it will continue to be used to support extravehicular activities (EVA) operations, as well as in grappling/docking of rendezvousing cargo vehicles (Ruttley *et al.* 2010). Training and operating the Shuttle and ISS telerobotic manipulator systems as well as telerobotically controlled surface rovers presents significant sensorimotor challenges (Currie & Peacock 2001; Lathan & Tracey 2002; Menchaca-Brandan *et al.* 2007). These systems are usually controlled using separate rotational and translational hand controllers, requiring bimanual coordination skills and the ability to plan trajectories and control the arm in some combination of end-effector or world reference frames. The abilities to visualize and anticipate the three-dimensional position, motion, clearance, and mechanical singularities of the arm and moving base are critical. Thus, operators must have the cognitive abilities to integrate visual spatial information from several different reference frames.

Often the video cameras are not ideally placed, and in some situations (e.g. ISS operations) the views may actually be inverted with respect to one another, so cognitive mental rotation and perspective taking skills are also important (Lathan & Tracey 2002; Menchaca-Brandan *et al.* 2007). Teleoperation is sufficiently difficult that several hundred hours of training are required to qualify, and all operations are monitored by a second qualified operator, backed up by a team of trainers and engineers on the ground. Recency is important, so ISS astronauts perform on-orbit refresher training.

Despite all the training and precautions, however, there have been four-five significant ISS teleoperation incidents (e.g., collisions with a payload bay door, significant violations, or close calls) over the course of the first 16 ISS increments (Williamson 2007). A review of these incidents is given in Appendix C. Procedures are updated after each incident, but there are generic common factors relating to spatial visualization skills, misperception of camera views, timeline pressures, and fatigue.

## 1.7. Driving Performance

Driving a vehicle is one of the most complex sensorimotor/cognitive tasks attempted by most humans, and driving performance is known to be impaired in vestibular patients (Cohen *et*

al. 2003). Page & Gresty (1985) reported that vestibular patients experience difficulty in driving cars, primarily on open, featureless roads or when cresting hills, and MacDougall & Moore (2005a) reported that the vertical vestibulo-ocular reflex contributes significantly to maintaining dynamic visual acuity while driving. Adaptive changes in sensorimotor function during space flight can compromise a crewmember's ability to optimize multi-sensory integration, leading to perceptual illusions that further compromise the ability to drive under challenging conditions. During the June 2006 Apollo Medical Operations Summit in Houston, TX, Apollo crewmembers reported that rover operations posed the greatest risk for injury among lunar surface EVA activities. During rover operations, crewmembers often misperceived the angles of sloped terrain, and the bouncing from craters at times caused a feeling of nearly overturning while traveling cross-slope, causing the crewmembers to reduce their rover speed as a result (Apollo 15 report). This is not surprising given the evidence of tilt-translation disturbances following G-transitions, as incorrect perceptions of vehicle accelerations, tilted terrain, and uneven (bumpy) surfaces may cause inappropriate responsive actions. While automatic control systems can compensate for some deficiencies in performance, lessons learned from the Apollo missions (Mindell 2008) suggest that manual takeover is required as a minimum safe guard, and therefore countermeasures must concentrate on mitigating risks associated with crewmembers in the control loop for rover operations.

### **1.8. Implications for Exploration Class Vehicle Design**

There are specific spatial disorientation issues to address with the exploration class vehicle (ECV) currently being designed for planetary exploration missions. As a capsule, this vehicle will differ from the Shuttle in opportunities to induce disorientation. In the ECV, crews will stow their seats after ascent, so there will be no up/down cues except for the cockpit panels and the windows. ECV, like Shuttle, will probably be manually docked when crewed, while the logistics (cargo) version may have auto-docking. The crew's ability to remain visually oriented with ISS during proximity operations is a concern (S. Robinson, Astronaut and member of Crew Exploration Vehicle Cockpit Team personal communication). The developers must undertake analyses to ensure that the integrated visual out-the-window, camera imagery, display information, and sensor data will be sufficient to perform the envisioned three-dimensional docking tasks reliably.

### **1.9. Walking on the Moon**

The Apollo crews trained for their missions in a 1/6 g Lunar Landing Training Vehicle, which did not simulate the vestibular effects of 1/6 g. Prior to their missions the only 1/6 g vestibular stimulation they received was during limited parabolic flight training. They did later admit feeling a little "wobbly" when they emerged to walk onto the lunar surface, but reported that coordination improved steadily during first few hours of lunar ambulation.

During the 6 Apollo missions on the Moon, 12 crewmembers performed 14 EVAs that lasted a total of 78 hours. During these 14 EVAs, there were 23 falls and 11 saves. The causes and consequences of these falls obtained from video analysis and from crew reports are indicated in Table 1.

<i>Mission</i>	<i>Falls</i>	<i>Saves</i>	<i>Lunar</i>	<i>EVA</i> s		<i>Comments</i>
Apollo 11	0	0	21.6 h	1 EVA 2 h 31 min		* tendency to tip over on high jumps, but no problems overall
Apollo 12	2	0	31.5 h	EVA 1 4 h	EVA 2 3 h 50	* never fell down flat, and were able to roll over and push themselves up
				TOTAL: 7 h 50 min		
Apollo 14	0	1	33.5 h	EVA 1 4 h 47	EVA 2 4 hr 34	* grabbed MESA before falling; no balance or stability problems overall * one-sixth-g training helped them out
				TOTAL: 9 h 25 min		
Apollo 15	3	5	66.9 h	EVA 1 6 h 33 mins	EVA 2 7 h 12 min	* problems hopping down the ladder * tripped on soft soil while taking pictures and also tripped on rocks * worried about getting dirty to push themselves up * possible fall dismounting Rover * lost balance while throwing a used pallet * not worried about suit popping or splitting even though fell down a couple of times; more concerned about rocks and unevenness * fell climbing steep, soft rim of Station 6 crater * potential for falling while down on one knee
				EVA 3 4 h 50 min		
				TOTAL: 18 h 30 min		
Apollo 16	10	4	71 h	EVA 1 7 h 11	EVA 2 7 h 23	* loss of balance at Rover seat tugging at Velcro * problems retrieving objects from lunar surface (rock, bag, dust brush, penetrometer)
				EVA 3 5 h 40 min		* stumbled and ran forward to keep from falling
				TOTAL: 20 h 14 min		* landed heavily on PLSS during jumping turn
Apollo 17	8	1	75 h	EVA 1 7 h 12	EVA 2 7 h 37	* problems retrieving objects from lunar surface (tool and rock) * fall with discus-like throw of bag * problems removing deep core with jack * fall getting on Rover and running across a slope
				EVA 3 7 h 15 min		
				TOTAL: 22 h 4 min		

*Table 1. List of falls and close calls (saves) during the EVAs on the lunar surface.*

Frame by frame motion analyses were conducted on the films of the falls and near-falls during the Apollo-15 and Apollo-16 lunar EVAs to identify the specific reasons for their occurrence (Kubis et al. 1972a, 1972b). The in-flight voice transcripts were also used with the films to establish the events that occurred before and after the actual fall. Their research concluded that most of the falls were due to feet slipping on the loose soil and the crewmembers being in an unstable position to begin with. More specifically, the possible causal factors for the falls or near falls were the following:

- (a) Slipping, tripping, or temporarily losing balance occurred in certain terrain conditions, such as minor depressions, frequently associated with loose soil at the edges; sloping terrain with soil of varying density; scattered rocks of various sizes; reflectivity; and other changing or unexpected EVA conditions.
- (b) The crewmembers were unable to recognize the unexpected surface characteristics (see above) because of limited visibility due to the angle, position and glare of the sun; an inaccuracy in judgment of distance and slope; or an obstruction of their visual field by their helmet visor or the objects they were carrying out.

- (c) The automatic postural reactions that occurred in response to tripping or stumbling occurred too fast in lunar gravity, so these reactions threw them off even more. Frictional forces between the boot and the lunar surface are 1/6 of what they are on Earth. Thus, when tripping on a rock, the low frictional force between boot and surface caused slipping much sooner, and the postural reactions (which presumably used strategies developed in Earth gravity) did not serve to correct the disequilibrium (Kubis *et al.* 1972a).
- (d) The limited mobility of the suit was such that insufficient joint flexing could be achieved to pick up objects (e.g. hammer) on the lunar surface. This restraint forced the crewmembers to extend their hand beyond their knee and in doing so to lean forward approximately 75° from vertical. This placed the crewmembers in an unstable position and when they started to bounce back up, their feet slid and they fell (Kubis *et al.* 1972b).

## 2. EVIDENCE OBTAINED FROM SCIENTIFIC INVESTIGATIONS IN FLIGHT

Both in-flight and post-flight sensorimotor responses are related to the length of time spent in a weightless environment (Lackner & DiZio 1993). During the flight, a number of acute and chronic sensorimotor events may occur, including: (a) space motion sickness; (b) spatial disorientation; (c) difficulty with acquiring and tracking visual targets; (d) vision changes due perhaps to reshaping of the eye; (e) modification of proprioception (e.g., pointing, knowledge of limb position); (f) modifications in vestibulo-spinal reflexes; and (g) loss of muscle tissue and motor efference (Lackner and DiZio, 2000). Some of these events may be due to changes in otoconial structure with associated modification of supporting neural tissue, as observed in animal studies.

Return to the Earth can also result in neurovestibular symptoms for some crewmembers. Limitations in performance can then be caused by: (a) post-flight motion sickness lasting from a few hours to more than a week; (b) hypo- or hypertonia of the major postural skeletal muscles which alter locomotion; (c) early muscle fatigability and potentiation of major postural muscle reflexes which lead to falls or other accidents; (d) frequent under- or overshooting when reaching for an object; (e) unilateral gaze nystagmus, which is associated with dizziness and vertigo; (f) saccadic intrusion during smooth pursuit tracking; and (g) postural ataxia, including an inability to perform a simple heel-to-toe tandem walk without falling.

The sections below review the evidence collected during on orbit and pre/post flight investigations.

## 2.1. Decrements in Sensation

### 2.1.1. Visual Acuity

Testing of Mercury, Gemini, and Apollo astronauts revealed few significant changes in visual function with the exception of the following: constriction of the visual field, changes in intraocular tension, and changes in the caliber of retinal vasculature. Some constriction of the visual field was noted post-flight as well as a decrease in unaided seven-meter visual acuity, although the latter was not statistically significant. Post-flight decrease in intraocular pressure was significant and returned to preflight levels more slowly than expected. Retinal photography revealed no lesions but did show a decreased size of retinal vessels (Clément & Reschke 2008; Clément 2011).

Anecdotal reports from early Shuttle crewmembers describing decreases in visual performance, such as difficulty in reading checklists and unstable focus in the cabin, led to additional ophthalmologic testing (Task & Genco 1987). In 1989, NASA incorporated a questionnaire into the post-flight eye examination, regarding subjective reports of in-flight visual changes during short- and long-duration space flight. Astronauts were queried as to whether they perceived a subjective improvement or degradation in distant or near vision. This post-flight survey of approximately 450 astronauts (including several who had one or more previous flights) documented that visual changes were commonly observed, particularly during long-duration space flight, with degradation in near vision being significantly more prevalent (Figure 2).

These near vision changes were noted to be hyperopic (farsighted) in nature and more clinically apparent in older astronauts with decreased lens accommodation (Ginsberg & Vanderploeg 1987). In response to documented anecdotal reports of changes in near vision during space flight, astronauts over the age of 40 are prescribed “Space Anticipation Glasses” in the event that they experience a hyperopic shift during the mission. This visual shift appears to occur gradually, is variable in magnitude, and may persist for years following return to Earth. The origin of the visual changes during space flight is thought to be posterior globe flattening and choroidal engorgement brought about by cephalad fluid shifts leading to a forward displacement of the retina. This shortening of the distance between the retina and the lens may account for the hyperopic shift in microgravity (Mader *et al.* 2011, 2013).




Figure 2. Anecdotal ocular findings among Space Shuttle (left) and ISS crewmembers (right) from post-flight questionnaire. (DVA=distance visual acuity; NVA=near visual acuity; Dec=decrease; Inc=increase)

### 2.1.2. Proprioception

Microgravity modifies the stimuli associated with proprioception and thus affects the astronauts' knowledge of position of their limb (Schmitt & Reid 1985; Reschke *et al.* 1986; Kornilova 1997). Altered proprioception has been observed to cause erroneous discrimination of mass while accelerating a ball up and down using whole-arm movements (Ross *et al.* 1984). Mass discrimination improves, however, with arm movements of higher acceleration, indication that the in-flight impairment is partly due to a reduction in the z-axis pressure stimulation that provides information about weight on Earth (Ross *et al.* 1986a, 1986b).

Two theories exist to explain the observed decrements in proprioceptive function: (a) a physical degradation in the proprioceptive sensory system either via neural fiber degeneration, muscle atrophy, fluid shift, or the sudden and prolonged release of a constant muscle tone; (b) a disturbance of an external or internal-based spatial map. The results of pointing experiments suggest that gravity is important for the maintenance of a stable external spatial map (Young *et al.* 1993; Berger *et al.* 1997; Watt 1997). An egocentric reference system may also be used to assess limb position and maintain a sense of verticality (Kurtzer *et al.* 2005; Lackner & DiZio 2005b). This was observed aboard Mir when cosmonauts were able to draw ellipses in the air, either parallel or perpendicular to the longitudinal axis of the body, without visual aid. However, conflicting results have suggested there is a body scheme disturbance in the absence of gravity (Berger *et al.* 1992).

Proprioceptive illusions generated by vibrating leg muscles are different in-orbit compared to preflight. The perceptual effect of vibrating leg muscles was investigated in two subjects restrained on a back support. Before flight, leg vibration induced a backward and forward tilt sensation depending on which muscle was vibrated. After 20 to 21 days in orbit, the same vibrations caused a different whole body sensation. When the back support was used to replicate the axial force of normal gravity, the preflight illusions of forward and backward tilt returned (Roll *et al.* 1993). The results from a similar setup in parabolic flight demonstrated that

illusory movements were diminished in weightlessness but increased in hypergravity relative to 1 g. The authors concluded that the receptor output per unit stretch of a muscle spindle is influenced by variations in gravitational strength suggesting that unloading the otoliths in freefall decreases their descending modulation of alpha and beta motor neurons. This results in decreased tonic vibration reflexes (Lackner & DiZio 1992).

Proprioceptive illusions that involve self- or surround-motion have been observed for several muscle groups. The type and magnitude of illusion is influenced by the duration of space flight, the use of tactile cues, and whether the subject is restrained or free-floating. When crewmembers, wearing a special harness, were dropped from a quick-release hook, the sensation was one of the floor coming up rather than the subject falling. Most subjects reported that the floor suddenly slapped them on the feet (Reschke *et al.* 1984). The stretch reflex resulting from this landing propelled the subjects higher in the air than the height from which they were originally dropped.

However, there was a large variability in individual perception styles. Some subjects felt anchored by tactile cues from bungee cord-induced foot pressure, while others felt a tendency toward self-rotation under the same circumstances (Watt *et al.* 1992; Young *et al.* 1993). The CNS system that compares motor commands with sensory inputs may not be able to correctly distinguish between self-motion and movement of the environment because of sensory reinterpretation that occurs in space flight (Melvill-Jones 1974, Young *et al.* 1984; Parker *et al.* 1985; Reschke *et al.* 1986; Reschke & Parker 1987).

### **2.1.3. Motion Perception**

Some cosmonauts have reported sensations of linear acceleration to occur after engine cut-off (Bryanov *et al.* 1986; Kornilova *et al.* 1983, 1995). The change in threshold for detecting linear acceleration during flight, during passive body motion using a sled or another crewmember, has been hard to determine (Arrott *et al.* 1986; 1990; Benson & Vieville 1986; Young *et al.* 1993). Considerable variability was seen across crewmembers between the in-flight and pre-flight responses. Threshold sensitivity does seem to be axis- and gravity- dependent, but the difference in unloading the saccular and utricular otolithic membranes has yet to be fully addressed (Clément & Reschke 1996).

During passive angular acceleration in orbit, perception of the angle of rotation was overestimated during both roll and pitch movements (Clément *et al.* 1987). Yaw was not affected because, much like on Earth, the otoliths do not play a large role in yaw stimulation (Armstrong *et al.* 2015). Using a multi-axis rotating chair to assess subjective response to passive angular motion in orbit, results showed that pitch motion resulted in a perceived tumbling sensation. One subject reported that roll rotation felt like yaw motion (Benson *et al.* 1986). Without a constant gravitational vector or visual cues, the subject's reference to the environment is established using intrinsic coordinates. If no sensory map is available to establish intrinsic coordinates, then the axis of rotation or orientation relative to extrinsic coordinates cannot be determined (Wright *et al.* 2006; Harm *et al.* 2015).

Rotating-dome experiments conducted during Spacelab missions showed that most subjects experienced increased intensity in their visually induced sensation of motion (called vection) in microgravity, with some reporting complete rotatory (saturated) vection during flight (Young *et al.* 1986; Young & Shelhamer 1990). The onset of linear vertical vection in free-floating cosmonauts was earlier than in ground-based tests. Subjects also reported the illusion of bending forward and pitch, rather than linear vection (Mueller *et al.* 1993). Visual-induced illusions become stronger in space, perhaps because the otolith organs neither confirm nor deny body tilt in microgravity.

## **2.2. Decrements in Gaze Control**

Eye-head coordination is critical to performing piloting tasks, and it is very important to controlling other vehicles (e.g., rovers and automobiles) and complex systems (e.g., robotic arms and other remote manipulators). Rapidly locating and reading instrument displays, identifying suitable landing locations, free of craters, rocks, etc., and tracking the motion of targets and/or objects being manipulated are among the tasks requiring good vision enabled by optimized eye movement control (Harm *et al.* 2007). A large body of evidence demonstrates that the G-transitions associated with space flight disrupt oculomotor performance. Highlights are summarized in the following subsections.

### **2.2.1. Smooth Pursuit Eye Movements**

Smooth pursuit eye movements are produced during voluntary visual tracking of moving targets without head movements. Evidence suggests that the basic mechanisms underlying smooth pursuit tracking are modified by exposure to space flight. When crewmembers were asked to visually track a simple point stimulus at 0.33 Hz in either horizontal or vertical planes the amplitude of eye movement was reduced and the number of corrective saccades was increased (Kornilova *et al.* 1991a, 1991b; Reschke *et al.* 1999; Reschke *et al.* 1996). They crewmembers clearly undershot the target. The performance deterioration was the most pronounced for a point stimulus moving vertically or diagonally, early in-flight (flight day 3), late in flight (flight days 50, 116, and 164), and early after flight. Thus, it appears that the saccadic system must be utilized extensively to maintain accurate target tracking, and vision is degraded by an inability to maintain the target focused on the fovea (Hopp & Fuchs 2006).

### **2.2.2. Vestibulo-Ocular Reflex**

During head and/or body movements, the gaze stabilization system maintains high visual acuity by coordinating movement of the eyes and head so as to stabilize the image of interest on the fovea. The vestibulo-ocular reflex (VOR), a servo system that uses head motion signals sensed by the semicircular canals and otolith organs to generate vision-stabilizing compensatory eye movements, is critical to this function. Blurred vision, oscillopsia (illusory movement of the visual world), and/or reduced dynamic visual acuity occur when this gaze compensation mechanism is disrupted. VOR function is plastic, meaning it can adapt to different environmental stimuli (Berthoz & Melvill Jones 1985). For example, the VOR gain (amount of eye rotation caused by a unit of head rotation) adapts when individuals begin wearing new prescription eyeglasses. A number of relevant flight experiments have

demonstrated that various VOR response properties are modified during and after space flight, and that the degree of adaptation varies among subjects and experimental conditions (reviewed Reschke *et al.* 1996).

No significant changes in yaw VOR gain were observed in response to voluntary (active) head oscillations at frequencies ranging from 0.25 to 1 Hz (Thornton *et al.* 1985, 1988, 1989; Watt *et al.* 1985; Viéville & Clément 1986; Benson & Viéville 1986). Unlike yaw plane head movements, pitch and roll plane head movements in normal gravity change the orientation of the head relative to the gravity vector, thereby modulating gravitational stimulation of the otolith organs (Wood *et al.* 2009). One might expect, therefore, that the pitch and roll plane VOR would be more affected by space flight than the yaw plane VOR. The pitch VOR response to voluntary head oscillations has been measured during and after space flight at frequencies comparable to those described above for yaw, and the results are inconclusive. Some studies have reported no changes in flight (Watt *et al.* 1985) while others found an increase in VOR gain and in the phase lag (delay between head motion and elicited eye motion) (Berthoz *et al.* 1986), and yet others reported a decrease in vertical VOR gain (Viéville *et al.* 1986; Clarke *et al.* 1993). Clarke *et al.* (2000) also reported changes in torsional VOR during voluntary head movements in the roll plane during and after space flight.


These findings suggest that VOR is presumably disrupted early after insertion into orbit. Fortunately, central adaptive processes re-establish VOR response properties over time in the new environment, resulting in recovery of accurate stabilization of vision during head and/or body movements in the new environment. However, critical mission activities requiring accurate gaze stabilization during head movements (e.g., piloting/landing a spacecraft) will likely be performed less skillfully during or soon after G-transitions.

### **2.2.3. Eye-Head Coordination and Target Acquisition**

Gaze is the direction of the visual axis in three-dimensional space. It is defined as the sum of eye position with respect to the head and head position with respect to space. Acquisition of new visual targets of interest is generally accomplished using coordinated eye-head movements consisting of a saccadic eye movement that shifts gaze onto the target combined with a VOR response that maintains the target on the fovea as the head moves to its final position. Space flight modifies eye-head coordination during target acquisition (Kozlovskaya *et al.* 1985; Thornton *et al.* 1988; Tomilovskaya *et al.* 2011) and ocular saccadic performance (Uri *et al.* 1989; Andre-Deshays *et al.* 1993; Reschke *et al.* 1996, 1999).

Different strategies for gaze and target acquisition have been observed during and post-flight. Head movement towards a target near or beyond ( $\geq \pm 50^\circ$ ) the effective oculomotor (EOM) range is delayed, resulting in a VOR that tends to pull gaze off target (Reschke *et al.* 1999). Reduced angular positioning of the head forces larger compensatory eye saccades to direct the eye back to the target (Gresty & Leech 1977; Zangemeister & Stark 1981, 1982; Zangemeister *et al.* 1988, 1991) (Figure 3). The slower head movement contributes to a near doubling of the latency required to fixate peripheral targets. Sirota *et al.* (1987) showed that during adaptation to space, non-human primates trained to perform a visual target acquisition

task requiring accurate perception of peripheral targets showed delays in the onset of the gaze response and made significantly more errors in identifying the visual characteristics of the peripheral targets.



*Figure 3. Eye, head and gaze (sum of eye and head) position during target acquisition at the limits of the effective oculomotor range (50° in the horizontal plane). Note that on the second flight day the head rotation is delayed, the head moves at a much lower velocity than preflight, and the VOR is infused with several saccades to bring the eye on target.*

#### **2.2.4. Optokinetic Nystagmus**

Optokinetic eye movements that are produced in response to scene motion, whether it is induced by self-motion or surround-motion, offer a unique way to investigate the effects of space flight on the vestibular system, as the vestibular cells in the brainstem cannot themselves differentiate between visual scene movement and head movement relative to the environment. The asymmetry in optokinetic nystagmus (OKN) gain normally observed on Earth was reversed during the first 3 days of space flight and a downward drift was still evident on the fifth day, despite the gain and the eye movement field being nearly normal. A decrease in vertical OKN gain was also noted during flight, which increased shortly after flight, along with a restoration of asymmetry (Clément & Berthoz 1988, 1990). An experiment using short-radius centrifugation to elicit linear acceleration in orbit also indicated that the eye rotation axis during OKN tended to align with this linear acceleration (Moore *et al.* 2005a, 2005b). The reduced vertical asymmetry might reflect adaptation from a gravitational to an idiotropic reference frame in situations where vertical asymmetries have no functional use (Dai *et al.* 1994).

#### **2.3. Decrements in Eye-Hand Coordination Performance**

Eye-hand coordination skills are also critically important to performing piloting tasks and controlling other vehicles and complex systems. Reaching to switches on instrument panels, smoothly guiding the trajectory of a flight- or ground-based vehicle, and carefully positioning the end-effector of a robotic arm are some of the tasks requiring high levels of eye-hand coordination (Bortolami *et al.* 2008). While not studied as intensively as oculomotor performance, a number of studies of eye-hand coordination have been performed during space flight missions.

### **2.3.1. Control of Aimed Arm Movements**

When astronauts first encounter an altered gravity environment, arm movements are often inappropriate and inaccurate (Gazenko *et al.* 1981; Johnson *et al.* 1975; Nicogossian *et al.* 1989). During the Neurolab Space Shuttle mission (STS-90), Bock *et al.* (2003) performed an experiment in which subjects pointed, without seeing their hands, to targets located at fixed distances but varying directions from a common starting point. Using a video-based technique to measure finger position they found that the mean response amplitude was not significantly changed during flight, but that movement variability, reaction time, and duration were all significantly increased. After landing, they found a significant increase in mean response amplitude during the first post-flight session, but no change in variability or timing compared with preflight values. In separate experiments, Watt *et al.* (1985, 1997) reported reduced accuracy during space flight when subjects pointed to memorized targets. This effect was much greater when the hand could not be seen before each pointing trial. When subjects pointed at memorized locations with eyes closed, the variability of their responses was substantially higher during space flight than during control sessions on Earth. In other studies (Berger *et al.* 1997; Papaxanthis *et al.* 1998), the investigators found that when crewmembers on the Mir station pointed to targets with eyes open, variability and mean response amplitude remained normal, but the movement duration increased by 10 to 20% over the course of the mission (flight day 2-162).

### **2.3.2. Reaching and Grasping**

Thornton & Rummel (1977) showed that basic tasks such as reaching and grasping were significantly impaired during the Skylab missions. Later, Bock *et al.* (1996a, 1996b; Bock *et al.* 1992; Bock 1996; Bock & Cheung 1998; Hudson *et al.* 2005) investigated pointing, grasping, and isometric responses during brief episodes of changed gravity, produced by parabolic flights or centrifugation. These experiments provided converging evidence suggesting that during either reduced or increased gravity, the mean amplitude of responses is larger than in normal gravity, while response variability and duration remains unchanged. During the Neurolab Space Shuttle mission, Bock *et al.* (2003) found that the accuracy during flight of grasping luminous discs between their thumb and index fingers was unchanged from preflight values, but task performance was slower.

### **2.3.3. Manual Tracking**

Changes in the ability of crewmembers to move their arms along prescribed trajectories have also been studied in space. For example, Gurfinkel *et al.* (1993) found no differences in orientation or overall shape when crewmembers with eyes closed drew imagined ellipses oriented parallel or perpendicular to their long body axes. In another study, Lipshits *et al.* (1993) examined the ability of crewmembers to maintain a cursor in a stationary position in the presence of external disturbances. They found no performance decrements when the disturbances were easily predictable. However, in follow-on experiment using more complex disturbances, Manzey *et al.* (1995, 1998) found that tracking errors were increased early in

flight, but gradually normalized within 2-3 weeks of exposure to the space environment. Step-tracking performance accuracy was also affected only marginally during flight in another experiment (Sangals *et al.* 1999). However, kinematic analyses revealed a considerable change in the underlying movement dynamics: too-small force and, thus, too-low velocity in the first part of the movement was mainly compensated by lengthening the deceleration phase of the primary movement, so that accuracy was regained at its end.

In another experiment where subjects tracked with their unseen finger a target moving along a circle at 0.5, 0.75, or 1.25 Hz, subjects' response paths were found to be elliptical rather than circular (Bock *et al.* 2003). The variability of finger positions about the best-fitting ellipse was significantly higher than preflight during the first in-flight session, and responses lagged significantly behind the target during the highest target speed condition. Performance normalized later during flight, but deficits, albeit less pronounced, reappeared during the first two post-flight test sessions. It should be noted that response slowing and increased variability were limited to the first in-flight session for the tracking paradigm, but were most pronounced during later in-flight sessions for the pointing paradigm. The investigators interpreted these observations as indicating an underestimation of mass during flight (Bock *et al.* 2003).

#### **2.3.4. Force Discrimination and Control**

During a Mir station mission the ability of a cosmonaut to reproduce several positions of a handle from memory was tested. The accuracy with which the handle was set to a given position was reduced. However, the temporal parameters of the movement and the number of discernable handle positions did not change (Lipshits *et al.* 1993; Reschke *et al.* 1996).

#### **2.3.5. Fine Motor Control**

Campbell *et al.* (2005) evaluated the feasibility of survival surgery performed on rats during the Neurolab Shuttle mission. Craniotomy, leg dissection, thoracotomy, laminectomy, and laparotomy were performed as a part of physiological investigations. Surgical techniques successfully demonstrated in rats during space flight include general anesthesia, wound closure and healing, hemostasis, control of surgical fluids, operator restraint, and control of surgical instruments. Although the crew noted no decrement in manual dexterity, the operative time was longer compared with the ground experience due to the need to maintain restraint of surgical supplies and instruments. In another study, Rafiq *et al.* (2006) measured the effect of microgravity on fine motor skills by investigating basic surgical task performance during parabolic flight. They found that forces applied to the laparoscopic tool handles during knot tying were increased force while knot quality was decreased during flight compared with ground control sessions. Also, Panait *et al.* (2006) studied the performance of basic laparoscopic skills (clip application, grasping, cutting, and suturing) during parabolic microgravity flights. When compared with one gravity performance, they found that there was a significant increase in tissue injury and task erosion and a decreased trend in the number of tasks successfully completed.

### **2.3.6. Dual-Tasking and Manual Performance**

Studies have demonstrated the effects of dual tasking on manual control during space flight. [Manzey et al. \(1998\)](#) found impairments in tracking performance and time-sharing efficiency during the first month in space. These impairments included larger single- versus dual-task differences of both memory search speed and tracking error in orbit compared to preflight. [Bock et al. \(2010\)](#) also found more tracking error for dual-task conditions inflight compared to preflight. Importantly, the increase in tracking error persisted for the duration of the mission and did not recover until four days after the return to Earth. [Manzey et al. \(1995\)](#) examined dual task performance for a crewmember throughout an 8-day space mission. Specifically, unstable tracking with concurrent memory search were tested 13 times during the course of the mission. The results indicated impairments in the single task memory and dual task performance. Based on the results from these studies, psychomotor processes and higher attentional functions are impaired when in the space environment.

Laboratory tasks might underestimate the actual deficits since they differ from a real-life scenario in a number of ways. For example, the slowing of aimed arm movements was 10-30% in experimental tasks, but was up to 67% during routine activities on Skylab as analyzed using time and motion studies ([Kubis et al. 1977](#)). Degradation of performance may be exacerbated in part due to postural instability, which may not play a role when a pilot controls a landing while strapped into a seat, but may have a greater impact if landing is performed while standing like during the Apollo lunar landings.

## **2.4. Decrements in Spatial Orientation Perception**

Spatial disorientation has been one of the most frequently studied aspects of sensorimotor adaptation during space flight. Returning crewmembers report that the most overt physiological phenomena associated with space flight are inversion illusions at main engine cut off, occasional in-flight disorientation, early-mission motion sickness, and head-movement-contingent disorientation during entry and landing. These neuro-vestibular phenomena occur during and after G-level transitions, which, unfortunately, also correspond to mission phases where physical and cognitive performance are particularly critical for crew safety and mission success. Accurate perception of self-in-space motion and self-motion relative to other objects are critical to piloting, driving, and remote manipulator operations.

### **2.4.1. Spatial Disorientation in Aviation**

Spatial Disorientation (SD) is traditionally defined as a “failure to correctly perceive attitude, position, and motion of the aircraft” ([Benson & Barnes 1978](#)), as displayed on the aircraft’s primary attitude and flight control displays. A history of aviation SD research, a taxonomy of the classic SD illusions, an explanation of the underlying physiology, and data on incidence of SD related accidents have been well-reviewed by [Gillingham & Previc \(1996\)](#), [Young \(2003\)](#), and [Previc & Ercoline \(2004\)](#). Classic SD (e.g., somatogravic illusions, leans, G-excess illusions, inversion illusions, or Coriolis illusions) can result from unusual vestibular

stimulation or from sparse or misleading visual cues (false horizons, ambiguous size, or surface slant cues). SD has been further categorized as Type I (unrecognized), Type II (recognized), or Type III (incapacitating). Type 1 SD is the most insidious because the pilot is unaware the aircraft is in a dangerous state (Gillingham & Wolfe 1986).

Spatial disorientation remains an enduring problem in aviation. Surveys consistently indicate that a majority of pilots have experienced significant SD, many more than once. Pilots spend hours training, hoping that when SD occurs they will recognize it and be able to fly through it using their instruments. Incidence of SD depends on the type of flying and weather conditions. The likelihood that SD will create an accident goes up when flying at low altitude, since there is less time to recognize and recover. In US general aviation, SD is a factor in 15% of fatal accidents, a rate of one every 100,000 flying hours (National Transportation Safety Board [NTSB] 2007). Hence there is one fatal SD related accident approximately every week. In US scheduled airline flying, most of which is at high altitude and on autopilot, the SD rate is far lower, though, as discussed below, controlled flight into terrain during approach remains a problem. For similar reasons, in military flying, SD rates are higher in fighter/attack aircraft and helicopter operations than in military transport flying. In the US Air Force, the overall SD rate is 0.5 per 100,000 flying hours. SD was a factor in 14% of all major accidents, and insufficient or misleading visual cues contributed to 61% of these (Lyons *et al.* 1994; Holmes *et al.* 2003). Many military aircraft HUD displays automatically declutter to better support recognition and recovery of extreme attitudes. In US Army helicopter operations, most of which take place at low altitudes, the overall SD rate has been about 3 per 100,000 hours. However, the rate rises by a factor of five at night, and by 20 when night vision goggles are used (Braithwaite *et al.* 1998; Durnford *et al.* 1995; Curry *et al.* 2007). Lack of reliable visual references has been the most frequent cause of helicopter SD fatalities, typically due to rain, fog, or blowing snow or dust during landing.

There is evidence from flight simulator experiments that instrument flying experience and recency (within 2 weeks) helps pilots “fly through” disorienting transients created using galvanic stimulation of the vestibular system (Malcik 1968). However, applying such stimulation to pilots flying visual approaches can trigger pilot induced oscillations (MacDougall & Moore 2005a).

#### **2.4.2. Geographic Disorientation and Terrain Awareness**

A second class of accidents and incidents are caused by “geographic disorientation,” defined as the failure to recognize and/or maintain the desired position relative to the external ground and airspace environment (Antunano & Hernandez 1989). Common examples include becoming lost in the air or on the ground and then straying into prohibited airspace, landing at wrong airport, or taking off, landing, or intruding into an inappropriate runway, causing collisions or overruns.

Loss of terrain awareness is almost always a factor in Controlled Flight Into Terrain (CFIT) accidents, where the pilot unintentionally flies into terrain, usually during the approach-and-landing phase. Ground proximity warning systems (GPWS), mandated since the 1970s in civil

transport aircraft, have reduced the number of CFIT accidents, but they still account for a third of all fatal accidents in this sector over the past decade (Boeing 2007). CFIT can result from classic spatial disorientation, but more commonly results from loss of terrain awareness due to preoccupation with other tasks, or incorrectly setting and/or inappropriately trusting the autopilot. Electronic cockpit map displays highlighting nearby terrain (e.g. terrain awareness and warning system [TAWS]) and “synthetic vision” backgrounds for attitude indicators, depicting a virtual “out the window” view are expected to reduce CFIT incidence.

Because the classic definition of spatial disorientation does not include Terrain Awareness or Geographic Disorientation, CFIT and geographic disorientation accidents are typically only categorized by investigators as due to “Loss of Situational Awareness.” At the most general level, situation awareness is defined as perception of elements defining a mental model of the current situation, comprehension of their meaning, and projection of their future status (Endsley 1995). Obviously, there are many dimensions to situation awareness, including all elements of spatial knowledge, as well as awareness of traffic, weather, autopilot mode, fuel, system, and weapon status, etc. The more general situation awareness concept has proven useful in understanding many types of accidents, partly because it emphasizes the importance of “confirmation bias” in attention, perception, and decision-making. Classic spatial disorientation accidents should also be properly co-coded as loss of situation awareness accidents (Previc *et al.* 1995), but traditionally are not.

Overall, one concludes that large transport aircraft operations are relatively safe – less than one spatial disorientation related accident every 106 approaches. However, in other aviation segments, where more of the flight is conducted at low altitude, in bad weather, and/or using sensory aids, the fatal accident rate rises by an order of magnitude or more.

### **2.4.3. Spatial Disorientation During Space Flight**

The literature on spatial disorientation events during space flight has been well reviewed by Oman (2003, 2007). Numerous detailed firsthand accounts by astronauts and cosmonauts have also appeared (Cooper 1976; Burrough 1988; Linenger 2000; Lu 2003; Jones 2006; Hadfield 2013). Almost all crewmembers describe a transient somatogravic tumbling illusion or momentary inversion illusion upon reaching orbit, when main engine cutoff causes a rapid deceleration to constant orbital velocity. About 10% subsequently experience a sense of gravitational inversion (what they call “the downs”) that persists for 2 to 3 days after launch regardless of relative body orientation in the cabin, even with eyes closed. Persistent inversion illusions are thought to result from the combined somatosensory effects of headward fluid shift, and saccular otolith unweighting (Mittelstaedt 1986a, 1986b, 1986c; Oman *et al.* 1986; Graybiel & Kellogg 1967).

On Earth, the ability to perceive verticality is quite good. This ability is dependent on input from visual, vestibular and somatosensory systems, and on a functioning CNS (Bortolami *et al.* 2006a, 2006b; Bryan *et al.* 2007). Many reports also exist of astronauts’ perceptions of pitch-forward, pitch-up, or pitch-down attitudes once they enter weightlessness (Harm *et al.* 1994), giving rise to an altered subjective vertical. The perception of subjective visual vertical was

investigated in cosmonauts on short- and long-duration flights, either seated upright or rolled laterally (Bokhov *et al.* 1969; Bokhov *et al.* 1973; Kornilova & Kaspransky 1994). Immediately post-flight, all cosmonauts returning from long-duration flights and 38% of those returning from short-duration flights inaccurately perceived the subjective visual vertical. In the lateral position, the magnitude of error increased sharply and in the majority of cases altered attitude, relative to that observed preflight. In 10% of short-term and 18% of long-term mission cosmonauts, the previously occurring Aubert phenomenon gave way to the Müller phenomenon. The recovery of subjective vertical perception occurred 10-14 days post-flight. Interestingly, the accuracy of subjective perception of the body's vertical axis was virtually unchanged between pre- and post-flight testing.

Clément *et al.* (1987) examined in-flight adaptive changes in perception of subjective body orientation. They observed that when subjects' feet were held in Shuttle foot restraints, perception of subjective body orientation depended greatly on visual cues. Rotation around the ankle joint had little or no effect on correcting tilt angle in the absence of vision, which may be related to the lack of proprioceptive input during static limb positioning. The flexor tone produced in the dorsiflexor muscles has been proposed to help maintain a virtual vertical projection of the body's center of mass (Massion *et al.* 1997). In other words, the CNS tries to recreate a condition in weightlessness that is similar to that on Earth. This interpretation is in agreement with an internal representation of gravity (Mittelstaedt & Glasauer 1993; Clément *et al.* 2001), which allows a coherent mental representation of the body in alignment with the longitudinal axis. This internal model of gravity would also serve as a reference frame for movement.

#### **2.4.4. Visual Reorientation Illusion**

The visual reorientation illusion (VRI) has been first described by astronauts on the Skylab and Spacelab-1 missions. When crew float about in the cabin, they experience a spontaneous change in the subjective identity of surrounding surfaces, such that the "surface beneath my feet seems somehow like a floor." Oman (2003, 2007) noted that astronauts must orient with respect to a vehicle frame of reference defined by local visual vertical cues. However, architectural symmetries of the cabin interior typically define multiple "visual vertical" directions, usually separated by 90°. The Earth can provide yet another visual reference frame when viewed through cockpit windows or while spacewalking. There is a natural tendency to perceive the subjective vertical as being aligned with the head-foot axis, generally referred to as the "idiotropic" effect (Mittelstaedt & Mittelstaedt 1996). Which visual reference frame the observer adopts thus depends strongly on relative body orientation and visual attention. VRI occur when the perceived visual vertical reference frame is not aligned with the actual, so that, for example, the overhead surface is perceived as a deck.

The type and magnitude of perceptual illusions may depend on whether a crewmember uses an idiotropic or visual reference frame. Individuals who are visually oriented with respect to external references perceive themselves to be inverted or sideways during flight. They report difficulty in switching rest frames and performing coordinate transformations, in addition to experiencing loss of orientation in the absence of visual cues. When an idiotropic reference

frame is used, the alignment of a vertical along the longitudinal body axis allows little disorientation and an easy switch between rest frames. Forty-six percent of astronauts and 58% of cosmonauts were classified as using an idiotropic reference frame, 46% of astronauts and 34% of cosmonauts as using predominantly visual reference, and 8% of both crews were classified as using a mixed of both (Young *et al.* 1986; Friederici & Levelt 1990; Harm & Parker 1993). Individual experiences with self- or surround-motion may vary, but commonly reported illusions include (a) exaggerated rate, amplitude, and positioning of body movement; (b) temporal disturbance to perception of motion; and (c) altered path perception (Harm *et al.* 1999).

Data from animal experiments in parabolic and orbital flight (Taube *et al.* 2004; Oman 2007) suggest that the VRI surface identity illusion physiologically corresponds to a realignment of the two-dimensional plane that limbic neurons use to code direction and location (see section 2.4.6). When VRI occur, crews lose their sense of direction with respect to the entire vehicle, and reach or look in the wrong direction for remembered objects. Susceptibility to VRI continues through the first weeks in space, and occasional illusions have been reported after many months on orbit. Strong sensations of height vertigo have been described during spacewalks. These might reflect sudden changes in the limbic horizontal frame of reference from the spacecraft to the surface of the Earth.

VRI can also occur on Earth, but reorientations usually occur only in yaw perception about the gravitational axis, e.g., when we emerge from a subway and discover we are facing in an unexpected direction. VRI about Earth-horizontal axes have been created using tumbling rooms and virtual reality techniques (Harm *et al.* 2008). For example, investigators have shown that the direction and strength of visual vertical cues depend on field of view, the relative orientation of familiar gravitationally “polarized” objects, and the orientation and symmetry of surfaces in the visual background (Hu *et al.* 1999; Howard & Hu 2001, Jenkin *et al.* 2007). Single planar surfaces or the longer surface in a rectangular room interior were most frequently identified as “down.” (Oman 2007) Prior visual experience and knowledge of the specific environment are also important factors. Even when VRI do not occur, the visual verticals of adjacent or docked spacecraft modules are often incongruently aligned. Astronauts typically orient to the reference frame of the local module, and significant cognitive effort is required to sort out these multiple vehicle frames of reference. Using virtual reality simulations, Oman 2007 and Aoki *et al.* (2006, 2007) have shown that subjects remember the interiors of each module in a canonical, visually upright orientation. When performing tasks that require subjects to interrelate different reference frames, additional time is required and workload imposed. The fastest responses occur when module verticals are congruently aligned. Significantly greater time is required to perform simulated emergency egress navigation tasks when module visual vertical reference frames are incongruently aligned (Oman 2006).

#### **2.4.5. Tilt-Translation and Tilt-Gain Illusions**

Arguably the greatest space flight-related challenge to the human internal navigation system results from the ambiguities between tilt and translation stimuli. Albert Einstein (1908) was the first to postulate “the complete physical equivalence of a gravitational field and a

corresponding acceleration of the reference system.” According to his equivalence principle, linear accelerations resulting from translational motions are physically indistinguishable from linear accelerations resulting from tilts with respect to gravity because the forces are identical in nature. The ability of the central nervous system to resolve tilt-translation ambiguities is critical to providing the spatial orientation awareness essential for controlling activities in everyday life.

Two hypothetical mechanisms that have been proposed for resolving tilt-translation ambiguities are frequency segregation and multi-sensory integration. The frequency segregation hypothesis suggests that low frequency linear accelerations are interpreted as tilt and high frequency accelerations as translation (Mayne 1974; Merfeld *et al.* 2005a, 2005b). This hypothesis appears consistent in principal with the response dynamics of the different primary otolith afferents (Fernandez *et al.* 1972; Fernandez & Goldberg 1976; Peterson & Chen-Huang 2002), secondary processing of otolith input in the vestibular nuclei (Xerri *et al.* 1987; King *et al.* 1999), and also with natural behavior (Pozzo *et al.* 1990). The multi-sensory integration hypothesis, on the other hand, suggests that the brain must rely on information from other sensors, such as canals and vision, to correctly discriminate between tilt and translation (Guedry 1974; Angelaki *et al.* 1999). More specifically, it suggests that the brain learns to anticipate a sequence of sensory feedback patterns for any given movement. This hypothesis generally involves the use of internal models, or neural representations of physical parameters, and combines efferent and afferent information to resolve sensory ambiguity (Young 1974; Oman 1982; Droulez & Darlot 1990; Zupan *et al.* 2000; Green & Angelaki 2004; Poon & Merfeld 2005; Zupan & Merfeld 2005).

Although multi-sensory integration and frequency segregation are typically posed as competing hypotheses, they are not mutually exclusive. The segregation of otolith-ocular responses as a function of frequency has been clearly demonstrated (Paige *et al.* 1996). Yet one implication of frequency segregation is that there must be a mid-frequency crossover region where it is difficult to distinguish tilt from translation. Paige & Seidman (1999) reported that the crossover frequency is approximately 0.5 Hz in primates, and Wood (2002) suggested that it occurs at about 0.3 Hz in humans. Multi-sensory integration may play a critical role near the crossover frequency.

Among the factors that facilitate sensorimotor adaptation, active voluntary motion may be one of the most important (Welch 1986). Performing visual tasks with the intent to override vestibular input may also catalyze adaptation (Guedry 1964; Shelhamer *et al.* 2010, 2012). Most sensory conflict theories related to sensorimotor adaptation have been derived from the concept of “efference copy”, which states that there are predicted sensory feedbacks for any given motor action (Von Holst & Mittelstaedt 1973; Reason 1978). Head movement kinematics on Earth yield invariant unique patterns of canal and otolith signals irrespective of other sensors (Guedry *et al.* 1998). During adaptation to altered gravito-inertial environments, though, new patterns of sensory feedback must become associated with head movements to reduce sensory conflict. The observation that some astronauts tend to restrict head-on-trunk movements on orbit, preferring to rotate more from the waist than the neck, reflects an

adaptive change in motor strategy that might further contribute to motion sickness (Watt 1987) and post-flight postural and dysfunction (Bloomberg *et al.* 1997).

Two separate human experiments conducted on orbit by Reschke *et al.* (1988) and Clément *et al.* (2001) investigated the effects of sustained linear accelerations during eccentric rotation created by short-radius centrifuges. Interestingly, subjects report no sense of translation in either experiment during the constant velocity centrifugation. In orbit exposure to 0.2 Gz at the head during 60 s of constant velocity was insufficient to provide a vertical reference (Benson *et al.* 1997), possibly because of the opposing G-gradient along the trunk and legs and/or the relatively small resultant force level (Mittelstaedt 1999). When subjects were exposed to greater force levels (0.5 Gy and 1.0 Gz) for up to 5 min, these forces did provide a vertical reference on orbit. The subjects perceived roll-tilt when the resultant force was directed along the interaural axis, and inversion when the resultant force was directed towards the head (Clément *et al.* 2001). Ocular counterrolling was also unchanged during this experiment (Moore *et al.* 2001).

#### **2.4.6. Physiological Basis for Spatial Disorientation**

The physiological basis of spatial orientation perception became better understood with the discovery in rat and primate limbic systems of place cells that code the direction the animal is facing, independent of head movement. Also discovered were grid and place cells that code various attributes of location relative to visual landmarks (Wiener & Taube 2005), analogous to a map of the local environment. All three classes of cells respond in a navigation coordinate frame normally defined by the plane of locomotion, even in 0 g and hypergravity (Knierim *et al.* 2000; Taube *et al.* 2004). How larger (geo) scale environmental knowledge is coded is not yet understood, but clinical evidence from patients with poor geospatial abilities suggests that these same limbic structures at least participate.

### **2.5 Decrements in Cognitive Function**

Controlling vehicles and other complex systems can place high demands on cognitive and psychomotor functions. Space flight might affect these functions through direct microgravity effects (such as those described in the preceding sections) or through stress effects associated with sleep loss, workload, or the physical and emotional burdens of adapting to the novel, hostile environment. Kanas & Manzey (2003) and Strangman *et al.* (2014) have provided a good overview of the relevant evidence. Data collected on astronauts in space and during simulated missions have relied on a variety of human performance measures and test batteries, including the MINICOG Rapid Assessment Battery, the Advisory Group for Aerospace Research and Development Stress Battery (AGARD), the NASA Performance Assessment Battery (NASA-PAB). These tests focused predominantly on attention and memory (concentration, working memory, visuo-spatial memory and searching, verbal learning and recall, reaction time). According to Strangman *et al.* (2014) the available evidence, however, fails to strongly

support or refute the existence of specific cognitive deficits in low Earth orbit during long-duration spaceflight. The large variability in task performance within and between individuals argues against overgeneralizing from small samples and anecdotal reports, a caveat that reinforces the need for careful characterization and systematic study of cognition in space flight. To address this issue, a new neurocognitive test battery was recently developed and is currently being used on the ISS and in analog environments. The COGNITION Test Battery for Space Flight consists of 10 brief neurocognitive tests aimed at detecting cognitive performance changes in: motor praxis, visual object learning, fractal 2-back, abstract matching, line orientation, emotion recognition, matrix reasoning, digit symbol substitution, balloon analog risk, and psychomotor vigilance. These tests are also known to engage specific brain regions as evidenced by functional neuroimaging ([Basner et al. 2015](#)).

More details on the cognitive effects due to extreme isolation and confinement, feelings of remoteness, monotony and boredom, lack of privacy and personal space, depression, and fatigue can be found in the Behavioral Health and Performance Evidence Report (Slack et al. 2016). In the following sections we focus on the cognitive effects that are mainly attributable to reversible vestibular changes induced by spaceflight. As should be clear from the evidence presented here, space flight induces many of the hallmarks of a (reversible) vestibular lesion. Cognitive deficits, such as poor concentration, short-term memory loss, and inability to multi-task occur frequently in patients with vestibular abnormalities ([Jacob & Furman 1996, 2001](#)). [Hanes & McCollum \(2006\)](#) published a thorough review of the literature suggesting broader interactions between vestibular and cognitive function (including oculomotor, motor coordination, and spatial perception/memory effects similar to those described above) and demonstrating a physiological basis through observations of neuronal projections from the vestibular nuclei to the cerebral cortex and hippocampus. These results suggest that cognitive abilities may be most compromised during landing, particularly if an off-nominal event occurred that had not been recently well-rehearsed.

### **2.5.1. Mental Rotation**

The face of a well-known person is not as easily recognized when presented upside down. This phenomenon suggests that people have difficulty recognizing familiar shapes when they are in an unfamiliar orientation ([Howard & Templeton 1966](#); [Howard 1982](#); [Finke & Shepard 1986](#); [Cohen, 2000](#)). In such circumstances, mental rotation is necessary for shape and facial recognition. Until crewmembers become adept at mentally rotating themselves and/or their environment, and/or develop new spatial maps, they can easily become disoriented. Poor ability to mentally rotate the visual environment could be an important factor in determining susceptibility to space motion sickness ([Parker & Harm 1992](#)).

Mental image rotation and reconstruction experiments were performed on orbit. During STS-51G, mental reconstruction after body tilt demonstrated that the critical angle was 65°,

which is comparable to values attained on Earth (Corballis *et al.* 1978). After 3 days of microgravity, the subjects could mentally rotate the environment even while they were in an inverted position (Clément *et al.* 1987). The investigators proposed that mental rotation could be a gravity-dependent process and that weightlessness, by releasing this constraint, facilitates processing of visual images in any orientation. In contrast, Léone *et al.* (1995) suggested that mental rotation depends on symmetry detection and an internal vertical reference. In the absence of gravity, detection of symmetry was less accurate and required more time. The discrepancies between findings may be attributed to procedural differences, use of subject restraint, and difficulty of mentally manipulating test objects.

Experiments using mental rotation of three-dimensional objects (Shepard & Metzler 1971) in the yaw and roll axes, while subjects were restrained, showed that response times and error rates were similar before and during flight. These results were consistent with those of Friederici & Levelt (1987, 1990) and Léone *et al.* (1995), and support the conclusion that mental rotation of visually presented 3-D objects is independent of gravity.

### **2.5.2. Distance and Size Perception**

Further space flight-related changes occur in cognitive visual-spatial processing, which helps in perception of distance and size of objects. Distance and size perception are skills learned through repetitive practice. Normally sighted, binocular and even totally monocular people develop and use effective distance and size perception skills (Beaton *et al.* 2015). Monocular depth cues include angular variations—or parallax—when moving the head; texture, luminosity, color, and shading variations of the visual scene; and perspective.

In microgravity, the environment is not structured with a gravitational reference and a visual horizon, so perspective is less relevant. Astronauts perceive heights and depths of objects as taller and shallower on orbit, respectively (Lathan *et al.* 2000; Clément *et al.* 2013), and there are changes in the perception of geometric illusions and perspective-reversible figures after 3 months in space (Clément *et al.* 2012, 2015). These changes may occur because the perspective cues for depth are less salient in microgravity (Clément *et al.* 2015a). Ground-based studies also showed that the occurrence of geometric illusions based on perspective is reduced when subjects are tilted relative to the gravitational normal (Clément & Eckardt 2005) and in patients with vestibular deficits (Clément *et al.* 2009). Consequently, the changes in 3D visual perception in orbit reduction are presumably due to the altered peripheral otolith input or to a central adaptation in the processing of visuo-spatial cognitive function (Clément & Reschke 2008).

On Earth, horizontal distances relative to the observer are accurately estimated up to 4 m, and underestimated by approximately 10% as distance increases (Daum & Hecht 2009). By contrast, vertical distances are overestimated by about 30%, especially when looking down (Stefanucci & Profitt 2009). During a recent study, 6 astronauts were presented with stereoscopic (anaglyphs) photographs of natural scenes. Small yellow targets were superimposed on easily recognizable landmarks within each scene, e.g., a remarkable building, the end of a bridge, the top of a mountain, or the bottom of a tower. The subjects were asked to estimate the absolute distance between themselves and the target (egocentric distance)

using a conventional metric of their choice (e.g., feet, yards, or meters). On average, the astronauts reported distances above 50 m to be about 20%-25% smaller in-flight than pre-flight (Figure 4). One interpretation for this underestimation of distance in flight is that the distance between the eyes and the floor varies when astronauts are free floating; therefore they cannot use the eye height scaling to estimate distance and size as on Earth (Clément *et al.* 2013).




Figure 4. Differences in egocentric distance estimations using natural scenes between inflight and preflight conditions (Clément *et al.* 2013).

Distortions of the visual space during space missions may influence astronauts' ability to accurately perform cognitive and sensorimotor tasks, such as those involved in robotic operations. Additionally, this misperception will alter how astronauts view their habitat and workspace volume. These are important considerations for future human planetary exploration missions, and warrant further investigation and consideration for countermeasure development (Clément *et al.* 2013).

### 2.5.3. Time Perception

Some astronauts and cosmonauts have reported a "time compression syndrome" in orbit, whereby time is subjectively sensed as compressed relative to the perceptions gained during training and simulation (Schmitt & Reid 1985; Albery & Repperger 1992). Another perception experienced by the astronauts is that longer time than normal is required to execute standard mental activity (Manzey & Lorenz 1998). Yet another reported syndrome is "space fog", which affects cognitive performance during the first weeks of a mission (Welch *et al.* 2009).

Following the historical one-orbit flight of Gagarin, Titov flew on board Vostok-2 for a full day (17 Earth orbits), and performed the first cognitive neuroscience experiments in orbit. The objective of one experiment was to assess his ability to evaluate time intervals. After starting a stopwatch, he began to count 20 s in his mind; when he estimated subjectively that 20 s had passed, he stopped the stopwatch and looked at the actual elapsed time. The average time estimates during the 4 in-flight sessions were not significantly different from those measured during training, but they were biased by the fact that he had continuous feedbacks on his performance (Leonov & Lebedev 1968).

Another experiment on time perception in microgravity was performed on four astronauts during a 4-day Space Shuttle mission. Subjects viewed a visual target traversing a display and, while it was obscured, estimated the time of its arrival at a predetermined point by any means other than counting. The time perception for short duration tasks (2 s) were consistently overestimated in 0 g. As the time duration of the task increased, the subjects tended to underestimate. These errors in duration estimates increased each day as the flight progressed. Three hours after landing the duration estimates were also significantly larger than on flight day 4 (Ratino *et al.* 1988). These results suggest that the ability to estimate brief intervals of time deteriorate during a short space mission and after landing.

Similar effects were also observed in subjects exposed to hypergravity in a centrifuge (Albery & Repperger 1992). The authors point out that one potential consequence of these effects is that crewmembers who need to make quick decisions and perform critical tasks in-flight and re-entry may exhibit some delays in their responses, which would compromised safety. Clément *et al.* have proposed an experiment on board the International Space Station whose objective is to investigate whether the subjective perception of time is affected during and after long-duration space flight.

## **2.6. Space Motion Sickness**

### **2.6.1. History**

Space motion sickness (SMS) is the most clinically significant neurosensory phenomenon experienced by crewmembers during the first few days of space flight, and its effect may be almost as significant immediately after returning to Earth (Lackner and DiZio, 2006). The neurosensory and motor systems and their relationship with the vestibular system were extensively studied during the early Space Shuttle missions. The major focus during that time was the prevention of SMS, which is characterized by a plethora of symptoms, such as somnolence, vomiting, stomach awareness, fatigue, and performance decrements. No correlation was found between SMS and terrestrial motion sickness in terms of crewmember susceptibility. As a result of these studies, many preflight sensorimotor tests were abandoned due to their misrepresentations of induced mechanisms during actual space flight. On the other hand, post-flight postural instability and ataxia were evaluated and studied in greater detail to better understand the prevention of injuries during locomotion or emergency egress immediately after landing.

### **2.6.2. Symptoms**

On Earth, exposure to provocative motion, whether real or apparent, leads to the progressive cardinal symptoms of terrestrial motion sickness, which typically include the following: pallor, increased body warmth, cold sweating, dizziness, drowsiness, nausea, and vomiting. Although similar, the symptoms associated with SMS differ slightly from those of acute terrestrial motion sickness, probably because of differences in the physical environment, such as the lack of normal air-currentvection and of gravitational force on the contents of the stomach. In particular, sweating, except for palmar sweating (Oman *et al.* 1990), is uncommon during space flight, and flushing is more common than pallor. SMS, as compared to acute

terrestrial motion sickness, typically is more often associated with stomach awareness, vomiting, headache (due perhaps to headward fluid shifts), impaired concentration, lack of motivation, and drowsiness (Thornton *et al.* 1987a; Davis *et al.* 1988; Oman *et al.* 1990). Vomiting is usually sudden and infrequent, and is often not marked by prodromal nausea. Bowel sounds, obtained by auscultation, are decreased or absent in crewmembers experiencing SMS (Thornton *et al.* 1987b; Harris *et al.* 1997). Despite these differences, nearly universal symptoms are malaise, anorexia or loss of appetite, lack of initiative, and (for some) increased irritability.

### **2.6.3. Incidence**

Historically, as the size of space vehicles has increased, so has the incidence of SMS. No SMS was reported in either Project Mercury or Project Gemini (Homick 1985), but 35% of the Apollo Program astronauts and 60% of the Skylab Program crewmembers developed symptoms of SMS (Davis *et al.* 1988). The incidence of SMS was 67% among first-time flyers on the first 24 Space Shuttle flights. Statistically, symptom occurrence was not different between career vs. non-career astronauts, commanders and pilots vs. mission specialists, men vs. women, different age groups, or first-time vs. repeat flyers. However, observations of U.S. astronauts suggest that the incidence of post-flight sickness may be higher in women than in men (R. Billica, *personal communication*, 1999). Also, an astronaut's susceptibility to SMS on his or her first flight correctly predicted susceptibility on the second flight in 77% of the cases (Davis *et al.* 1988). Later estimates indicated that 80% to 90% of all Shuttle crewmembers experienced some symptoms of motion sickness (R. Billica, *personal communication*, 1999).

Russian researchers report that 54% of cosmonauts have symptoms lasting 1 to 3 days, 25% have symptoms lasting 14 days or longer, and 8 of 46 cosmonauts on long-duration missions (85-365 days) periodically developed vertigo and queasiness, especially during the last 10 to 14 days of the mission, when their activity increased (Bryanov *et al.* 1986; Kornilova *et al.* 1995).

No reports of post-flight motion sickness (PFMS) were noted in the Space Shuttle program through the mid-1980s (Thornton *et al.* 1987a). However, it now appears that this syndrome affects a similar percentage of both U.S. and Russian crews. The Russian reports indicate that PFMS symptoms generally occur in cosmonauts who have SMS in-flight. However, 11% of those who experience little or no SMS on orbit do experience mal de débarquement (Bryanov *et al.* 1986). Postflight medical debriefs were examined for Shuttle missions from the beginning of the program, in April 1981, through January 1999, which involved 241 crewmembers having flown between one and six missions. Postflight, 32% of crewmembers reported vertigo, 14.7% reported nausea, and 8% vomiting (Bacal *et al.* 2003).

PFMS onset occurs in a time pattern similar to that of SMS. Within minutes of g-force onset during re-entry symptoms may already be developing. Crewmembers who have no symptoms during re-entry and landing may develop symptoms as soon as they stand up to exit the vehicle. The severity of the symptoms and the functional recovery seem to be directly proportional to the time on orbit. There have been reports of a "relapse" phenomenon in the post-landing

recovery course. Astronauts who are exposed to certain types of inertial environments, like turning a corner in a car or lying in bed in the dark, can bring on a sudden return to an early postflight state of maladaptation, which may elicit 'mild' to 'severe' PFMS symptoms several days up to a week after return to Earth. Recovery from this "relapse" generally occurs more rapidly than the recovery immediately after returning from orbit (Ortega & Harm 2008).

Like SMS, PFMS does not appear to correlate with gender, age, crew position, or number of previous flights (Reschke *et al.* 2014). Past experience with postflight re-adaptation does not seem to affect incidence (Bacal *et al.* 2003). PFMS is likely complicated by the relative dehydration upon return and orthostatic intolerance following flight.

#### **2.6.4. Provocative Stimuli**

Microgravity by itself does not induce space sickness. The larger a volume of spacecraft and the mobility of their inhabitants, the higher chance of SMS. Specifically, factors that may initiate or worsen SMS include distasteful, unpleasant, or uncomfortable sights, noxious odors, certain foods, excessive warmth, loss of 1-g orientation, and head or whole-body movements (Jennings 1998). Similarly, post-flight symptoms may be induced and/or exacerbated by warmth and head movements during reentry and immediately after landing. Hypersensitivity to angular head motions is also common; crew members have reported that head movements in the pitch plane are initially more provocative than those made in other planes (Thornton *et al.* 1987a; Oman *et al.* 1990).

#### **2.6.5. Theories and Hypotheses**

Many theories and hypotheses have been proposed to explain SMS. The *fluid shift* theory (Barrett & Lokhandwala 1981; Parker *et al.* 1983) postulates that headward fluid shifts accompanying weightlessness produce changes that alter the response properties of vestibular receptors. However, this theory is not ideal in that it fails to adequately address the development of motion sickness during space flight.

The *sensory conflict* theory from Reason & Brand (1975) assumes that human orientation in three-dimensional space, under normal gravitational conditions, is based on at least four sensory inputs (otolith organs; semicircular canals; visual system; and touch, pressure, and somatosensory systems) to the CNS. Motion sickness may result when the environment is altered in such a way that information from sensory systems is not compatible and does not match previously stored neural patterns. It is important to note that it is the combination, rather than a single course, of these conflicts that somehow produces sickness, although the exact physiological mechanisms remain unknown.

In the *poison* theory, Treisman (1977) suggested that the purpose of mechanisms underlying motion sickness, from an evolutionary perspective, was not to produce vomiting in response to motion, but to remove poisons from the stomach. Money and colleagues (1970, 1996) concluded that the mechanism to facilitate vomiting in response to toxins is partly vestibular.

The *otolith mass asymmetry* hypothesis describes a mechanism complementary to the sensory

conflict theory that explains adaptation to weightlessness, readaptation to 1 g, and individual differences in susceptibility to SMS (von Baumgarten & Thuemler 1978; von Baumgarten *et al.* 1982; Kornilova *et al.* 1983). According to this theory, some individuals possess slight functional imbalances between right and left otolith receptors that are compensated for by the CNS in 1 g. *Sensory compensation* (Parker & Parker 1990) occurs when the input from one sensory system is attenuated and signals from others are augmented.

Finally, the *Otolith Tilt-Translation Reinterpretation (OTTR)* hypothesis assumes that because of the fundamental equivalence between linear acceleration and gravity, graviceptors signal both the head orientation with respect to gravity (tilt) and a linear acceleration of the head that is perceived as translation. As a consequence of the absence of sensed gravity during orbital flight, graviceptors do not respond to static pitch or roll in weightlessness; however, they do respond to linear acceleration. Because stimulation from gravity is absent during space flight, interpretation of the graviceptor signals as tilt is meaningless. Therefore, during adaption to weightlessness, the brain reinterprets all graviceptor output to indicate translation (Young *et al.* 1984; Parker *et al.* 1985). The *sensory compensation* and *OTTR* hypotheses have both been further refined by Merfeld (2003) and Clément & Reschke (2008).

#### **2.6.6. Predicting Susceptibility**

The prediction of susceptibility to motion sickness has long been of interest to space flight researchers. Since most motion sickness treatments are more effective when they are administered before symptoms develop, the identification of individuals susceptible to SMS would allow preventive measures to be taken only by those requiring them, and would free insusceptible persons from the undesirable side effects of anti-motion sickness medications and/or the scheduling requirements of pre-training (Diamond & Markham 1991).

A number of predictors for motion sickness have been investigated and can be grouped into the following categories: exposure history, physiological predisposition, psychological predisposition, plasticity, provocative tests, and operational measures (Clément & Reschke 2008). The real test of any predictive method relies on the use of data from crewmembers. Ground-based measures on normative subjects, while useful, are not true measures of the criterion of interest, SMS. Until enough flight data become available, along with ground-based tests for flight personnel, the relationships between various predictors and SMS susceptibility will remain unclear.

#### **2.7. Impacts of Vestibular Changes to Orthostatic Intolerance**

The vestibular system is required for motion sickness to occur, but the exact neural pathways involved are still unknown. Understanding the role of the vestibular system in autonomic regulation is essential to understanding SMS (Davis *et al.* 1988, 1993). Some authors argue that unusual motion or direct vestibular stimulation triggers a poison response (Money *et al.* 1996). The poison response consists of a stress response and stomach emptying. Vomiting is primarily associated with increases in parasympathetic activity, whereas stress responses are primarily associated with increases in sympathetic activity or decreases in parasympathetic activity (or both). The most direct pathway for vestibular modulation of autonomic responses

involved in motion sickness is efferent projections from the medial and inferior vestibular nuclei to the nucleus tractus solitarius (NTS) in the brainstem and the dorsal motor nucleus of the vagus. The NTS plays an integral role in both gastric motility and emesis related to motion sickness (Ito & Honjo 1990). The NTS receives input from peripheral and central ascending fibers (Onai *et al.* 1987; Barron & Chokroverty 1993), and in turn influences vagal stimulation of the stomach and heart and activation of the sympathetic nervous system (Yates 1992; Previc 1993; Yates *et al.* 1993; Yates & Miller 1996). The cerebellum may be another route through which vestibular inputs may modulate autonomic activity (Wood *et al.* 1994; Balaban 1996).

## 2.8. Peripheral and Central Neural Changes Associated with Space Flight

The radiation environment in exploration class missions poses a greater threat to the CNS than the environment for ISS astronauts since they are partially protected by the Earth. During a short-duration mission, astronauts may experience detrimental CNS changes in cognition, short-term memory, motor function, and behavior. Late responses to radiation damages can include premature aging, Alzheimer's disease, or other dementia (Cucinotta 2014). Radiation exposure in low Earth orbit is ultimately caused by galactic cosmic rays (GCR) and solar particle events (SPEs). Although all dose regimens have the potential to cause decrements in performance and cognition, it is the low to moderate levels (~ 1 to 2 Gy) of charged-particle exposure that will define the space radiation environment for crewmembers during long-duration space flights (Nelson 2009). In a study in which mice were exposed to 0.1 and 1 Gy of whole body proton irradiation, dendritic complexity was significantly dependent upon dose reduction, and the number and density of dendritic spines along hippocampal neurons of the dentate gyrus was significantly reduced (Parihar 2014).

Vertebrates sense gravito-inertial acceleration by mechanoreceptors in the otolith organs of the inner ear. These structures consist of ciliated sensory hair cells with crystals of calcium carbonate called otoconia placed on top of the hair cells that stimulate the hair cells when moved due to linear acceleration or tilt. Inner ear structures can regulate their function through adaptive processes by increasing or decreasing production of calcium carbonate in response to a sustained decrease or increase in the amplitude of gravity, altering otolith mass and subsequent transduction gain of the system. Boyle *et al.* (2015) used electron microscopic techniques to image otoconia mass obtained from (a) mice exposed to both 91-days of weightlessness in the ISS and 91-days of 2 g centrifugation on ground, and (b) mice flown on short-duration Shuttle missions. Results from ISS showed a clear restructuring of individual otoconia with increased deposition and mass while the 2 g counterparts showed a decrease in otoconial mass. Conversely, for shorter duration exposures to weightlessness (13-days) the otoconia appeared to be normal. Therefore, long-duration exposure to space flight may induce adaptive mechanisms that lead to structural alterations in peripheral end organ transduction of motion contributing to behavioral disturbances.



The brain may also undergo structural remodeling as a result of microgravity exposure, radiation exposure, and vascular changes associated with space flight (Carpenter *et al.* 2010). The brain's structural organization is not fixed but rather can undergo extensive remodeling, even in the adult brain. These changes occur in response to skill learning (Karni *et al.* 1995),

recovery from brain insult such as stroke (Cramer *et al.* 1997), and as a result of cognitive training (Olesen *et al.* 2004). Moreover, brain plasticity occurs as a result of disuse such as prolonged bed rest (Roberts *et al.* 2010). Animal studies have shown that microgravity exposure results in structural brain changes. Research with rats has demonstrated that brain structural changes occur as a result of microgravity exposure, particularly in the somatosensory cortex (Krasnov 1994; Newberg 1994; D'Amelio *et al.* 1998) and cerebellum (Holstein 1999; Holstein & Martinelli 2003). These changes include decreased synapses and degeneration of axonal terminals. It has been demonstrated that hair cells in the rat utricular macula undergo extensive plasticity as a result of space flight, with a large (40-55%) increase in synapse number (Ross 1993, 1994, 2000, 2003). This plasticity remained evident following the flights, even after posture control in the rats had returned to normal. Therefore, in humans potential structural changes associated with long-duration space flight could have behavioral implications for both space flight operations and long-term health of crewmembers. Studies are currently being conducted with ISS crewmembers that will identify potential changes in brain structure and function following long-duration space flight using magnetic resonance imaging techniques to assess the risks of changes in brain structure and the impact on sensorimotor and cognitive function (Erdeniz *et al.* 2013; Koppelmans *et al.* 2013).

### 3. EVIDENCE OBTAINED FROM SCIENTIFIC INVESTIGATIONS DURING REENTRY AND AFTER LANDING

#### 3.1. Decrements in Postural Equilibrium

The first studies, designed to quantify post-flight postural ataxia, required astronauts, upon landing, to tandemly stand on narrow rails of various widths with their eyes either open or closed and arms folded across their chests (Berry & Homick 1973; Homick & Miller 1975; Homick *et al.* 1977; Homick & Reschke 1977; Kenyon & Young 1986). Other studies have used static force plates for stabilometry and simpler tests, such as the clinical Romberg test, a sharpened (toe-to-heel) Romberg test, and vertical posture with varying head positions, to assess postural ataxia immediately after flight (Bryanov *et al.* 1976; Yegorov 1979; Clément *et al.* 1984). Other postural performance studies have relied on dynamic posture platforms that translate the subject (Reschke *et al.* 1984; Clément *et al.* 1985; Anderson *et al.* 1986), tilt the subject (Kenyon & Young 1986; Reschke *et al.* 1991), or provide more sophisticated means of posture control such as stabilization of ankle rotation and/or vision (Paloski *et al.* 1993). Pre- and post-flight studies of vestibulo-spinal reflexes (Baker *et al.* 1977; Reschke *et al.* 1984; Kozlovskaya *et al.* 1984; Watt *et al.* 1986) and postural responses to voluntary body movements (Clément *et al.* 1984; Reschke *et al.* 1988) have also been performed.



*Figure 5. Equilibrium (EQ) score during computerized dynamic posturography evaluation after ISS missions. High EQ score means more stability. Data points show responses for individual crewmembers preflight and postflight. The gray area represents the 90% confidence limit region around the median score (red line) for all crewmembers. Adapted with permission from Wood et al. 2015.*

The greatest decrease in stability occurs when subjects must rely on vestibular information alone, when proprioceptive or visual feedback is either altered or absent. Decrements in postural stability with eyes closed are well documented from Skylab missions and were observed to persist for up to 1-2 weeks post-flight (Homick et al. 1977). Computerized dynamic posturography (CDP) evaluations conducted after ISS flights clearly demonstrated that postural

stability was diminished (Figure 5). Recovery of postural stability after landing occurs in two phases: initial rapid improvement followed by a gradual recovery. Both the initial decrement and the time required for recovery vary as a function of flight duration. Stability is further compromised during dynamic head tilts (0.33Hz @  $\pm 20^\circ$  in pitch plane), with a time constant of recovery of 19hr for head erect versus 111hr for head moving, so that the majority of crewmembers are unable to maintain quiet stance for 20 sec. The extent of the postural stability decrement may also reflect previous space flight experience (Clark & Bacal 2008). Paloski *et al.* (1990, 1993, 1994) reported that veteran astronauts seem to have better post-flight postural performance than first-time fliers. A more recent within-subject analysis of CDP data comparing performance decrements on a similar post-flight day suggests that ~60% of veteran astronauts were more stable following repeat flights (2010, not publicly available).

A few physiological factors are known to contribute to space flight-induced postural instability (Forth *et al.* 2011). Firstly, the dorsiflexor muscles play a larger role in space than the extensor muscles, which are used to counteract gravity on Earth (Clément *et al.* 1984). This causes astronauts to assume a forward tilted posture when asked to stand perpendicular to the spacecraft floor. A small flexor tone is generated in order to maintain the feet at a right angle to the leg, as this is the normal neutral posture of the ankle (Clément & Lestienne 1988). Another explanation for this posture in microgravity is that the normal excitatory drive exerted by input from the otoliths, by way of the vestibular nuclei, on the extensor muscles is inhibited due to the lack of gravity.

Changes in vestibulo-spinal reflexes may also contribute to postural decrements. The Hoffmann reflex and otolith-spinal reflexes were dramatically reduced during space, but differences between pre- and post-flight responses were not significant (Watt *et al.* 1986). Reschke *et al.* (1986) observed a potentiation of the Hoffman reflex 40 ms after astronauts underwent an unexpected drop (Earth-vertical fall with bungee cords) during flight; this potentiation disappeared after 7 more days. Immediately after space flight, there was significant potentiation again compared to preflight responses. Such changes in the Hoffman reflex are predictive of change in the gain of the spinal reflex pathway. How gain changes in this pathway are linked to preprogrammed muscular activity such as the maintenance of posture is not clear.

## **3.2. Decrements in Locomotion**

### **3.2.1. Locomotor Control and Segmental Activation**

Most crewmembers experience some degree of locomotor dysfunction when they return to Earth after space flight. Some of these post-flight alterations are ataxia with the sensation of turning while attempting to walk a straight path; sudden loss of postural stability when rounding corners or after unexpected perturbations; and sudden loss of orientation in unstructured visual environments. In addition, some astronauts report oscillopsia (illusory movement of the visual field) while walking.

Foot contact with the ground, the transfer of weight from one foot to the other, and the pushoff with the toe from the ground are critical phases, as these interactions with the support

surface result in forces that create vibrations, which if unattenuated could interfere with the visual-vestibular sensory systems in the head (Voloshin 1988; Smeathers 1989; Pozzo *et al.* 1990; Valiant 1990; Lafortune *et al.* 1996; Ito *et al.* 1997; McDonald *et al.* 1997; Whittle 1999; Mulavara *et al.* 2005a; Mulavara & Bloomberg 2003). The musculoskeletal system controls these vibrations; muscles and joints act as filters to minimize the perturbing effects of impacts with the ground and help to maintain a stable trajectory at the head (Holt *et al.* 1995, McDonald *et al.* 1997). During treadmill walking after returning from long-duration space flight, astronauts' knee flexion during the stance phase significantly increased, but it returned to normal within 6–10 days (Bloomberg & Mulavara 2003). An increase in knee flexion during locomotion will result in reduction of the axial stiffness of the lower-limb complex during the critical stance phase following heel strike, leading to reduced perturbations being transmitted to the head during locomotion.

Distinct post-flight performance decrements in gait have been observed in cosmonauts returning from Soyuz missions lasting 2 to 63 days (Chekirda *et al.* 1971; Bryanov *et al.* 1976; Chekirda & Yermin 1977). In most cases, the duration of post-flight effects correlated with the duration of the mission. Post-flight changes in locomotion included increased angular amplitude of motion at the knee and ankle, and increased vertical accelerations of the center of mass (Hernandez-Korwo *et al.* 1983).

Post-flight locomotor control and segmental coordination show alterations in muscle activation. Layne *et al.* (1997, 1998b) reported that the temporal relationship and relative amplitude of muscle activation are modified by space flight, particularly for the events of heel-strike and toe-off that are complementary to evidence of changes in kinematics during locomotion at these events of the gait cycle (McDonald *et al.*, 1996; Miller *et al.* 2010). The loss of neuromuscular coordination may cause difficulty in achieving optimal transitions between muscles while walking (Courtine *et al.* 2002). This loss in coordination between muscles may also affect the ability to maintain stable head movement (Layne *et al.* 2004, 2001) and are complimentary to evidence seen in alterations in head-trunk coordination during walking (Bloomberg and Mulavara, 2003; Bloomberg *et al.*, 1997; Mulavara *et al.*, 2012); reduced visual acuity during walking (Peters *et al.*, 2011) and impairment in the ability to coordinate effective landing strategies during jump tasks (Newman *et al.*, 1997; Courtine and Pozzo, 2004). Other neurophysiological changes including proprioceptive hyper-reactivity, such as increased Hoffman reflex amplitudes; reduced ability to perform graded muscle contractions; and decreased muscle stiffness (Grigoryeva & Kozlovskaya 1987) may also have a contributory influence on muscle coordination during locomotion activity.

Activation of ankle-joint muscles post-flight has been heavily investigated. Zangemeister *et al.* (1991) concluded that space flight-related adaptive modifications in neural processing of vestibular input could negatively influence activation of lower limbs. This change, in combination with altered strength between ankle plantar flexors and dorsiflexors (Hayes *et al.* 1992), can cause difficulty in walking.

Changes in ankle musculature activation are suggested to result in reinterpretation of proprioceptive input (Roll *et al.* 1993). Ankle proprioception is no longer interpreted as coding

anterior-posterior body sway while upright, but instead codes either whole body axial transportation (i.e., pushing off the support surface) or foot movement exclusively. In addition, the extent of reinterpretation correlates with duration of mission. Despite being appropriate for weightlessness, these changes are maladaptive on return to Earth and may contribute to post-flight decrements in locomotion.

### **3.2.2. Head-Trunk Coordination**

Gaze and head movement control stabilizes vertical and horizontal vision during various motor tasks such as jumping, walking, running, hopping and/or tasks requiring a person to maintain equilibrium on a beam or a moving platform ([Assaiante et al. 1993](#); [Pozzo et al. 1995](#)). The head serves as a stable platform to provide a veridical reference frame for visual, vestibular, and proprioceptive integration, facilitating the organization of postural and locomotor control patterns ([Pozzo et al. 1990](#); [Bloomberg et al. 1992](#); [Bloomberg and Mulavara 2003](#); [Mulavara et al. 2002](#); [Mulavara et al. 2003](#); [MacDougall & Moore 2005b](#)). Another system that benefits from head stabilization with respect to the environment is the system for maintaining gaze during body movement. Head movements actually contribute to gaze stabilization during locomotion ([Bloomberg et al. 1992](#)). An example of this is the pitch head rotation (as in nodding the head), which compensates for the vertical translation of the trunk that occurs with each step during the gait cycle. The magnitude of these head rotations was observed to be controlled as a function of the distance of the visual target to the eyes. The goal-directed nature of these head movements during concurrent locomotion and visual target fixation suggests that head movements are not completely dependent on passive inertial and viscoelastic properties of the head-neck system but are actively modulated to respond to altered gaze control requirements. Thus, head stabilization mechanisms help adjust posture, to maintain balance of the moving body, and maintain visual acuity for navigational control through a constantly varying environment during locomotion.

In an initial study, [Bloomberg et al. \(1997\)](#) examined whether short-duration exposure (7–16 days) to the microgravity environment of space flight during Shuttle missions induces alteration in post-flight head-trunk coordination during locomotion. Before space flight, pitch head movements acted in a compensatory fashion to oppose vertical trunk translation during locomotion. As the trunk translated upward, the head pitched forward and downward, assisting in maintaining target fixation. Following space flight, coordination between compensatory pitch angular head movements and vertical trunk translation was significantly altered. Like patients with vestibular deficits ([Keshner 1994, 1995](#)) and children prior to development of their mature head stabilization response ([Assaiante et al. 1993](#)), head movements were restricted during locomotion. This change in head-trunk coordination strategy may account, in part, for the reported oscillopsia that occurs during post-flight locomotion and may contribute to disruption in descending control of locomotor function. Comparison of responses from multi- and first-time astronauts indicated that astronauts who had experienced more than one space flight demonstrated less post-flight alteration in the frequency spectra of pitch head movements than subjects on their first flights.

Changes in head-trunk coordination during locomotion were also characterized in returning Mir crewmembers (Bloomberg & Mulavara 2003). These subjects walked (6.4 km/h) on a treadmill before and after space flight while visually fixating on an earth-fixed target. At this walking speed, head pitch movements compensate for the vertical trunk movements that occur during each step (Pozzo *et al.* 1990, 1995; Bloomberg *et al.* 1997). Subjects showed a reduction in head movements in the frequency range of 1.5-2.5 Hz reflecting the contributions of reflexive head stabilization mechanisms (Keshner *et al.* 1995) during post-flight locomotion followed by a recovery trend spanning several days. This reduction in head pitch movement occurred despite no significant change in trunk pitch or vertical movement. Therefore, during post-flight locomotion head movement amplitude with respect to space was reduced.

Bloomberg *et al.* (1997), Davids *et al.* (2003), Mulavara *et al.* (2005a) and Madansingh and Bloomberg (2015) have shown that tasks requiring sensorimotor integration after an adaptive exposure are associated with a wide range of adaptive behavioral responses. Specifically, after short-duration space flight, astronauts showed diverse responses, with some showing increases and others showing decreases in the magnitude of head pitch movement during walking (Bloomberg *et al.* 1997). A report from Mulavara *et al.* (2012) confirms and extends this observation of response variability one day after return from long-duration space flight on board the ISS. Subjects were classified into two groups according to the pre- and post-flight averages of the magnitude of their head pitch movements during locomotion: a “decreaser” group wherein subjects’ post-flight average head pitch movements decreased with respect to their preflight average, and an “increaser” group, wherein subjects’ post-flight average exceeded their preflight average. The vertical torso translation was not significantly different after exposure to space flight compared to preflight.

Zangemeister *et al.* (1991) demonstrated that normal locomotion performed with the head in a retroflexed position induces alterations in lower limb muscle activity patterns. They concluded that a functional linkage exists between otolith signals generated by various head positions and the muscle activity patterns generated in the lower limbs during locomotion. Appropriate attenuation of energy transmission during locomotion, achieved by the lower limbs’ joint configuration coupled with appropriate eye-head-trunk coordination strategies, was demonstrated as being fundamental feature of an integrated gaze-stabilization system during locomotion (Bloomberg and Mulavara 2003; Mulavara *et al.* 2002/2003; Mulavara *et al.* 2005). From this point of view, the whole body is an integrated gaze-stabilization system to which several subsystems contribute, leading to accurate visual acuity during body motion. Given these functional linkages, it can be argued that space flight induces adaptive modification in segmental coordination and disrupt coordinated body movement during post-flight terrestrial locomotion. It follows that active body movement in the unique inertial environment encountered during space flight may require subjects to adaptively acquire novel head-trunk and lower body segmental control strategies. Adaptation to long-duration space flight leads to modified reflexive head stabilization mechanisms; modified transmission characteristics of the shock wave at heel strike; and increased total knee movement during the subsequent stance phase during post-flight walking. These strategies, however, may be maladaptive for

locomotion in a terrestrial 1 g environment leading to impairment of locomotor function during the readaptation period following the return to Earth.

Previous space flight data have shown post-flight increases in vestibulo-spinal reflexes in humans and increased utricular afferent sensitivity to translation shown in toadfish (Reschke *et al.* 1984; Boyle *et al.* 2001). Upregulation of the sensitivity of the utricular afferents after adaptation to 0 g may be manifested in the “increaser” group of astronaut subjects. However, a similar increase in vestibular sensitivity would tend to generate larger head pitch for the same trunk vertical translation during locomotion (Moore *et al.* 2006). A similar increase in head movements in the pitch direction was observed in all subjects who had undergone 30 minutes of adaptation to locomotion while unloaded to 60% of body weight with no changes in trunk vertical translation during locomotion (Mulavara *et al.* 2012). This was not the case for the subjects in the “decreaser” group of astronaut subjects. Data from experiments with labyrinthine-deficient patients and experiments in which galvanic vestibular stimulation (GVS) was used to induce acute vestibular disturbances indicated that a reduction in head movement response could be a voluntary strategic response to reduce sensory conflict (Mulavara *et al.* 2012). Thus, the “decreaser” group of subjects may show an increased weighting of vestibular signals and hence sensory weighting may be a marker of post-flight disturbance. This strategy may also reflect the response of a control system looking for a new equilibrium point. The goal of establishing this new end point would be to reduce potential canal-otolith ambiguities.

The lack of change in vertical trunk translation indicates that the input disturbances to the gaze control system remain unchanged. Taken together, the kinematic and gaze stabilization findings indicate that body load-sensing somatosensory input centrally modulates vestibular input and can adaptively modify vestibularly mediated head-movement control during locomotion. Thus, space flight may cause a central adaptation of the converging vestibular system and the body load-sensing somatosensory system, leading to alterations in head-movement control.


### **3.2.3. Dynamic Visual Acuity**

Gaze control orchestrated by the CNS is critical to dynamic visual acuity (DVA), the ability to see an object clearly when the object, the observer, or both are moving. Deficient gaze control experienced following G-transitions cause oscillopsia, or blurred vision, and decrements in dynamic visual acuity, with stationary objects appearing to bounce up and down or move back and forth during head movements. Decreased dynamic visual acuity caused by space flight can lead to misperception of sensory information and poses a unique set of problems for crewmembers, especially during entry, approach, and landing on planetary surfaces. Visual disturbances could adversely affect entry and landing task performance, such as reading instruments, locating switches on a control panel, or evacuating a vehicle in suboptimal visual conditions (e.g., smoke in the cabin). Post-flight oscillopsia and decreased dynamic visual acuity could decrease crewmember safety when returning to normal duties (e.g., driving a rover, scuba diving, or piloting an aircraft) or activities of daily living (e.g., driving, contact sports, climbing ladders, etc.) after flight.

Measures of dynamic visual acuity (DVA) have been used as a diagnostic tool for identifying vestibular dysfunction (Lee *et al.* 1997; Hillman *et al.* 1999; Schubert *et al.* 2001, 2002; Tian *et al.* 2001). However, even persons with healthy vestibular function can experience, under certain conditions, compromised visual performance (Deshpande *et al.* 2013). Human factors (i.e. ergonomics) investigations looking at the effects of whole-body vibration have documented changes in visual performance over a wide range of stimulus conditions (Meddick & Griffin 1976; Moseley & Griffin 1986; Boff & Lincoln 1988; Griffin 1990; Demer *et al.* 1993). An important factor for determining the visual performance in these investigations is the transmissibility of the vibration to the head. Factors such as the subject's posture and muscle tone, as well as their coupling to contact surfaces or added masses, can have an effect on visual performance. The coupling between astronauts and their spacecraft during critical phases of the mission (e.g., entry, landing) could therefore affect their ability to see clearly.

McDonald *et al.* (1997) discussed the implications to gaze control of adaptive changes in musculoskeletal impedance and posture after space flight. Musculo-skeletal impedance is also affected by G-loading, which in turn affects vibration sensitivity; G- and vibration-loading often occur together during launch and entry/landing. Visual performance may well be degraded while standing during piloting, as potentially proposed for future planned space flight missions and previously employed during the Apollo program.

Decreased DVA performance was demonstrated in astronauts following return from long-duration space flight (Bloomberg and Mulavara 2003; Peters *et al.* 2011). A second-generation test using Landolt C characters instead of numbers also documented decrements in DVA performance as a function of time after flight in crewmembers returning from long-duration space missions (Peters *et al.* 2011) (Figure 6). For some subjects the decrement was greater than the mean acuity decrement seen in a population of vestibular impaired patients collected using a similar protocol. The population mean showed a consistent improvement in DVA performance during the two-week post-flight recovery period. These results may significantly underestimate the decrements in visual performance that are actually experienced during and immediately following landing because all DVA data (with the exception of one subject) were collected no earlier than 24 hours after landing. Given how rapidly VOR function and gaze control re-adapts the decrement in visual acuity at the actual time of landing was likely much higher than measured during the first postflight data collection session.



*Figure 6. Dynamic visual acuity data during treadmill walking from 14 crewmembers after return from 6-month ISS flights. Data are normalized to the subjects' preflight DVA values, which are represented on the y-axis at 0. These data show a decrement in postflight walking acuity followed by an improvement in performance during the postflight recovery period.*

Changes in dynamic visual acuity also contribute to functional changes (on the ground) in patients with vestibular disorders (Herdman 1994). For various reasons, physicians often caution patients with vestibular disorders against driving (Cohen et al. 2003). One such patient, referred to a rehabilitation program, specifically identified an inability to stabilize visually the car's instrument panel as a reason for self-limiting driving (H. Cohen, personal communication). Clark & Rupert (1992) report on a case study involving a student naval aviator with a similar complaint. Turbulence caused the aviator to become unable to see the instrument panel clearly. Testing revealed that the student had defective VOR function. As a result, his eye movements were not able to compensate adequately for the motions of his body in turbulent conditions.

### 3.3. Decrements in Jump Performance

Following space flight crewmembers also experience changes in the otolith-spinal reflex mechanisms that are essential for the preprogrammed motor strategies used for impact absorption after a jump. Watt et al. (1986) tested Shuttle astronauts during sudden "drops" and reported that all subjects were unsteady post-flight. The otolith-spinal reflex, which helps prepare the leg musculature for impact in response to sudden falls, is dramatically reduced during space flight. Reschke et al. (1986) used the Hoffman-reflex to examine the effect of drops on the sensitivity of the lumbosacral motoneuron pool, which is presumably set by descending otolith control signals. A large potentiation of the Hoffman-reflex recorded in the soleus muscle was found beginning approximately 40 ms following an unexpected drop. This potentiation of the Hoffman reflex during drops vanished on the 7th day of space flight. Immediately following space flight, two of four subjects demonstrated a significant increase in potentiation during the drop compared with pre-flight testing.

Shuttle astronauts were also tested pre-flight and post-flight (< 4 hr after return) on voluntary 2-footed downward hops from a 30 cm high step. Motion analysis of the jump

indicated impairment in the ability to coordinate effective landing strategies (Newman *et al.* 1997; Courtine & Pozzo 2004). A decrease in hip flexion and changes in the center of mass position relative to the feet were observed. The majority of crewmembers fell backwards (likely due to a potentiated stretch reflex) on the first of three jumps, and there was a greater use of arms for balance. These data provide further evidence for post-flight changes in motor programming during the jump aerial phase and impaired ability to prepare the limb muscles for the impact phase of the jump.

Other work indicates that space flight may affect proprioception of limb position (see section IV, 2.1.2). Watt *et al.* (1997) found a considerable decline in arm-pointing accuracy while blindfolded during and immediately following space flight. When they performed deep rhythmic knee and arm bends after space flight, subjects reported that floors and walls moved toward them, and that their knees bent more rapidly than intended. This can be partly explained by an abnormal level of muscle spindle receptor activation on return to 1 g, which results in misinterpretation of muscle length and subsequent abnormal flexion. Therefore, altered jumping performance seen post-flight may reflect decrements in limb proprioception sense, combined with altered central interpretation of otolith acceleration cues and vestibulo-spinal reflexes.

### **3.4. Perceptual Changes**

#### **3.4.1. Crew Reports**

After Shuttle missions, crew members were asked to indicate whether or not they experienced any of the 21 neurovestibular symptoms listed on a standardized questionnaire. It is important to keep in mind the subjective nature of these reports, as well as the possibility that some crewmembers might not have reported truthfully for personal reasons. It is unknown exactly at what point the astronauts actually felt the reported symptoms.

Nevertheless, out of 88 commanders reporting postflight:

- 9.09% reported persistence of illusory motion phenomenon
- 51.1% reported malaise
- 52.2% reported dry heaves
- 52.3% reported stomach awareness
- 60.2% reported sweating
- 61.4% had nausea
- 61.4% vomited
- 61.4% had persisting sensation aftereffects
- 68.1% had blurred vision
- 68.1% had headaches
- 65.9% had vertigo while walking
- 72.7% had difficulty walking in a straight line
- 69.3% had vertigo while standing
- 77.2% noticed clumsiness in their movements

Crewmembers also report a number of perceptual changes after space flight that may hinder their work performance: (a) *modified sensations of movement*, i.e., a feeling that they or the visual surround are moving following a simple head movement in any plane; (b) *sensations of excessive body weight* up to 2 to 2.5 times normal weight or that *objects, when handled, are heavier than normal*; (c) *sensations of greater bodily tilt* when the body is tilted from the vertical axis; and (d) *sensations of forced movement* of the body when turning corners.

Most perceptual disturbances and other symptoms resolve rather quickly, within 48 hours. Others, such as postural ataxia, may last with varying degrees of effect for months. During this prolonged recovery phase, tasks that require integration of multiple sensory inputs, e.g., head tilts while on unstable support surfaces, often reveal an underlying deficit that may not be revealed with more restricted movements. Interaction with the environment can shorten recovery times.

### **3.4.2. Proprioceptive Changes**

Overall, the post-flight reports of increased heaviness of static objects suggest that some central rescaling of static pressure systems occurs. A Spacelab D-1 mission required subjects to use higher accelerative shaking forces, which improved their ability to discriminate mass but not weight. Additionally, video recordings of astronauts showed that shaking was faster in-flight compared to preflight and slowed after landing, returning to baseline after three days ([Ross et al. 1986b](#)). Error in weight or mass perception may be due to a basic failure of reafference or to inadequate monitoring of command signals and inappropriate scaling of afferent signals.

### **3.4.3. Perceived Tilt and Translation**

Crews typically report that when they tilt their heads, they feel that the “gain” of their head tilt sensation is increased, as if their head had rotated farther than expected. A typical pilot comment is: “That really tumbled my gyros” ([see Section IV 1.1](#)). The sensation is thus reminiscent of the conventional hypergravic G-excess illusion. Other returning astronauts describe a transient sensation of horizontal or slightly upwards linear translation as a result of head tilt ([Parker et al. 1985](#); [Reschke et al. 1994](#); [Harm et al. 1999](#)). One of the most common post-flight illusions is of perceived translation, either of self or surround, during a tilting motion ([Harm & Parker 1993](#)). In one of the first post-flight experiments to investigate this phenomenon, a parallel swing was used to provide horizontal (interaural axis) translation and/or roll rotation about the head naso-occipital axis. All six astronauts participating in this study reported an increase in perceived lateral translation during passive roll rotation after flight ([Reschke & Parker 1987](#)).

On the basis of these observations, and similar ones reported by [Young et al. \(1984\)](#), the otolith tilt-translation reinterpretation hypothesis (OTTR) was proposed. The OTTR hypothesis is based on the premise that interpreting otolith signals as indicating tilt is inappropriate during space flight ([see Section IV, 2.4.6](#)). Therefore, during adaptation to weightlessness, the brain reinterprets otolith signals as indicating translation only. Relevant to driving tasks on sloped terrains, it is interesting to note that performance during roll-tilt closed-loop nulling tasks is

decreased for several days post-flight (Merfeld 1996a), while performance during translation closed-loop nulling experiments appears to be improved (Arrott *et al.* 1990).

An alternative hypothesis proposed by Guedry *et al.* (1998) suggests that rather than a reinterpretation of otolith signals, adaptation to space flight might involve ‘shutting down’ the search for position (tilt) signals from the otolith system in order to avoid vestibular conflict. This is based on the observation that on Earth the initial head position relative to gravity before a head turn foretells the unique combination of canal and otolith signals that will occur during the turn. The absence of a meaningful initial position signal from the otoliths on orbit may therefore be functionally disruptive, and eventually neglected. Guedry’s hypothesis also explains the post-flight tilt-translation disruptions described above, as well as the increased immunity to Coriolis stimuli observed following the Skylab missions (Graybiel *et al.* 1977).

Differences between active and passive motions may help explain some of the apparently contradictory observations regarding post-flight tilt-translation disturbances. For example, Golding *et al.* (2003) observed striking differences in motion sickness sensitivity between active and passive tilts. It is likely that the new ‘expected’ patterns of sensory cues adopted during head tilts on orbit will differentially influence responses during reentry depending on whether the motion is self-generated.

Merfeld (2003) noted that the OTTR hypothesis assumes that the utricular otolith mediates all tilt sensation, and that if otolith cues were simply reinterpreted as linear acceleration, a sustained head tilt should produce a sustained acceleration sensation—not what is usually observed. He hypothesized that both types of illusions could result from a change in the effect of semicircular canal cues on estimating transient rotations of the direction of “down” relative to the head. Unless the CNS estimate of angular velocity is aligned with the estimated direction of gravity, a conflict occurs. His hypothesis, known as the Rotation Otolith Tilt-Translation Reinterpretation (ROTTR) hypothesis, suggests that the CNS resolves this conflict by rotating the direction of its internal estimate of gravity at a rate proportional to the vector cross product of the estimated angular velocity and gravity vectors. These rate constants determined the dynamics of the resulting illusion (Merfeld *et al.* 1993).

Tilt-translation illusions can occur during spacecraft pitching or rolling maneuvers, even if the pilot’s head remains stationary relative to the cockpit, and could lead to an incorrect manual control responses. For example, a Tilt Gain illusion might result in an under-response to a Shuttle wing drop, a sensation that a wind gust was pushing the nose up unexpectedly, resulting in under-rotation during the critical landing flare maneuver. An OTTR illusion might produce an over-response to a wing-drop, and perhaps the sensation that a gust had suddenly pushed the Shuttle off runway centerline. One implication of ROTTR theory is that the tendency toward Tilt Gain or OTTR illusions may be a personal characteristic. If so this could account for the diversity in the anecdotal descriptions by astronauts.

Some evidence exists that provides insight into the physiological basis of these illusions. For example, in a series of rodent experiments, Ross (1993, 1994, 2000, 2003) showed increased numbers of synapses in type II hair cells of the utricular maculae during and just after

space flight. The findings of increased synaptic plasticity are consistent with the human behavioral studies suggesting an increased gain of the otolith organs. These findings were also supported by an experiment performed by [Boyle et al. \(2001\)](#) aboard the Neurolab mission, in which the primary utricular afferent information was shown to be highly potentiated (up-regulated) during the first few hours after space flight in oyster toadfish (*Opsanus tau*) subjected to linear translations in various planes. These data were similar to those reported by [Reschke et al. \(1986\)](#), who found an enormous potentiation of the monosynaptic Hoffman reflex response early after flight in human subjects from the Spacelab-1 mission subjected to linear translational acceleration stimuli. This Hoffman reflex response, which is modulated by descending signals from the vestibular otolith organs and normally aids in preparing the anti-gravity muscles for stable landing following a jump (or fall), had completely disappeared in these same subjects by the sixth flight day of the mission.

Further evidence was obtained by [Holstein & Martinelli \(2003\)](#), who found in rodents flown aboard the Neurolab mission ultrastructural signs of plasticity in the otolith recipient zone of cerebellar cortex (nodulus), an area thought to be critical for motor control, coordination, timing of movements, and motor learning. Rats flown for 5-18 days in the Russian Cosmos Biosatellite Program also showed morphological changes in neural structure, including decreased lengths in dendrites directed from cells in the reticular formation toward structures in the vestibular nuclei and morphological changes in cerebellar structures including mossy fiber terminals in the granular layer of the nodular cortex ([Krasnov 1994](#)). [Pompeiano \(2003\)](#) also studied rodents flown aboard the Neurolab mission. He found biochemical evidence of plasticity (expression of the immediate early gene c-fos and presence of fos-related antigens) in multiple regions of the brain, including the vestibular nuclei, which play a role in controlling posture and eye movements, the nucleus of the tractus solitarius (NTS), which is involved in regulation of cardiovascular and respiratory function, the area postrema, which plays a role in motion sickness, the amygdala, cortical and subcortical areas involved in body orientation and perception, and the locus coeruleus, which is involved in regulation of the sleep-wake cycle.

#### **3.4.4. Otolith Asymmetry**


Another underlying mechanisms that may contribute to inflight and postflight disturbances in perception and spatial orientation is natural asymmetry in otoconial mass between the right and left saccule and utricle. These asymmetries would lead to different output signals from the left and right otoliths. On Earth the CNS is thought to compensate for this inherent imbalance between otoliths. However, exposure to a weightless environment may lead to decompensation of this process leading to asymmetrical output from the right and left otolith organs and subsequent disturbances in perception of motion and spatial orientation ([Clarke et al. 2013](#)). A study was conducted using Shuttle crewmembers that examined otolith asymmetries as a potential underlying factor leading contributing to inflight and postflight perceptual and motor control disturbances ([Clarke et al. 2010](#)). The study utilized the vestibular evoked myogenic potentials (VEMP) as an indicator of otolith saccular function through measurement of a vestibulo-collic reflex. Results showed a general increase in asymmetry of otolith responses on landing day relative to the preflight baseline data. There was a subsequent reversal in asymmetry within 2-3 days. Recovery back to preflight levels occurred within the

first week following the return to Earth. These findings indicate that space flight results in adaptive changes in neural integration of otolith inputs contributing to perceptual illusions during gravitational transitions.

### 3.5. Gaze Changes

#### 3.5.1. Vestibulo-Ocular Reflex

Evidence exists that the VOR could be severely compromised during the transition between low Earth orbit and the return to Earth's gravity. Using a gaze stabilization protocol, eye and head movements were recorded from crewmembers inside of the Landing and Entry Helmet during descent (Reschke *et al.* 1999). This protocol required the subject to view a target presented to the subject at a distance of approximately 0.5 m from the eyes. Immediately upon target presentation, vision was occluded and the subject acquired the remembered target. One second later the target was exposed and the subject made any correction to the target with only the eyes. This protocol permitted the determination of the VOR gain. Interestingly, the data showed that the VOR is not functional immediately after landing (Figure 7).



*Figure 7. Gaze stabilization paradigm. The VOR, nearly perfect preflight, is disrupted during entry. The eye is apparently locked with the head rather than moving equally and opposite the head movement (stabilizing vision). Approximately one hour after landing (Postflight) the VOR has returned to normal.*

Experiments during the D-1, SLS-1, and SLS-2 Spacelab missions utilized passive body movements provided by step changes in the angular velocity of rotating chairs to stimulate the VOR. During parabolic flight, the persistence of the yaw VOR response after the chair motion stopped was decreased in eight astronauts tested just before space flight (Oman *et al.* 1993, 1996) and in normal subjects (DiZio & Lackner 1988). However, after 4-10 days in orbital flight, the yaw VOR persistence was no different from preflight values in five of the eight astronauts tested, although active head pitch movements (“dumping”) did not interfere with the VOR persistence, as it consistently did on Earth. Early after flight (1-2 days), the persistence was decreased relative to preflight in nine of 12 astronauts tested, but it eventually returned to preflight values in all (Oman *et al.* 1988). These findings suggest that transitions to and from weightlessness temporarily reduce the contribution of brainstem mechanisms that normally extend the low frequency bandwidth of the human angular VOR response.

### 3.5.2. Ocular Counterrolling

The otoliths, detectors of linear acceleration, contribute to eye stabilization and spatial orientation during space flight. In the absence of gravity, head tilt has insignificant meaning, and therefore, torsional eye movements should be minimal or absent. However, statolith input may be reinterpreted. As a result, ocular counter-rolling (OCR) induced by tilt has been investigated before and after space flight as an indicator of otolith adaptation. Head tilt after Gemini missions induced no changes in OCR (Graybiel *et al.* 1967).

Clément *et al.* (2007) have reviewed the evidence of changes in OCR during static whole body tilt in postflight studies following Space Shuttle missions, and they noted that the findings are inconsistent. Some studies report decreases in astronauts' OCR after the flight relative to preflight, while others have shown postflight increases of OCR or no changes at all (Reschke *et al.* 1985; Vogel & Kass 1986; Young & Sinah 1998; Diamond & Markham 1998; Hofstetter-Degen *et al.* 1993; Moore *et al.* 2001; Clarke & Kornilova 2007; Kornilova *et al.* 2011). When averaged across all these space studies, the difference between pre- and postflight OCR measurements was found to be less than 0.6° for body tilt angles ranging from 15-45° (Figure 8). The inconsistency in these results may be due to the various experimental procedures employed, including flash afterimages, flash photography of the eyes, or video-oculography, and to the high variance of the OCR across individuals.




Figure 8. A. Individual (open circles) and averaged (closed circles) OCR amplitude during static whole body tilt recorded preflight in 18 astronauts. B. Differences between pre- and postflight OCR measurements (Clément *et al.* 2007).

### 3.5.3. Eye Movements during Coriolis and Cross-Coupled Acceleration

As part of the Human Vestibular Function (M-131) study during the three Skylab manned missions, eight crewmembers performed head and body movements during yaw rotation ranging from 12.5-30 rpm (Graybiel *et al.* 1977). This stimulus generated Coriolis, cross-coupled accelerations, which were at the origin of motion sickness preflight. All crewmembers could perform more head body movements and had reduced motion sickness from flight day 6 and

beyond. In addition, 7 out of the 8 crewmembers had reduced motion sickness postflight from the day after landing (R+1) through R+17. The remaining crewmember had mild and severe MS for two of the 3 days at sea following splash-down. The other crewmembers had taken anti-motion sickness medication taken during recovery and no motion sickness.

Other experiments demonstrated that most subjects perceive passive vestibular stimulation to be less provocative on landing day than preflight. For example, 9 of 10 crewmembers tested had no symptoms of motion sickness when exposed to passive yaw rotation or Coriolis acceleration on R+0 (Thornton *et al.* 1987a; Oman *et al.* 1996; Clément *et al.* 1999). In those experiments using passive rotation on R+1 and later, 26 out of 29 crewmembers showed a decreased of susceptibility to MS postflight relative to preflight (Graybiel *et al.* 1977; Wetzig *et al.* 1993; Harm *et al.* 1994; Moore *et al.* 2001). One crewmember was tested using Coriolis and off vertical axis rotation (OVAR) following the 10-day Apollo-9 mission. Reports indicated that the single subject tested had an average susceptibility to motion sickness preflight and that there was an increase tolerance with repeated exposures postflight (Homick & Miller 1975). In another study, 58 crewmembers participating in 16 flights of the Space Shuttle (duration < 11 days) were also tested during Coriolis (12.5-30 rpm) and during OVAR (20 rpm up to 30° tilt) before (L-3 to L-6 months) and after (R+0 to R+3 months) their space flight. This experiment focused on predictive values of ground tests (Homick *et al.* 1987). Two crewmembers tested immediately postflight showed reduced motion sickness to Coriolis (up to several weeks), although one subject was found to be hypersensitive on landing day (Thornton *et al.* 1987a). Past experiments indicate that velocity storage (Cohen *et al.* 1977), the central integration of semicircular canal signals, is attenuated during exposure to nonterrestrial gravitoinertial force backgrounds and that this effect carries over to the postflight period (DiZio & Lackner 1988; Oman & Balkwill 1993; Oman *et al.* 1996). Converging evidence from ground-based and space flight experiment also points to a relation between motion sickness and the properties of the velocity storage (DiZio & Lackner 1991; Oman 1998; Dai *et al.* 1996, 2010). Perception of the vertical and of body angular movements indicate that spatial integration of canal afferent signals is disturbed after adaptation to microgravity (Clément *et al.* 2007, 2001), suggesting that velocity storage is reduced.

#### **3.5.4. Eye movements during Linear Acceleration**

Two subjects exposed to transient lateral acceleration 3-5 hours after the landing of the SL-1 mission demonstrated smaller torsional amplitudes than three of the four preflight measures of sinusoidal ocular torsion for these subjects. Torsional amplitude in these subjects steadily increased over most of the post-flight tests, but changes were not statistically significant because of high variability in the preflight measurements (Arrott & Young 1986). Y-axis linear translation enhanced horizontal eye movements in crewmembers 2-3 days after landing (Parker *et al.* 1986). These results are consistent with the OTTR hypothesis, which predicts reduced eye torsion immediately after landing due to reinterpretation of otolith signals as linear translation (Parker *et al.* 1985; Liao *et al.* 2011).

In two recent studies, eye movements and perceived motion were evaluated in astronauts returning from Space Shuttle missions during OVAR (Clément & Wood 2013) and during short-


radius centrifugation (Clément & Wood 2014). No changes were seen on the compensatory eye movements to these linear accelerations between pre- and post-flight. However, the crewmembers reported an overestimation of the sensation of roll tilt (but not pitch tilt) and an overestimation of the sensation of translation immediately post-flight. These results confirm that some VORs elicited during passive motion may not be altered by short-duration space flight, or may readapt very quickly, and that the resolution of sensory conflict associated with post-flight recovery involves higher-order neural processes (Holly *et al.* 2010).

### **3.6. Decrements in Functional Performance**

#### **3.6.1. Functional Mobility**

Pre- and postflight functional mobility assessment was performed on ISS crewmembers to determine their ability to complete challenging locomotor maneuvers similar to those encountered during an egress from a space vehicle (Mulavara *et al.* 2010). To perform the Functional Mobility Test (FMT) subjects walked at a self-selected pace through an obstacle course set up on a base of medium density foam that increased the challenge of the test. There was a 48% increase in time to traverse the course one day after landing, and recovery of function took an average of 15 days to return to within 95% of their preflight level of performance (Figure 9).

The results also showed that post-flight recovery can be divided into two processes: rapid strategic learning over the six trials on the first day after return, and a slower process taking over 2 weeks to recover to a pre-flight level of performance. It is believed that training can promote or enable these strategic or early learning responses for facilitating faster re-adaptation to Earth's 1-g environment on return from space flight. Additionally, a significant positive correlation between measures of long-term recovery and early motor learning (strategic learning) indicating that the two types of recovery processes influence an astronaut's ability to re-adapt to Earth's gravity environment. Early motor learning helps astronauts make rapid modifications in their motor control strategies during the first hours after landing. Further, this early motor learning appears to reinforce the adaptive realignment, facilitating re-adaptation to Earth's 1-g environment on return from space flight.



**Figure 9.** Left: scatter plot of time to complete the course (TCC) for 18 long-duration subjects showing a 48% increase in time to traverse the obstacle course one day after landing. Right: Recovery of function took an average of 15 days to return to within 95% of their preflight level of performance.

Adaptation to different gravitational environments may affect planetary extravehicular activities during the initial adaptation period due to postural and locomotor dysfunction. These alterations may also lead to decrement in the ability to multi-task along with increasing the metabolic costs of ambulation. A recent study examined subjects walking on a destabilizing support surface using a treadmill mounted on a six-degree-of-freedom motion base that provided an oscillating support-surface during walking (Peters *et al.* 2013b). Results demonstrated that measures of locomotor stability, cognitive load, and metabolic cost were all significantly greater during support-surface motion than during baseline walking conditions and showed a trend toward recovery to baseline levels during locomotor adaptation. These decrements are operationally meaningful because they indicate broader functional implications for postflight locomotor instability (Brady *et al.* 2009, 2012). Until recently, locomotor adaptation to discordant sensory conditions has been characterized primarily in terms of impact on the underlying mechanisms contributing to locomotor stability. These results indicate that uncoordinated walking during periods of adaptive change may also come at significant cognitive and metabolic costs to the crew. Cognitive load increases and metabolic cost rises because of new demands on attention and additional physical work required to maintain balance while walking. Energetic cost is a key contributor to the duration and intensity of EVAs performed by suited astronauts, and previous research on suited locomotion has explored the effects of load, slope, and walking vs. running (Carr & Newman 2007a, 2007b). Metabolic cost associated with locomotor instability is a factor that should be accounted for in the continuing efforts to improve extravehicular suit design.

### 3.6.2. Functional Tasks

To understand how changes in physiological function affect functional performance, an interdisciplinary pre- and post-flight testing regimen, Functional Task Test (FTT), was developed

to systematically evaluate both astronaut functional performance and related physiological changes (Spiering *et al.* 2011; Arzeno *et al.* 2013; Ryder *et al.* 2013; Bloomberg *et al.* 2015a). Ultimately this information will be used to assess performance risks and inform the design of countermeasures for exploration class missions. This FTT study was conducted on Shuttle and ISS crewmembers before and after 6-month expeditions and is currently being conducted for 1-year expeditions. Additionally, in a corresponding study, the FTT protocol was used on subjects before and after 70 days of 6° head-down bed-rest as an analog for space flight. Bed-rest provides the opportunity to investigate the role of prolonged axial body unloading in isolation from the other physiological effects produced by exposure to microgravity (Reschke *et al.* 2009). Therefore, the bed rest analog allowed the investigation of the impact of body unloading on both functional tasks and on the underlying physiological factors that lead to decrement in performance and then compare those with the results obtained in the space flight study.

Functional tests included ladder climbing, hatch opening, jump down, manual manipulation of objects and tool use, seat egress and obstacle avoidance, recovery from a fall, and object translation tasks (Bloomberg *et al.* 2015a). Physiological measures included assessments of postural and gait control, dynamic visual acuity, fine motor control, plasma volume, heart rate, blood pressure, orthostatic intolerance, upper- and lower-body muscle strength, power, endurance, control, and neuromuscular drive. ISS crewmembers were tested three times before flight, and 1, 6, and 30 days after landing. Subjects in bed rest studies were tested three times before bed rest and immediately after getting up from bed rest as well as 1, 6, and 12 days after reambulation.

Astronaut data showed that functional tests requiring a greater demand for dynamic control of postural equilibrium (i.e., fall recovery, seat egress and obstacle avoidance during walking, object translation, jump down) showed the greatest decrements in performance (Bloomberg *et al.* 2015a). Functional tests with reduced requirements for dynamic postural stability (i.e., hatch opening, ladder climb, manual manipulation of objects and tool use) showed less reduction in performance. Similarly, subjects exposed to prolonged bed rest (> 20 days) showed the same trends in performance change as astronauts, namely a reduction in performance on functional tests requiring a greater demand for dynamic control of postural equilibrium. For both spaceflight and bed rest subjects these changes in functional performance were paralleled by similar decrements in physiological tests designed to specifically assess postural equilibrium and dynamic gait control.

Taken together, the space flight and bed rest results indicate that the unloading of body support structures (major postural muscles) that is experienced during space flight plays a larger role than expected in post-flight alteration of functional task performance and balance control. These data point to the importance of providing significant axial body loading during in-flight treadmill and resistive exercise. In addition, these data indicate that balance training should be used to supplement current in-flight aerobic and resistive exercise activities.

### 3.6.3. Field Test

Testing of crew responses following long-duration flights has not been previously possible until a minimum of +24 hours after landing. As a result, it has not been possible to determine the trend of the early recovery process, nor has it been possible to accurately assess the full impact of the decrements associated with long-duration flight. To overcome these limitations, both the Russian and U.S. programs have implemented joint testing at the Soyuz landing site. This ISS research effort has been identified as the Field Test, and represents data collect on NASA, Russian, ESA, and JAXA crews.

The primary goal of this research is to determine functional abilities associated with long-duration space flight crews beginning as soon after landing as possible on the day of landing (typically within 1 to 1.5 hr). This goal has both sensorimotor and cardiovascular elements. To date, a total of 15 subjects have participated in a 'pilot' version of the full 'field test'. The full version of the 'field test' will assess functional sensorimotor measurements included hand/eye coordination, standing from a seated position (sit-to-stand), walking normally without falling, measurement of dynamic visual acuity, discriminating different forces generated with the hands (both strength and ability to judge just noticeable differences of force), standing from a prone position, coordinated walking involving tandem heel-to-toe placement (tested with eyes both closed and open), walking normally while avoiding obstacles of differing heights, and determining postural ataxia while standing (measurement of quiet stance). A number of these tests have been utilized and tested for sensitivity and specificity in a relatively large cohort of normal and clinical patients populations ([Cohen \*et al.\* 2012a, 2012b, 2013, 2014](#); [Peters \*et al.\* 2012b](#); [Peters \*et al.\* 2013a](#), [Mulavara \*et al.\* 2013](#))

Sensorimotor performance has been obtained using video records, and data from body worn inertial sensors. The cardiovascular portion of the investigation has measured blood pressure and heart rate during a timed stand test in conjunction with postural ataxia testing (quiet stance sway) as well as cardiovascular responses during sensorimotor testing on all of the above measures. Motion sickness data associated with each of the postflight tests has also been collected. When possible a rudimentary cerebellar assessment was undertaken. In addition to the immediate postlanding collection of data, postflight data has been acquired twice more within 24 hours after landing and measurements continue until sensorimotor and cardiovascular responses have returned to preflight normative values (approximately 60 days postflight).

The level of functional deficit observed in the crew tested to date is more severe than expected, clearly triggered by the return to gravity loads immediately after landing when the demands for crew intervention in response to emergency operations will be greatest. Measureable performance parameters such as ability to perform a seat egress, recover from a fall or the ability to see clearly when walking, and related physiologic data (orthostatic responses) are required to provide an evidence base for characterizing programmatic risks and the degree of variability among crewmembers for exploration missions where the crew will be unassisted after landing. Overall, these early functional and related physiologic measurements

will allow the estimation of nonlinear sensorimotor and cardiovascular recovery trends that have not been previously captured ([Reschke et al. 2015](#)).

#### **3.6.4. Navigation**

To quantify performance in orienting during free walking after space flight, astronaut subjects were asked to walk, preflight and post-flight, a previously seen triangular path with normal vision and vision occluded ([Glasauer et al. 1995](#)). The path, marked on the ground by a cross at each corner, consisted of a right triangle with two legs 3-m long. The trajectories of three infrared-reflective markers fixed on a helmet were recorded using a video-based motion analysis system. Subjects showed inter-individual differences, especially for directional deviations from the path, in the vision-occluded condition even preflight; the characteristics of these differences persisted throughout all experimental sessions. The absolute directional errors turned out to be larger post-flight, which means that subjects had larger directional errors but in different directions. In the post-flight condition, however, there was a trend to a larger underestimation of the angle turned at each corner. In contrast to directional errors, the length of the legs walked was similar pre- and post-flight. These data suggest that the perception of self-displacement during turning, but not during linear motion, was changed by the stay in weightlessness. A possible explanation could be the development of a mismatch between information from otoliths and semicircular canals during whole-body turns in microgravity. This change in canal-otolith interaction may underlie the disturbances in locomotion experienced by returning astronauts.

#### **3.7. Decrements in Manual Control**

In studies performed immediately after two Spacelab missions, returning astronauts were seated on a rail-mounted sled, and asked to use a joystick to null a random linear disturbance movement along their interaural ([Arrott & Young 1986](#)) and/or longitudinal ([Merfeld et al. 1996b](#)) body axes. Four of the seven subjects tested showed improved post-flight performance on the nulling task. Also, [Merfeld et al. \(1996a\)](#) tested the early post-flight performance of astronauts trying to maintain a flight simulator in an upright orientation in the presence of pseudorandom motion disturbances about a tilt axis located below their seat. On landing day, both subjects showed impaired ability to control their tilt in the dark, but displayed normal responses when visual motion cues were provided. Results confirm that returning crews have difficulty estimating their tilt orientation with respect to the gravitational vertical on landing day. The absence of change with visual cues shows that neuromuscular and fatigue factors were not major contributors to the effect. It is important to note that the subjects in these experiments all knew whether tilt or translation motions were possible. Subsequent ground-based experiments ([Park et al. 2006](#); [Wetheim et al. 2001](#)) showed that when subjects must resolve tilt-translation ambiguities, and are naïve to the possible motion, large misperceptions of tilt could result.

#### **3.8. Impact of Vestibular Changes to Orthostatic Intolerance**

Historically, factors that have been thought to contribute to post-space flight orthostatic intolerance (PSOI) include space flight-related volume depletion ([Fischer et al. 1967](#); [Johnson et](#)

*al.* 1977; Bungo *et al.* 1985; Charles & Lathers 1991) and excessive venous pooling in the lower extremities (Johnson *et al.* 1976; Buckey *et al.* 1992) or splanchnic circulation. Later findings, however, suggest that abnormalities in autonomic cardiovascular control, including a loss of carotid-cardiac baroreflex range and slope during and after space flight (Fritsch *et al.* 1992; Fritsch-Yelle *et al.* 1994) and a deficit in peripheral vasoconstriction in the upright position on landing day may play a primary role in PSOI (Buckey *et al.* 1996; Fritsch-Yelle *et al.* 1996). These autonomic factors were reemphasized given the finding that little correlation exists between deficits in plasma volume and deficits in orthostatic tolerance in returning astronauts. The mechanisms underlying the autonomic abnormalities associated with PSOI, however, remain poorly understood.

Over the past several years, much evidence has accumulated to suggest that the vestibular organs can provide “-” signals that supplement feedback information provided by baroreceptors in maintaining orthostatic tolerance (Doba & Reis 1974; Gillingham *et al.* 1977, Essandoh *et al.* 1988; Satake *et al.* 1991; Yates 1992; Previc 1993; Yates *et al.* 1993; Convertino *et al.* 1997; Ray *et al.* 1997; Shortt & Ray 1997; Woodring *et al.* 1997; Yates & Kerman 1998; Serrador *et al.* 2009a, Serrador *et al.* 2009b). The neuroanatomic and neurophysiologic bases for vestibular contributions to autonomic cardiovascular control were reviewed thoroughly by Yates and colleagues (Yates 1992, 1998; Yates & Kerman 1998; Yates & Miller 1996, 1998; Yates *et al.* 1998).

The following, taken together, provide strong evidence that changes in vestibular (and especially otolith) function during space flight contribute to PSOI: (1) changes in otolith function (Vogel & Kass 1986; Young *et al.* 1986) and structure (Ross 1993, 1994) have been described and verified in multiple studies during and after space flight; (2) signals from intact otolith organs clearly contribute to sympathetically mediated peripheral vasoconstriction in animals (Doba & Reis 1974; Woodring *et al.* 1997; Yates 1992, 1998; Yates & Kerman 1998; Yates & Miller 1998) and likely contribute to sympathetically mediated peripheral vasoconstriction in humans (Essandoh *et al.* 1988; Normand *et al.* 1997; Ray *et al.* 1997; Shortt & Ray 1997); and (3) the key autonomic defect associated with PSOI in most returning crewmembers is inadequate sympathetically mediated peripheral vasoconstriction (Buckey *et al.* 1996; Fritsch-Yelle *et al.* 1996). In spite of all of this evidence, however, the precise role of vestibular-autonomic factors in PSOI remains to be defined. Further ground-based studies using both labyrinthine-deficient and -intact humans and animals will be required before any definitive treatment or prophylactic regimens can be designed based on an assumption of underlying vestibular-autonomic pathology.

## V. COUNTERMEASURES

### 1. PHARMACOLOGICAL COUNTERMEASURES

Many drugs have been tested for their effectiveness against motion sickness. Although some drugs have proven somewhat effective, no drug or drug combination has been identified that protects all individuals.

#### 1.1. Drug Classes

The mechanism(s) of action of the effective anti-motion sickness drugs is unclear, but it has been noted that the action(s) responsible for their anti-motion sickness efficacy tends to differ from the drug's primary action ([Money 1970](#)). Numerous studies have found scopolamine, an anticholinergic (parasympatholytic) drug, to be effective in treating motion sickness ([Wood & Graybiel 1968, 1970, 1972](#); [Graybiel et al. 1976, 1981](#); [Graybiel & Knepton 1977](#); [McCauley et al. 1979](#); [Laitinen et al. 1981](#); [Shashkov & Sabayev 1981](#); [Homick et al. 1983](#); [Noy et al. 1984](#); [Levy & Rapaport 1985](#); [Pyykkö et al. 1985](#); [van Marion et al. 1985](#); [Grigoriev et al. 1986](#); [Offenloch et al. 1986](#); [Wood et al. 1986, 1987a](#); [Attias et al. 1987](#); [Graybiel & Lackner 1987](#); [How et al. 1988](#); [Karkishchenko 1989](#); [Shupak et al. 1989](#); [Woodard et al. 2014](#)). Although most of the antihistamines tested for anti-motion sickness properties have had some benefit, they tend to be less effective than scopolamine. Promethazine, the most effective of the antihistamines, approaches scopolamine in efficacy ([Wood & Graybiel 1972](#)). The few sympatholytic drugs that are effective against motion sickness are of marginal benefit and have less effect than the least effective antihistamine ([Wood & Graybiel 1968](#)). The combination of a parasympatholytic drug (scopolamine) and a sympathomimetic has also been more effective than these classes of drugs taken alone.

Anti-motion sickness drug research has been reviewed by [Wood \(1979, 1990\)](#). The vast majority of anti-motion sickness drugs have been given orally. However, the complications of oral medications force frequent dosing and the use of secondary medications via other routes of administration to overcome decreased absorption ([Chess et al. 1975](#); [Reason & Brand 1975](#); [Graybiel et al. 1976](#); [McCauley et al. 1979](#); [Graybiel et al. 1981](#); [Homick et al. 1983](#); [Becker et al. 1984](#); [Levy & Rapaport 1985](#); [Pyykkö et al. 1985](#); [van Marion et al. 1985](#); [Offenloch et al. 1986](#); [Attias et al. 1987](#); [Graybiel & Lackner 1987](#); [Wood et al. 1987b](#); [How et al. 1988](#); [Bagian 1991](#); [Davis et al. 1993a, 1993b](#)). The occurrence of side effects also precludes the effective use of many of these drugs. Two newer classes of antiemetic drugs, 5HT<sub>3</sub> and NK<sub>1</sub> receptor antagonists, may be promising candidates for the treatment of motion sickness ([Stott et al. 1989](#); [Koch et al. 1994](#); [Gardner et al. 1995](#)).

#### 1.2. In-Flight Medication

A reported 28–30% of all Shuttle crewmembers received medication for relief of SMS during flight ([Santy & Bungo 1991](#)). In the U.S. program, scopolamine, scopolamine with dextroamphetamine (scop-dex), promethazine, promethazine with ephedrine, metoclopramide, naloxone, and Compazine have all been used to treat SMS, with varying degrees of success ([Graybiel 1980](#); [Thornton et al. 1987a](#); [Davis et al. 1993a, 1993b](#); [Bagian &](#)

[Ward 1994](#)). Reports are available on the prophylactic and in-flight use of scop-dex taken orally ([Davis et al. 1993a](#)). Only 3 of 19 crewmembers who took this medication experienced no SMS symptoms.

Intramuscular (IM) promethazine has been used successfully to treat SMS symptoms. [Davis et al. \(1993b\)](#) reported that of 20 crewmembers given 25–50 mg of promethazine IM (dose adjusted for body weight) on flight day 1, 25% were still classified as “sick” on the second flight day. In contrast, 50% of the 74 crewmembers reporting SMS on the first day of flight who did not receive promethazine, or received other anti-motion sickness medications, were still “sick” on flight day 2. Ninety percent of those who received IM promethazine reported relief from SMS symptoms within 1 to 2 hours of dosing and only three crewmembers needed a second dose. Three of the IM promethazine recipients reported drowsiness after administration, but the injection is often given immediately before the sleep period. An IM injection of 25–50 mg of promethazine is now the recommended treatment for moderate to severe cases of SMS in the U.S. space program, whereas oral and suppository routes are used for less severe symptoms. Some crewmembers have taken a prophylactic combination of promethazine and dextroamphetamine before launch.

The variable success of anti-motion sickness drugs administered during flight may be due to changes in drug absorption or metabolism by factors such as dehydration, reduced gastrointestinal motility, changes in body chemistry, changes in cabin pressure, and disruption of normal sleep/wake cycles. The concomitant administration of medications for other indications is another confounding factor ([Santy & Bungo 1991](#)). The success of IM promethazine administration is encouraging. Questions still remain, however, as to whether the effectiveness of promethazine is due to its pharmacologic effect or its route of administration ([Bagian 1991](#)).

## **2. TRAINING AND REHABILITATION**

### **2.1 Vestibular Training**

Vestibular training has been used in attempts to prevent or control the symptoms of SMS. Vestibular training techniques investigated thus far have been based on one of two suppositions: (a) adaptation to stressful motion can be hastened through previous exposure to conflicting sensory inputs; or (b) symptoms can be avoided by learned control of autonomic responses.

[Lapayev and Vorobyev \(1986\)](#) have hypothesized that motion sickness susceptibility is proportional to the ratio of signals from the vestibular system and other sensory (e.g., visual, proprioceptive) systems. They propose that the most effective method of increasing tolerance to motion sickness is to train the vestibular system while stimulating the other senses. The Russian space program primarily uses Coriolis and cross-coupled angular acceleration as preflight vestibular training. However, this method does not duplicate the sensory conflicts or fluid shifts encountered in weightlessness. [Aizikov et al. \(1991\)](#) observed that using a

predetermined sequence of muscle tension and relaxation increased tolerance to experimentally induced motion sickness by reducing the number of symptoms and shortening the recovery time. These investigators theorize that the afferent propriotactile information generated by the muscle-tension regimen provides enough information on body position to override coincident, possibly inaccurate, vestibular information.

## 2.2. Preflight Adaptation Training

Training devices and procedures to adapt astronauts to the sensory-stimulus rearrangements of microgravity before flight are being developed in the US. Preflight adaptation training is based on the following postulates: (a) that microgravity rearranges sensory stimuli and astronauts adapt to the rearranged stimuli (sensory conflict theory); (b) that adaptation may result from sensory compensation, reinterpretation of stimuli, or both (sensory compensation and OTTR hypotheses) (see Section IV, 2.4.6); and (c) that people can learn and store perceptual, sensory, and sensorimotor responses appropriate to different sensory stimulus conditions, and can learn to invoke these alternative responses when necessary.

Two training devices are used to provide a variety of stimulus rearrangements and train sensorimotor reflexes: the device for orientation and motion environments (DOME) that achieves graviceptor stabilization, and the tilt-translation device (TTD) that produces graviceptor-visual rearrangement. Theoretically, training with these devices would produce the necessary responses to weightlessness and for the return to a 1-g environment, such as compensatory eye movements, postural-muscle reflexes, and self-motion and orientation experiences in relation to visual scene movements.

Ground-based studies using the TTD trainer have revealed that a 270° phase relation between tilt and surround-motion in the TTD best supports reinterpretation of otolith signals as linear translation (Reschke *et al.* 1988; Harm & Parker 1993, 1994). Exposure to this profile also results in decreased compensatory eye movement gain, net gaze compensation, and decreased postural stability (Michaud *et al.* 1989; Paloski *et al.* 1990; Harm *et al.* 1991; Harm & Parker 1993). These results are consistent with the OTTR model of sensory adaptation and are consistent with observations of astronauts and cosmonauts during or after flight.

Flight investigations involving these training devices thus far have focused on providing an experience of sensory rearrangement that results in illusions of linear or angular self- or surround-motion. The primary purpose of these studies is to teach astronauts to describe perceptual phenomena systematically by using a quantitative “motion perception vocabulary” related to anatomy and physiology so that they can properly describe perceptual illusions.

Post-flight training on the TTD has provided evidence that the simulated perceptual experiences are similar to those experienced in-flight. SMS symptoms and visual disturbances have been reelicited, and perceptual illusions of linear self-motion and tilt angle are intensified relative to preflight stimulation (Harm & Parker 1993; Harm *et al.* 1999). These results generally support the OTTR model of sensory adaptation to microgravity and suggest that the training devices can simulate the appropriate sensory rearrangements. Moreover, crewmembers who

participated in this study showed an average 33.5% improvement in SMS symptoms compared with those who did not participate ([Harm et al. 1999](#)).

### **2.3. Sensorimotor Adaptability Training**

Research is currently being conducted to develop a comprehensive sensorimotor adaptability (SA) training program as a countermeasure to facilitate rapid adaptation to novel gravitational environments and readaptation to Earth's gravity ([Bloomberg et al. 2015b](#)). The human brain is highly adaptable, enabling individuals to modify their behavior to match the prevailing environment. It has been previously shown that subjects trained to adapt to varied sensorimotor challenges can adapt faster to new sensory environments that they have never experienced before ([Roller et al. 2001](#); [Cohen et al. 2005](#); [Mulavara et al. 2009](#); [Roller et al. 2009](#); [Seidler et al. 2006](#));). This is a process known as adaptive generalization that allows you to enhance the ability to “learn how to learn” to adapt to novel environments ([Krakauer, 2006](#); [Seidler 2010](#); [Bloomberg et al. 2015b](#)). This learning paradigm is similar to structural learning in motor systems that use abstract learning strategies that are applicable to a wide range of novel environments ([Braun et al. 2010](#)).

By applying these motor-learning concepts for training astronauts these training programs can enhance their ability to rapidly adapt their behavioral responses following a gravitational transition. To minimize cost and demands on crew time current training concepts are integrated SA training with existing exercise activities, namely treadmill walking. The SA training program being developed entails manipulating the sensory conditions of treadmill exercise to systematically and simultaneously challenge multiple sensorimotor systems while conducting nominal exercise activities. To provide SA training investigators have mounted a treadmill on a six degree-of-freedom motion base to produce variation in the support surface offering subjects balance challenges during walking ([Peters et al. 2012](#); [Peters et al. 2013](#); [Brady et al. 2009](#)). Additional sensorimotor challenges are provided by exposing subjects to variation in visual input during walking using a projected virtual scene that produces variation in visual flow ([Richards et al. 2004](#); [Mulavara et al. 2005](#); [Nomura et al. 2005](#); [Richards et al. 2007](#); [Bucello-Stout et al. 2008](#); [Brady et al. 2012](#)). [Bloomberg et al. \(2015b\)](#) have reviewed several studies that have shown that subjects who received SA training adapted faster than controls when presented with a novel discordant sensory environment because they were able to apply adaptive skills that were learned during their earlier training sessions. Importantly, the training improved performance across a number of modalities including enhanced locomotor function and increased multi-tasking capability when walking in a novel, discordant sensory environment not previously experienced by the subjects. This improved performance could be retained over a 6-month period and perhaps longer, indicating that a component of this training could take place before long-duration missions ([Batson et al. 2011](#)).

Facilitating adaptation to altered gravitational environments using treadmill walking while experiencing discordant sensory input in 1-G has been targeted as a potential countermeasure to reduce post spaceflight postural and locomotor dysfunction. This training would serve to enhance the potential repertoire of adaptive behaviors available for individual astronauts by enhancing adaptive generalization capability (learning to learn). In addition, the training would

enhance the subject's ability to become more adept at sensory reweighting by systemically exposing them to different combinations of surrogate sensory disturbances during the training. Therefore the goal is not to expose subjects to actual gravitational transitions –because prolonged 0-g to 1-g or partial g transitions is not possible - but to enhance adaptive generalization capabilities along with increased ability to reweigh veridical sensory information during periods of sensory conflict.

Preflight adaptability training could play a central role in facilitating crewmember adaptive response to new gravitational environments in support of both short and long-duration space flight. Given that training may be retained for many months preflight training countermeasures would probably be sufficient to increase adaptability, even for long-duration flight. Therefore one can conceive of this training more in terms of a preflight “inoculation” that will not require significant amounts of crew time preflight and may only require infrequent “booster” training to maintain the training effect. Finally, a collateral benefit of the application VR technology, in this context, will be to make training programs more interesting, ultimately leading to increased adherence to prescribed training regimens.

Astronauts show significant inter-subject variations in their abilities to adapt to microgravity and to readapt to Earth's gravitational environment. Developing predictive measures of sensorimotor adaptability would allow optimization of training prescriptions by designing and implementing sensorimotor adaptability training countermeasures that would be customized for each astronaut's unique sensory bias and individual adaptive capabilities (Seidler *et al.*, 2015). Customization would allow more efficient use of crew time during training and produce better outcomes. This is a topic of current research in NASA's Human Research Program.

Healthy adults integrate visual, vestibular, and somatosensory information to produce appropriate motor output. The degree to which these sensory inputs are weighted and reorganized in discordant conditions varies by individual. Sensory weighting preferences have been reported for special populations and observed in healthy individuals under certain conditions. For example, higher visual dependence has been documented for stroke patients (Bonan *et al.*, 2004; de Haart *et al.*, 2004), astronauts (Young and Shelhamer, 1990) elderly people (van Hedel and Dietz, 2004), and vestibularly-intact but anxious normals (Viaud-Delmon *et al.*, 2000). Vestibular weighting has been shown to increase just before initiating a turn while walking (Kennedy *et al.*, 2005) and may be more prevalent in individuals who suffer chronic headaches (So and Bent, 2009). Autistic children (Masterton and Biederman, 1983) and individuals susceptible to mal de débarquement symptoms (Nachum *et al.*, 2004) depend more heavily on somatosensory cues. Approximately 30% of healthy normals are “highly” visually dependent (Brady *et al.*, 2009; Keshner *et al.*, 2004; Streepey *et al.*, 2007; Warren *et al.*, 1996). Healthy adults walking in novel discordant conditions display inherent differences in how they weigh visual information (Brady *et al.*, 2009). Adaptation in walking performance during exposure to a new discordant sensory environment can be predicted based on individual inherent level of visual dependency (Brady *et al.*, 2012). These data indicate that subjects who are more reliant on vision for control of movement have more difficulty adapting their walking

strategies in new environments. A high level of visual dependency might therefore predict a decreased ability to adapt to novel environments. Other individual sensory biases including vestibular and proprioceptive biases may also predict sensorimotor adaptability and could conceivably be used as predictive metrics. Therefore an individualized adaptability training program designed to decrease dependency on a single sensory source and to promote use of multiple sensory modalities and facilitate the process of sensory reweighting would enhance the individual's ability to adapt to novel discordant sensory environments like spaceflight (Carriot *et al.*, 2015).

#### **2.4. Biofeedback Training**

The autonomic nervous system initially responds to motion-induced stress through sympathetic activation followed by parasympathetic activation. If nausea and vomiting are parasympathetic reactions to sympathetic activation, then motion sickness symptoms might be prevented by training an individual to maintain autonomic regulation at baseline (Cowings *et al.* 1986).

Autogenic feedback training (AFT) combines cognitive imagery and body exercises to produce a desired change in autonomic activity. Sensory feedback regarding information about selected autonomic activities is provided to the subject through visual or auditory cues (Cowings 1990). Two crewmembers in the U.S. Shuttle program performed AFT before flight and two other crewmembers on the same flight served as untrained controls. During flight, the crewmember who showed mixed success in achieving autonomic control during training experienced one severe episode of SMS. Limited space flight testing showed that a crewmember exhibiting greater autonomic control during training before flight reported no severe symptoms. Two untrained crewmembers had multiple episodes of severe symptoms despite the administration of anti-motion sickness drugs (Cowings 1990).

Because AFT does not change the perception of vestibular stimulation, this type of training must interrupt the autonomic response after the sensory conflict has already occurred. Therefore, AFT and parasympatholytic drugs like scopolamine may achieve the same effect but through different mechanisms. AFT reduces sympathetic activity, thereby eliminating the parasympathetic reaction and its resultant symptoms. Scopolamine reduces the parasympathetic response to the increased sympathetic activity that has already occurred (Cowings *et al.* 1986).

#### **2.5. In-Flight Exercise**

Physical exercise has been a critical component of countermeasures for long-duration flights. The ISS exercise program consists of both resistive and aerobic exercise. Aerobic exercise has been performed on both the Treadmill Vibration Isolation System (TVIS) and the Cycle Ergometer with Vibration Isolation System (CEVIS) and now more recently using the Combined Operational Load Bearing External Resistance Treadmill (COLBERT or T2). For resistive exercise the interim Resistive Exercise Device (iRED) was initially implemented on the ISS but this system had some limitations including limited loading and resistance that was not constant. The newer resistance exercise device named the Advanced Resistive Exercise Device

(ARED) utilizes vacuum cylinders and inertial flywheels to simulate constant mass and inertia of free weight exercise and provides twice the loading of the iRED. In addition to attenuating loss of bone mineral density and muscle mass (Smith *et al.* 2012) crewmembers that exercised on the ARED also has less decrement in postflight postural stability and agility scores compared to subjects using the iRED (Wood *et al.* 2011). The increased body loading during ARED exercises may have provided greater postural challenges during exercise improving postflight balance performance.

## **2.6. Head Movements during and after Reentry**

The process of readapting to Earth's gravity may be facilitated at the time of reentry when crewmembers perform systematic head movements. Because of operational constraints, scientific study at this time has not been possible, but anecdotal reports from Shuttle crewmembers have indicated that performing head movements that slowly increase in amplitude can minimize motion illusions and motion sickness. The head movements were performed in the yaw plane initially and then in the pitch and/or roll planes, using progressively larger head tilts. Each crewmember should only perform movements at amplitudes and rates that he or she can tolerate at the time, but crewmembers should not restrict head movements too much or they may have problems when they must move to exit the spacecraft. The configuration of crewmembers in the Soyuz at landing, the volume of the spacecraft, and the higher G profile makes moving the head systematically more difficult than it was in the Shuttle, but making systematic head movements during and after reentry is still recommended to crewmembers.

## **2.7. Postflight Balance Rehabilitation Therapy**

Balance rehabilitation therapy is the general name for an exercise-based program for reducing the symptoms of disequilibrium and dizziness associated with vestibular disease or disorder. A common neuro-otological approach for managing such symptoms is to prescribe medication that suppresses vestibular function. However, in the long term, such suppressants can interfere with a person's ability to make necessary adaptations. In addition, many of these medications cause drowsiness that may limit a person's ability to be active. Medications, with their side effects, can be avoided by using balance rehabilitation therapy instead.

Vestibular rehabilitation therapy (VRT) is a type of balance rehabilitation therapy. It includes specific exercises that can eliminate or significantly reduce symptoms by promoting CNS compensation for inner-ear deficits. The program is designed to achieve these goals: (a) decrease dizziness and visual symptoms; (b) increase balance and walking functions; and (c) increase general activity levels. The program may include exercises for: (a) coordinating eye and head movements; (b) stimulating the symptoms of dizziness in order to desensitize the vestibular system; (c) improving balance and walking ability; and (d) improving fitness and endurance. Exercises vary depending on the type of symptoms. Balance retraining exercises are designed to achieve steadier walking and standing through improvements in coordination of muscle responses and organization of sensory information (e.g., vision, proprioception). Such treatment is part of the "post-flight reconditioning" of astronauts for 2 hours each day after

long-duration space flight. Reconditioning specialists supervise individualized post-flight reconditioning activities, adjusting the level of task difficulty according to the crewmember's level of balance recovery. Most of the activities benefit many body systems even if they target specific functions ([Wood et al. 2011](#)).

The exercises selected for post-flight reconditioning are based on activities of daily living that crewmembers have reported to be challenging, such as bending over to pick up an object, stooping down to tie one's shoes, and tilting the head backward. Signs of sensorimotor deconditioning are making wide turns, having difficulty changing direction, and making deliberate and slow motions that involve coordination of body segments. Slowed reaction times, difficulty in judging distances, and misperception of force also impair crewmembers' abilities to perform their normal activities.

Post-flight reconditioning activities progress from lower risk to higher risk and simple to complex as crewmembers master particular skills and as particular movements become less provocative. Some exercises are performed every day and others are performed every other day. Posture and stability exercises are performed every day while standing. They include head movements in different planes, toe touches followed by hyperextensions, and trunk twists. Standing with both legs on a stable surface progresses to standing on one leg and standing on an unstable surface. Because mobility facilitates the resolution of sensorimotor symptoms after space flight, astronauts are encouraged to walk as much as possible, beginning on landing day. The limits of the individual's motion tolerance are continually challenged; a "walking toe touch" is one of the most provocative activities. Some activities involve catching and throwing balls of increasing weight, sometimes while walking or shuffling.

Reconditioning specialists and flight surgeons use quantitative measures to evaluate post-flight recovery of each crewmember. Posture is assessed by computerized dynamic posturography and sensory organization tests, including some performed with dynamic pitch head tilts. Agility is measured by testing crewmembers' ability to move forward, backward, and to either side around cones on the floor. Preflight and post-flight tests of posture and agility are used to measure the effectiveness of countermeasures such as in-flight exercise.

### **3. MECHANICAL DEVICES**

Various mechanical and electrical devices have been explored in the former Soviet Union to alleviate the symptoms of space motion sickness. The mechanical devices are designed to prevent the complete adaptation of the body to weightlessness by counteracting deconditioning during long missions as well as relieving motion sickness symptoms during the first days of flight. The mechanisms in which electrical devices operate are still unclear.

#### **3.1. Dynamic Foot Stimulation**

It has been hypothesized that the body unloading and subsequent loss of support afferentation experienced during space flight induces a cascade of neuromotor alterations leading to neuromuscular dysfunction including loss of tonic muscle activation and subsequent post-flight postural and locomotor instability ([Kozlovskaya et al. 2006](#); [Kozlovskaya et al. 2007](#)).

Research has been conducted to examine the use of dynamic foot stimulation as a potential supplement to more traditional in-flight exercise countermeasures to mitigate neuromotor degradation associated with long-duration space flight. Several studies have shown that dynamic foot stimulation restores absent neuromuscular activation during space flight throughout the entire lower-limb musculature (Layne *et al.* 1998a). Mechanical stimulation of the feet was first applied on a Russian crewmember on the Salyut-6 space station using footwear with pneumatic bladders. It was reported that this stimulation preserved lower limb strength and locomotor function (Hernandez-Korwo *et al.* 1983).

Layne *et al.* (1998a) examined whether applying foot pressure using inflatable bladders inside boots during space flight could enhance the neuromuscular activation associated with posture perturbing rapid arm movements. The boots were worn during an 81-day mission for two crewmembers and 115 days for the two other crewmembers on the Mir space station. The crewmembers were required to perform rapid, unilateral shoulder flexions while free floating. The results showed that the application of foot pressure elicited lower limb and trunk activation typically observed in 1 g but that was not present without this stimulus in microgravity. Importantly, the lower-limb musculature response produced by foot stimulation was maintained through flight day 188 during a Mir mission indicating no habituation to the stimulus (Layne & Forth 2008). Another study examined mechano-stimulation of the soles support zones during exposure to 7 days of simulated microgravity conditions using the dry immersion technique that provides more gravitational unloading along the long axis of the body compared to bed rest. Foot pressure was applied to the heel and forefoot every day for 20 minutes. Subjects completed slow 10 minute pacing followed by 10 minutes of fast pace walking simulations. The results showed that electromyographic activities of leg extensors improved with the use of foot stimulation. In addition there was an increase in flexor muscle activity (Layne & Forth 2008).

Mechano-stimulation of the soles of the feet has been shown to improve postural equilibrium control and functional mobility. In a previous study subjects were asked to negotiate a complex obstacle course before and after 60 days of bed rest (Reschke *et al.* 2009). In this study bed rest subjects performed the Functional Mobility Test (FMT) which entailed walking at a self-selected pace through an obstacle course set up on a base of medium density foam. Also, unique to this study was that daily foot massages were given to all bed rest subjects, after the first three subjects were completed to ameliorate the tenderness of the soles of feet. So the first three subjects tested did not receive any foot stimulation for the duration of the study. This enabled a comparison in performance between these two groups. Subjects in both groups showed a significant increase in their time to complete the course after bed rest. However, subjects who did not receive foot massage performed much worse after bed rest (94% increases in time) compared to those who did receive foot massage (27% increase in time). These data point to the importance of foot tactile and pressure input as a central component in the control of posture and locomotion.

A more recent study investigated the impact of space flight induced changes in the level of foot skin sensitivity on post-flight postural disequilibrium. Skin sensitivity of Shuttle astronauts was measured as vibration perception at the great toe, fifth metatarsal and heel. Both

increased and decreased skin sensitivity was observed post-flight. Increased skin sensitivity was associated with a greater reduction in postural equilibrium control using computerized dynamic posturography (CDP) pointing to the importance of foot skin receptors in maintaining balance control and its role in post-flight postural disequilibrium (Lowrey *et al.* 2014).

In support of the observations above, stimulation of the plantar surface of hindlimb-suspended rats resulted in preservation of soleus muscle mass cross sectional area (Kyparos *et al.* 2005). These studies used a pneumatic cuff that rapidly inflated and deflated to provide dynamic stimulation to the soles. The rats in the experimental group received this foot stimulation for only 5.6% of the entire period they were suspended, therefore, this relatively short time period of exposure was sufficient to maintain muscle size despite the limb unloading. Therefore, restoring support-loading afferentation by providing foot tactile input can be used to prevent muscle atrophy associated with space flight.

### **3.2. Load Suits**

Load suits are composed of adjustable elastic bands that produce tension over the chest, back, abdomen, side, and leg seams. The suits were first worn by the crews of the Soyuz-13 and Soyuz-14/Salyut-3. Although the load suits were considered “pleasant,” illusions, headward fluid shifts, and symptoms of SMS were still present (Gurovskii *et al.* 1975; Vorob'ev *et al.* 1975).

New systems are currently being developed. For example, the Variable Vector Countermeasure Suit (V2Suit) uses inertial measurement units and control moment gyroscopes within miniaturized modules placed on body segments to provide a viscous resistance during movements against a specified direction of “down” (Duda *et al.* 2015). This suit could act as a countermeasure to the sensorimotor adaptation performance decrements during long-duration spaceflight, including post-flight recovery and rehabilitation. This type of countermeasure system also has Earth benefits, particularly in gait or movement stabilization and rehabilitation.

### **3.3. Stroboscopic Goggles to Mitigate Motion Sickness**

Motion sickness in the general population is a significant problem driven by increasingly sophisticated modes of transportation, visual displays, and virtual-reality environments. It is important to investigate nonpharmacological alternatives for the prevention of motion sickness for individuals who cannot tolerate the available anti-motion sickness drugs, or who are precluded from medication because of operational environments. Both NASA and the U.S. Army have been investigating stroboscopic vision as a way to provide a simple and easily managed treatment for motion sickness (Reschke *et al.* 2006). Specifically, a five-part study was designed to investigate the effect of stroboscopic vision (with either a strobe light or liquid crystal display [LCD] shutter glasses) on motion sickness while (1) using visual field reversal; (2) reading while riding in a car (with or without external vision present); (3) making large pitch head movements during the 0 g phase of parabolic flight; (4) exposed to rough seas in a small boat; and (5) seated and reading in the cabin area of a UH60 Black Hawk helicopter during provocative flight patterns. A total of 69 subjects participated in selected phases of the study. Fewer subjects suffered from motion sickness under stroboscopic conditions. Stroboscopic

illumination prevents retinal slip, thereby treating motion sickness symptoms. Shutter glasses with a cycle frequency of 4 or 8 Hz and a short dwell (glasses clear) time (10–20 ms) are as effective as a strobe light, producing a useful adaptation during either self- or surround-motion without the penalty of using disabling MS drugs (Reschke *et al.* 2007b).

### **3.4. Tactile Spatial Awareness System**

A recent joint ESA-NASA pre- and post-flight experiment was designed to examine both the physiological basis and operational implications for disorientation and tilt-translation disturbances in astronauts after short-duration space flights. Changes in eye movements and motion perception were investigated during independent tilt and translation motion profiles in eight crewmembers returning from Space Shuttle missions. Roll motion was provided by a variable-radius centrifuge, while pitch motion was provided by NASA's Tilt-Translation Sled, in which the resultant gravitoinertial vector remains aligned with the body's longitudinal axis during tilt motion. Subjects were told to use a joystick to nullify tilt motion disturbances on these two devices. The stimuli consisted of random steps or sum-of-sinusoids.

Results showed that the ability to control tilt orientation was compromised following space flight, with an increased number of errors corresponding to changes in self-motion perception. However, the use of a tactile spatial awareness system (TSAS) improved manual control performance. During the closed-loop nulling task on both devices, small tactors placed around the torso vibrated according to the actual body tilt angle relative to gravity. Performance on the closed-loop tilt control task improved with the tactile display feedback of tilt orientation during both pre- and post-flight testing. In fact, with the TSAS, the performance during early post-flight tests was comparable to that without TSAS during the preflight tests (Clément *et al.* 2014).

### **3.5. Electrical Devices**

#### **3.5.1. Galvanic Vestibular Stimulation**

Galvanic Vestibular Stimulation (GVS) entails electrical stimulation to the vestibular labyrinth via surface electrodes placed over the mastoid bones that pass small currents that activate primary vestibular afferents. It has been used to stimulate the vestibular labyrinth artificially for laboratory studies of human vestibular cortex, spatial orientation, postural control and locomotion (Moore *et al.* 2011, 2015). GVS produces behavioral changes in balance function, gaze, head-trunk coordination and locomotor disturbances that are similar to those observed in post-flight astronauts (Moore *et al.* 2006). GVS has been validated in the Vertical Motion Simulator at NASA Ames during high-fidelity Shuttle landing simulations. When exposed to GVS, pilot subjects (including a veteran shuttle commander of 3 flights), experienced spatial disorientation and subsequent decrements in landing performance equivalent to that observed in actual Shuttle landings (Moore *et al.* 2011). The GVS analog accurately reproduces the effects of microgravity exposure on the central nervous system, and might be used to improve training of astronauts for future long-duration missions (Dilda *et al.* 2014). Dilda *et al.* have shown that subjects exhibit central adaptation phenomenon with repeated GVS exposure in a study of normal subjects for an extended period of up to 12 weeks (120 min of total exposure). During each trial subjects performed computerized dynamic posturography and eye movements were

measured using digital video-oculography. Follow up tests were conducted 6 weeks and 6 months after the 12-week adaptation period. Postural performance was significantly impaired during GVS at first exposure, but recovered to baseline over a period of 7–8 weeks (70–80 min GVS exposure). This postural recovery was maintained 6 months after adaptation. In contrast, the roll vestibulo-ocular reflex response to GVS was not attenuated by repeated exposure. This suggests that GVS adaptation did not occur at the vestibular end-organs or involve changes in low-level (brainstem-mediated) vestibulo-ocular or vestibulo-spinal reflexes. Faced with unreliable vestibular input, the cerebellum reweighted sensory input to emphasize veridical extra-vestibular information, such as somatosensation, vision and visceral stretch receptors, to regain postural function. After a period of recovery subjects exhibited dual adaptation and the ability to rapidly switch between the perturbed (GVS) and natural vestibular state for up to 6 months. In a follow up study (Moore *et al.* 2015) found that pre-adaptation to GVS is associated with enhanced sensorimotor performance in a flight simulator and asked to null the roll motion of a visual bar presented on a screen using a joystick compared to subjects who were not adapted to GVS. Thus GVS may be used as a training modality to enhance adaptability to aid recovery after space flight (Dilda *et al.* 2014; Moore *et al.* 2015).

### **3.5.2. Vestibular Stochastic Resonance**

A general phenomenon of neural systems is that paradoxically if noise is added to neural sensory systems, their ability to detect sub-threshold signals improves, which is a mechanism termed Stochastic Resonance (SR). SR thus enables the enhanced detection of relevant sensory signals. SR can be thought of simply as “noise benefit” by increasing information transfer in the presence of non-zero level of noise (for reviews see (Collins *et al.* 2003; Moss *et al.* 2004; McDonnell and Abbott 2009; Aihara *et al.* 2010). SR has been observed in human hearing (Jaramillo and Wiesenfeld 1998; Zeng *et al.* 2000; Ward *et al.* 2002) and has been identified as an important component in cochlear coding strategy (Morse and Evans, 1996). The presence of stochastic noise to sensory input has been shown to improve: visual contrast sensitivity and detection (Simonotto *et al.* 1997; Ward *et al.* 2002); the degree of association between the heart rate responses and weak periodic oscillatory variation in central venous pressure (Soma *et al.* 2003); letter recognition (Piana *et al.* 2000); perception of ambiguous figures (Riani and Simonotto 1994); and visual depth perception (Ditzinger *et al.* 2000). SR in tactile sensation has been demonstrated in the response to weak mechanical stimuli (Collins *et al.* 1996a; b; Collins *et al.* 1997; Ivey *et al.* 1998; Richardson *et al.* 1998). The application of mechanical noise to the feet has been shown to improve balance control through the reduction of sway in young and elderly subjects (Priplata *et al.* 2002; Priplata *et al.* 2003), in patients with diabetes and stroke (Priplata *et al.* 2006) and patients with functional ankle joint instabilities (Ross and Guskiewicz 2006; Ross *et al.* 2013). Similarly, balance improvement has been demonstrated with electrical noise applied to the back of the knee (Gravelle *et al.* 2002). Vibratory noise applied to the fingertip also enhanced balance performance based on SR phenomenon (Magalhaes and Kohn 2011). These same authors have also shown that the application of imperceptible electrical noise to the triceps surae during a seated task reduced force fluctuations in a force matching task of isometric plantar flexion force which were correlated to subsequent reductions in postural sway during quiet stance based on SR phenomenon (Magalhaes and Kohn 2012). There

have been a few studies that showed the effectiveness of applying sub-sensory vibratory noise to the soles of the feet during over-ground walking comparing elderly population with young control subjects (Galica *et al.* 2009). In a follow-up study, this group also showed the effectiveness of applying sub-sensory vibratory noise to the soles of the feet during treadmill walking in a set of control subjects (Stephen *et al.* 2012). SR using white noise based electrical stimulation at imperceptible amplitudes (at or below peri-threshold levels) of the vestibular system, applied using surface electrodes on the mastoid, leads to significantly improved balance, and locomotor performance during periods of novel sensory challenges in normals and Parkinsonian Disease patients (Mulavara *et al.* 2011, 2015a, 2015b, Goel *et al.* 2015, Samoudhi *et al.* 2015).

The methodology of using sub-threshold electrical stimulation of the vestibular system or similar stimulation of the proprioceptive systems at the sole of the feet will help retain and enhance the use of vestibular or proprioceptive information in performance of specific functions. Studies in the literature have investigated the usefulness of SR in conjunction with traditional training paradigms to improve performance (Ross and Guskiewicz 2006; Ross 2007; Ross *et al.* 2007; Ross *et al.* 2013). These investigators have shown significant improvement in postural balance control aiding recovery when electrical or mechanical SR stimulation to the muscles across the ankle joints was given in conjunction with conventional coordination training compared to training alone (Ross and Guskiewicz 2006; Ross 2007; Ross *et al.* 2007; Ross *et al.* 2013). Therefore, in general an individualized sensorimotor training program in conjunction with SR designed to promote the use of multiple sensory modalities can enhance the ability to adapt postural control and walking stability when exposed to a novel discordant sensory environment in the astronaut population.

#### **4. ARTIFICIAL GRAVITY**

Sensorimotor performance, but also bone loss, muscle weakening, and cardiovascular deconditioning, among other deficits, are all known deconditioning due to microgravity. The longer the flight duration, the more serious the health consequences become (Clément 2011). The current countermeasures on board the ISS (exercise, pharmaceuticals, food complements) address each of these physiological systems in a piece-meal fashion. Artificial gravity, i.e., a sustained centripetal acceleration generated by centrifugation, represents a novel and integrated approach to addressing the detrimental effects of reduced gravity on the human body (Clément & Bukley 2007). All body systems are challenged simultaneously by its application, not simply one physiological system at a time. In addition, artificial gravity is an improvement upon the current ISS countermeasures as it addresses the root causes of the deconditioning phenomenon instead of treating its end-effects system-by-system as the current countermeasures do.

More recently, a special focus of concern is the deficit in vision acuity in astronauts on board the ISS, which is hypothesized to be caused by weightlessness-induced fluid shifts to the upper body leading to intracranial hypertension (Mader *et al.* 2011). If this hypothesis is confirmed, it could be an impediment for future long-duration deep space missions. Thus an effective countermeasure against these effects will be required. Because it enables re-

establishing g-induced hydrostatic gradients, centrifugation might be the most efficient countermeasure.

Generating centrifugal force equivalent to the gravitational force on Earth during long-duration exploration missions can be obtained by rotating the entire spacecraft. However, this solution is costly in terms of power and mass, and it creates issues with navigation and control, communication, and docking. Another, more affordable solution, is to rotate only one part of the spacecraft, or to utilize an on-board human centrifuge.

To help inform the final decision on whether to conduct continuous spin of the whole space vehicle or to intermittently expose the crewmember to short-radius centrifugation, the limits of human adaptation in a rotating environment must be revisited. We need to identify the acceptable and/or optimal ranges for radius and rotation rate to avoid unacceptable crew health and performance consequences (Lackner & DiZio 2005a; Arya *et al.* 2007; Fong *et al.* 2007; Reinertson *et al.* 2007; Warren *et al.* 2007; Symons *et al.* 2009; Zwart *et al.* 2009). For intermittent applications, we need to identify what level, duration, frequency, and time of day of exposure to artificial gravity are optimal (Young *et al.* 2007). We also need to investigate the physiological responses to transitions between artificial gravity, microgravity, and Moon or Mars gravity because such studies would be useful in assessing whether dual adaptation to a rotating and a non-rotating environment is possible (Lackner & DiZio 2003).

NASA and other space agencies are working on a global research program on artificial gravity that would leverage the facilities available around the world (e.g., short- and long-radius centrifuges, slow rotating rooms, bed rest/dry immersion facilities, suspension systems, etc.) and integrate studies on human, animal, and cell models. Standardization of measures performed before and after each artificial gravity intervention will allow for more compatible assessment across various studies. The biomedical measurements will focus on countermeasure validation, medical events, and subject acceptance and comfort.

Regarding sensorimotor performance, artificial gravity projects that could be performed in the near future include the following: (a) test gravity level values along Gz within the range from microgravity to 1 g, using the methods described above, to reasonably reach conclusions on the threshold, optimal stimulus-response, and saturation for the effects of centrifugation on sensorimotor performance; (b) test the effects of gravity levels higher than 1 G to assess whether increasing the intensity of the Gz stimulus actually reduces the time of exposure needed; (c) compare whether exposure to centrifugation for intermittent, short periods of time in one or multiple sessions is as beneficial as continuous exposure to Earth's gravity; (d) investigate whether Gz centrifugation reduces intracranial pressure and possibly mitigates the visual impairment due to intracranial pressure (VIIP) syndrome; (e) assess whether centrifugation can possibly mitigate post-flight decrease in performance by studying the effect of centrifugation on cognitive and functional tasks; and (f) assess the effects of gravity gradient on spatial orientation by comparing the responses in subjects placed at various distances from the axis of rotation on a long radius centrifuge (Clément *et al.* 2015b).

Please see Table 2 below which provides an executive summary of the sensorimotor evidence, providing links from the physiological deficits to subsequent performance implications as well as to countermeasures currently in place, potential countermeasures previously developed but requiring verification, and potential future countermeasures under development.

<b>Evidence (Physiological Deficits)</b>	<b>Performance Implications (Functional Deficits)</b>	<b>Current Countermeasures</b>	<b>Potential Countermeasures</b>
<p><b>Sensation</b></p> <p>(Inappropriate sensory cues lead to blurry vision, perceived self/surround motion, inability to determine body position/force, altered mass discrimination, etc.)</p>	<p>Mobility, Driving, Flying, Navigation, Landing a vehicle, Rendezvous/Docking, Robotics, Fall/Fall Recovery, Seat egress, Vehicle egress, Obstacle avoidance, Object translation</p>	N/A	<p>Vestibular Training, Preflight Adaptation Training, Sensorimotor Adaptability Training, In-flight Exercise on Treadmill w/Head Movements, In-flight Exercise with Balance Training, Head Movements During/After Reentry, Balance Rehabilitation, Dynamic Foot Stimulation, Load Suits, Tactile Spatial Awareness System, Galvanic Vestibular Stimulation, Vestibular Stochastic Resonance, Artificial Gravity</p>
<p><b>Gaze Control</b></p> <p>(Blurry vision, slow/inaccurate target acquisition/tracking)</p>	<p>Mobility, Driving, Flying, Navigation, Landing a vehicle, Rendezvous/Docking, Robotics, Fall/Fall Recovery, Seat egress, Vehicle egress, Obstacle avoidance, Object translation</p>	N/A	<p>Vestibular Training, Preflight Adaptation Training, Sensorimotor Adaptability Training, In-flight Exercise on Treadmill w/Head Movements, Head Movements During/After Reentry, Balance Rehabilitation, Galvanic Vestibular Stimulation, Vestibular Stochastic Resonance, Artificial Gravity</p>
<p><b>Eye-hand Coordination</b></p> <p>(Slow and/or inaccurate: limb motion, target acquisition/tracking, dual-tasking)</p>	<p>Driving, Flying, Navigation, Landing a vehicle, Rendezvous/Docking, Robotics, Vehicle egress, Object translation</p>	N/A	<p>Preflight Adaptation Training, Sensorimotor Adaptability Training, Tactile Spatial Awareness System, Artificial Gravity</p>
<p><b>Spatial Orientation Perception</b></p> <p>(Self and geographical disorientation, self-/surround-motion)</p>	<p>Mobility, Driving, Flying, Navigation, Landing a vehicle, Rendezvous/Docking, Robotics, Fall/Fall Recovery, Seat egress, Vehicle egress, Obstacle avoidance, Object translation</p>	N/A	<p>Vestibular Training, Preflight Adaptation Training, Sensorimotor Adaptability Training, Tactile Spatial Awareness System, Galvanic Vestibular Stimulation, Vestibular Stochastic Resonance, Artificial Gravity</p>
<p><b>Cognitive Function</b></p> <p>(Poor concentration, memory loss, decreased multi-tasking, inaccurate distance/size/time perception)</p>	<p>Driving, Flying, Navigation, Landing a vehicle, Rendezvous/Docking, Robotics</p>	N/A	<p>Preflight Adaptation Training, Sensorimotor Adaptability Training</p>

<b>Evidence (Physiological Deficits)</b>	<b>Performance Implications (Functional Deficits)</b>	<b>Current Countermeasures</b>	<b>Potential Countermeasures</b>
<b>Space Motion Sickness</b> (Nausea, vomiting, drowsiness, dizziness, headache, etc.)	Mobility, Driving, Flying, Navigation, Landing a vehicle, Rendezvous/Docking, Robotics, Fall/Fall Recovery, Seat egress, Vehicle egress, Obstacle avoidance, Object translation	Pharmacological	Vestibular Training, Preflight Adaptation Training, Sensorimotor Adaptability Training, Biofeedback, Head Movements During/After Reentry, Stroboscopic Goggles, Galvanic Vestibular Stimulation, Vestibular Stochastic Resonance, Artificial Gravity
<b>Orthostatic Intolerance</b> (Dizziness, nausea, fainting)	Mobility, Fall/Fall Recovery, Seat egress, Vehicle egress, Obstacle avoidance, Object translation	Fluid Loading, Compression Garments	Vestibular Training, Artificial Gravity
<b>Peripheral/Central Neural Changes</b> (Changes to: vestibular organ, brain, nerve conduction, muscle coordination)	Mobility, Driving, Flying, Navigation, Landing a vehicle, Rendezvous/Docking, Robotics, Fall/Fall Recovery, Seat egress, Vehicle egress, Obstacle avoidance, Object translation	N/A	Artificial Gravity
<b>Postural Equilibrium</b> (Inability to maintain balance)	Mobility, Fall/Fall Recovery, Seat egress, Vehicle egress, Obstacle avoidance, Object translation	N/A	Vestibular Training, Sensorimotor Adaptability Training, In-flight Exercise on Treadmill w/Head Movements, In-flight Exercise with Balance Training, Head Movements During/After Reentry, Balance Rehabilitation, Dynamic Foot Stimulation, Load Suits, Tactile Spatial Awareness System, Galvanic Vestibular Stimulation, Vestibular Stochastic Resonance, Artificial Gravity
<b>Locomotion/ Jump Performance</b> (Inability to walk/run, inability to jump off a relatively low surface and land safely on another; i.e. bottom rung of ladder, traversing uneven surfaces)	Mobility, Fall/Fall Recovery, Seat egress, Vehicle egress, Obstacle avoidance, Object translation	N/A	Vestibular Training, Sensorimotor Adaptability Training, In-flight Exercise on Treadmill w/Head Movements, In-flight Exercise with Balance Training, Head Movements During/After Reentry, Balance Rehabilitation, Dynamic Foot Stimulation, Load Suits, Tactile Spatial Awareness System, Galvanic Vestibular Stimulation, Vestibular Stochastic Resonance, Artificial Gravity

*Table 2. Summary of the evidence providing links amongst the physiological deficits, performance implications, current countermeasures, and potential countermeasures.*

## VI. COMPUTER-BASED MODELING AND SIMULATION

While there are few robust ground-based models available for experimental investigations of the impacts of space flight on a crewmember's mobility and ability to maintain control of vehicles and complex systems, many computer-based models of the vestibular system and sensorimotor control have been developed. Since these may be useful in simulating and/or predicting the impacts of physiological adaptations on operational performance, particularly under off-nominal conditions, a brief review of the relevant aspects of the field is provided in this section. Before they can be used in design and verification, though, these (and other) models must be quantitatively validated and certified using targeted empirical studies

### 1. MODELS OF VESTIBULAR FUNCTION AND SPATIAL ORIENTATION

Vestibular neuroscientists have developed quantitative mathematical models for semicircular canal and otolith function, eye movements, and central nervous system (CNS) estimation of angular and linear motion perception. For example, Fernandez & Goldberg (51) modeled the firing frequency  $f_i$  of individual semicircular canal afferents using a linear transfer function model of the form  $f_i(s)/\omega(s) = s^2 K_i [(K_r s + 1)/(\tau_d s + 1)(\tau_c s + 1)(\tau_a + 1)]$ , where  $\omega$  is the component of head angular velocity in the canal plane,  $K_i$  is mid frequency gain,  $K_r$  is high frequency gain,  $\tau_d$  and  $\tau_c$  are the time constants of endolymph flow drag development and cupula-endolymph return (Groen 1957; Oman *et al.* 1987), and  $\tau_a$  is a time constant describing neural adaptation (Young *et al.* 1969).

Young *et al.* (1967) originally suggested that the CNS functions like an adaptive (Kalman) filter when combining sensory cues, and introduced additional dynamics into vestibular responses due to these central processes. Adapting inertial guidance theory, Young (1969; Young *et al.* 1973; Young *et al.* 1984) noted that laws of physics dictate that the body's graviceptors respond to the net gravito-inertial specific force ( $\mathbf{f} = \mathbf{g} - \mathbf{a}$ ), the physical quantity tracked by a pendulum or measured by a linear accelerometer (where  $\mathbf{a}$  = linear acceleration vector and  $\mathbf{g}$  = gravitational acceleration vector). A variety of different orientations and accelerations can cause in the same graviceptor stimulus. The CNS must therefore use other cues to distinguish the components caused by gravity from those caused by linear acceleration. The CNS may estimate linear acceleration by maintaining an internal estimate of the direction and magnitude of ( $\hat{\mathbf{g}}$ ) and subtracting off the graviceptor cue vector ( $\hat{\mathbf{a}} = \hat{\mathbf{g}} - \mathbf{f}$ ). The direction of down,  $\hat{\mathbf{g}}$  is estimated at low frequencies based on the average direction of graviceptor cues,  $\mathbf{f}$ , and also visual cues, if available. Visual inputs are angular and linear velocity of the visual surround with respect to the observer. At high frequencies, semicircular canal cues and body movement commands are used. If the direction of  $\hat{\mathbf{g}}$  is misestimated, dramatic misperceptions of orientation and linear acceleration can result.

Although "optimality" of the human observer (in the Kalman sense) has since been discounted, the notion remains widely accepted that the CNS functions as an "observer," in the control engineering sense (Borak *et al.* 1979), estimating head orientation based on internal representations of the direction of gravity and sensory organ dynamics. Others have elaborated CNS observer-based models for semicircular canal-otolith interaction. For example, Raphan *et*

*al.* (1979), Robinson (1981), and Merfeld *et al.* (1993) developed influential observer class models for CNS estimation of head angular velocity and tilt, now often referred to as “central velocity storage” theories. Merfeld’s models for canal-otolith cue interaction in “down” estimation (Merfeld & Zupan 2002) successfully predict canal-otolith cue interaction in a variety of experimental situations. They are now widely utilized in research and the diagnosis of clinical vestibular disorders. These models have occasionally been applied to aircraft accident investigation, albeit in a limited way, since they do not (yet) incorporate effects of visual cues, and data on aircraft accidents is frequently lacking.

## 2. MODELS OF MANUAL CONTROL PERFORMANCE

Manual Control theory was originally developed in the 1960s, when feedback control engineers sought to analyze and predict the performance of humans in control loops, and describe both the human (the operator) and the controlled system (the plant) within the same mathematical framework. The premise was that human operator performance could be approximated well using a “describing function.” Both compensatory tasks (where the operator sees only an error signal) and pursuit tasks (where both the goal and plant outputs are available) have been modeled this way.

A simple and widely used principle is the “crossover model” (McRuer 1972), which posits that the operator will instinctively adopt an appropriate control strategy such that at the open loop transfer function of the operator and plant taken together resembles that of a simple integral process and a time delay in the region of the crossover frequency. The operator can perceive the rate of change of plant output, and create anticipatory phase lead that counteracts phase lags due to the plant. If the plant is a vehicle, vestibular motion cues allow the operator to improve performance by creating additional phase lead. However, the operator’s transfer function is constrained. Some effective time delay is always present due to perceptual, cognitive, and muscle activation effects. Also, operators cannot respond to the second or higher derivatives of plant output. The crossover model structure and parameter values thus quantify the operator’s control strategy. The model also has important emergent properties: It predicts manual control gain and bandwidth limits. It also explains why humans cannot successfully stabilize higher than second order integral plant dynamics, unless the operator is able to monitor intermediate system outputs, in effect transforming the task into concurrent (multi-loop) lower order tasks. This is why an operator cannot successfully stabilize a hovering lunar lander or a helicopter (approximately triple integral plants) over a landing spot without reference to a real or artificial horizon, and why motion cues can have a dramatic effect on controlling marginally stable plants (Young 1967; Shirley & Young 1968). The crossover model has been extended to multi-loop control and validated across a wide variety of plant dynamics, and extensively applied in many domains, particularly in the area of vehicle handling quality standards (Young *et al.* 1973).

In the late 1960s, newer estimation and optimal control concepts, such as the Kalman observer and controller, were used to extend manual control theory. The optimal control model (Baron 1969) posited that the human observer’s control strategy utilized an internal dynamic mental model for the plant, and it weighted feedback information based on prior knowledge of

uncertainties. (Concurrent efforts by neuroscientists led to the present generation of Observer Theory models for orientation and sensory conflict in motion sickness described earlier.) Early applications included helicopter hovering and attention sharing. Results demonstrated the importance of vestibular motion cueing (Curry *et al.* 1976; Baron 1983).

When performing maneuvers such as flaring an aircraft on landing, a highly skilled human operator uses a “precognitive control strategy,” and generates open loop, preprogrammed commands based on a mental model of the plant. The preprogrammed command accomplishes most of the maneuver, but the operator completes the task by switching back to conventional compensatory manual control for final error reduction. The Shuttle landing flare is an example of a task accomplished using precognitive control (Ashkenas *et al.* 1983). Landing performance depends critically on proper timing of the preprogrammed manual flare command, and correct estimation of the aircraft state at that moment. Incorrect precognitive manual commands result in greater need for subsequent compensatory error reduction. After the flare, the pilot exerts “tight” control over aircraft altitude and altitude rate in order to achieve a smooth touchdown, employing relatively high control gain. Because the Shuttle flight control system has inherent phase delays and rate limits, excessively large pilot control gain can make the combined pilot-vehicle system unstable, and trigger pilot induced oscillations (McRuer 1972). At the time it was not generally recognized that misperceptions of vehicle pitch attitude and rate could also potentially cause over control and pilot induced oscillations (PIO), but they were detected during the Shuttle Enterprise Approach and Landing Test flight test program, where disorientation was presumably not a factor. Since control system delays could not be eliminated, a stopgap solution was to detect large oscillatory control stick commands using a suitable nonlinear filter, and adaptively reduce pilot control authority (Smith & Edwards 1980). Adaptively reducing control authority worked for “conventional” PIO. However, as described earlier, STS-3 subsequently experienced a PIO despite the PIO suppression filter. The only solution for disorientation induced PIO is to provide strong visual cues to pitch and pitch rate via a HUD, and restricting landings to conditions of good visibility. If the Shuttle were required to land in brownout/grayout conditions (e.g. as are Lunar Landers), PIO would be a continuing concern.

Landing on a planetary body requires a number of operational tasks including identification of an appropriate location that is level and free of hazards while maintaining a stable controlled descent to the surface. Various sensorimotor challenges may interfere with crewmembers’ performance. These include the astronauts’ first exposure to partial gravity following microgravity adaptation, the unique vehicle motions experienced on decent, and dust blowback from the descent engine thruster that may obscure vision. Models of human spatial orientation perception have been developed that can be used to predict the potential for disorientation in partial gravity environments (Clark *et al.* 2015). These models predict that spatial misperceptions are likely to occur during landings in partial gravity environments, particularly with limited or incomplete visual cues. For example a powered descent acceleration profile creates the misperception that the landing vehicle is upright, even when the vehicle has a large pitch or roll angle. When full visual information is provided these perceptual illusions are largely suppressed, however, dust blowback during landing may obscure visual cues out the window

and exacerbate spatial disorientation. These model predictions have been validated empirically using the NASA Ames Vertical Motion Simulator in which subjects self-reported their perceptions of vehicle motions during lunar-landing-like motions (Clark *et al.* 2014). Current research is focused on development of advanced display systems that could be implemented as countermeasures for landing spatial disorientation that include enhanced situation awareness displays and synthetic terrain displays that may help reduce potential landing misperceptions.

## VII. RISK IN CONTEXT OF EXPLORATION MISSION OPERATIONAL SCENARIOS

### 1. SENSORIMOTOR STANDARD

A sensorimotor standard has been drafted (NASA Standard 3001) for exploration class missions: *“Pre-flight sensorimotor functioning shall be assessed and be within normal values for age and sex of the astronaut population. In-flight Fitness-for-Duty standards shall be guided by the nature of mission-associated high-risk activities. In-flight Fitness-for-Duty standards shall be assessed using metrics that are task specific. Sensorimotor performance limits for each metric shall be operationally defined. Countermeasures shall maintain function within performance limits. Post-flight reconditioning shall be monitored and aimed at returning to baseline sensorimotor function.”* However, operational performance limits related to mobility (e.g., emergency egress), flight vehicle control (particularly for post-adaptation activities, such as rendezvous/docking and entry/landing), ground vehicle control (e.g., Lunar or Martian rovers), and remote manipulator/teleoperation activities have not yet been established.

The very first astronaut candidates underwent rigorous selection tests, including neurological tests, but NASA conducted little screening for vestibular or sensorimotor problems in subsequent groups of new astronauts, including those in the Space Shuttle program. As evidence evolved from early space programs through the Shuttle missions, sensorimotor and CNS problems began to become prevalent medical findings. Current medical history and exams for astronaut selection include: (a) no history of serious ongoing neurological disease, (b) examination by a neurologist, and (c) magnetic resonance imaging of the brain and magnetic resonance angiography of the head.

Abnormalities detected by these methods help to screen out some asymptomatic individuals, some of whom have potentially serious problems. NASA had already required the use of electroencephalography as a part of the selection process before instituting these additional rigorous neurological examinations. Attempts were made historically to use susceptibility to motion sickness, abnormal visual-vestibular function, and postural problems to screen out candidates. Much of this testing, performed by in-house laboratories at the NASA Johnson Space Center, was seldom weighted highly, primarily because researchers in the Neurosciences Laboratories were hesitant to recommend serious limits based on standards that did not reflect operational requirements.

The high degree of variability across crewmembers in terms of the severity of neurological symptoms, given the current knowledge base, suggests that medical selection and retention standards could be quite effective in minimizing the operational impacts of sensorimotor adaptation. However, the lack of validated assessment tools for predicting sensorimotor adaptation, or an individual’s inability to adapt, has hindered the development of relevant selection standards. The capability for clinical diagnosis of vestibular disorders greatly advanced during the Space Shuttle era from direct spin-offs of Shuttle projects (e.g., postural testing, eye measurement technology, etc.) (Reschke *et al.* 2013). Although selection and retention standards for sensorimotor function have remained limited to neurological screenings of reflex

functions consistent with standard aviator flight physical examinations, extensive selection standards are now in place for vision, audition, and other sensory functions.

Flight rules have been used to minimize the operational consequences of vestibular and sensorimotor changes associated with microgravity. These rules primarily limited crew activities after G-transitions, particularly during the early days of flight, to allow the crew to adapt to SMS. Examples of rules are prohibition of extravehicular activities until the third day on-orbit because of concerns related to emesis in the spacesuits, and restriction of driving or flying until the third day after short-duration flights. When Shuttle flights resumed after the loss of Space Shuttle *Columbia*, computerized dynamic posturography testing was implemented as an aid for the return-to-duty assessment to supplement preflight and post-flight neurological examinations.

## **2. RISKS DURING PILOTED LANDINGS**

Piloting a spacecraft through entry and landing is one of the most difficult tasks associated with space flight. The consequences of failing to complete this task successfully could be catastrophic, resulting in loss of life, vehicle, or other assets. While all piloted landings from space have been successful to date, the evidence presented above suggests that the landing performance has been lower than desired for both the Shuttle and the Lunar Lander. To the (currently unknown) extent physiological adaptations play a role in these performance decrements, we can anticipate that the risk of failure will become much greater during Mars missions. There is strong evidence that the six-month outbound trip (without artificial gravity) will cause a much more profound sensorimotor adaptation to 0 g than occurs during a 1-2 week Shuttle mission. This will likely cause a more profound physiological response to the G-transition during entry/landing; however, the impact of the reduced amplitude (3/8 g vs. 1 g) of the transition is unknown. Furthermore, piloting recency will decrease from 1-2 weeks during the Shuttle program to approximately six months during a Mars mission, decreasing the probability that a pilot will be able to fly through any spatial disorientation that accompanies the G-transition. Even piloted landings on the Moon present some unique risks, owing to the effects of the novel gravitational environment on spatial and geographic orientation and the potential for lunar dust obscuring vision during critical phases of landing.

## **3. RISKS DURING VEHICLE EGRESS AND EXTRAVEHICULAR ACTIVITIES**

Ensuring that crewmembers are able to egress the vehicle in the event an emergency occurs during the post-landing timeframe is essential to allowing them to survive or avoid serious injury during such an event. The crewmembers should also be able to function in 1-g environment in case of return to Earth, or in a 1/6-g or 3/8-g environment in case of lunar or Mars landing. Factors that affect egress in a timely manner include (a) visually determining hazards outside the vehicle, such as the presence of fire or debris; (b) having a hatch that can be operable by a single crewmember without the use of tools; and (c) having an egress path that allow egress of all occupants in a timely manner. Determining if it is safe to egress the vehicle and having an egress path requires good situation awareness, spatial orientation, and mental representation of space. Opening the hatch, egressing the vehicle, and walking in enough time

to protect from post-landing hazards may be compromised if crewmembers are incapacitated or in a deconditioned state. Orion and other commercial vehicles are currently designed for a parachuted landing on water after long-duration missions. In these water landing scenarios, the interaction between the adapted microgravity sensorimotor state and the prevailing unstable support surface induced by various sea state conditions will increase the risk associated with an emergency egress situation.

During Expedition 6 to the ISS, a series of unplanned events serendipitously created an analog mission for a trip to and landing on Mars. A spacecraft malfunction causing a ballistic entry which displaced the landing site about 475 kilometers off course resulting in about a 5 hour delay for arrival of the ground support team. This gave the crew an opportunity to perform spacecraft safing, egress, and set up survival gear without any outside help (Pettit 2010). The crew performed spacecraft safing, involving reading procedures, flipping switches, and pushing buttons on the control panel to power down unneeded equipment so that battery life for radio operations can be extended. Since the Soyuz capsule ended on its side, these operations were done from a position of being strapped into a seat fixed on a slanted ceiling. The crew opened the hatch, unstrapped, and crawled out. Following egress, they deployed the survival gear that was stowed in numerous small bundles throughout the spacecraft. Included were warm woollen cloths, food, water, a medical kit, a portable radio, and a signalling kit.

One crewmember reported that “performing these basic survival tasks was not easy. Moving was provocative. [...] Walking was labored but was done as needed shortly after landing. I had trouble walking but could crawl. [...] There were no systemic aches or pains associated with movement. We had good muscle strength. [...] My limbs felt heavy because my brain was not yet compensating for their weight. [...] Upon returning, the brain had not yet kicked in this compensation which takes about 10 to 15 hours. Slow and deliberate motions were readily made with sufficient motor control to connect electrical wire harnesses, antennas, cycle switches on control panels, and shoot a shotgun pistol. Motor control for operating the spacecraft mechanisms and survival gear was not a problem. However, fast coordinated movement was not possible for me.” (Pettit 2010). In addition, this crewmember recommended that “a well-designed [Mars] mission should have minimal demands on the crew after landing, giving them a few days for adaptation before engaging in significant operational tasks.”

#### **4. RISKS DURING ROVER OPERATIONS**

The risk of performance failure (i.e., loss of vehicle control) while driving an automobile is high for those having vestibular deficiencies and for those whose cognitive and/or sensorimotor functions are impaired by ethanol, fatigue, or certain medications. Crewmembers readapting to Earth-gravity following return from space flight exhibit similar performance decrements, and, as a result, are currently restricted from driving automobiles for a short time (2-4 days) after Shuttle missions and a longer time (8-12 days) after ISS missions. The impact of sensorimotor adaptations on driving rovers on either Moon or Mars is unknown. While the potential consequences of performance failure while driving a rover are less than those of piloting a space craft through entry and landing, the possibility of crew injury (or death) or loss of the

rover exists, particularly in the vicinity of steep-sided craters. The duration of the initial adaptation period to the Lunar or Martian gravity environment is also unknown, and, while likely to be proportional to the time spent in 0-g transit, cannot be determined until it can be measured on the planetary surface. Thus, the amplitude and duration of increased risk during rover driving are currently unknown.

## **5. RISKS DURING RENDEZVOUS/DOCKING AND REMOTE MANIPULATOR SYSTEM OPERATIONS**

Apart from the Spektr incident, performance data on rendezvous/docking has so far eluded the authors. However, evidence provided above suggests that the incidence of performance failure during remote manipulator operations aboard the Shuttle and ISS has been fairly well characterized (at least operationally). There is no reason to suspect that performance of these 0-g operations will be any different from our ISS experience during an outbound transit to Mars. Thus, we would not expect the risk to increase. However, the risk impacts of an additional 18 months at Mars gravity followed by six months at 0 g during return transit are unknown, and may well lead to an unacceptable range.

## **6. RISKS WHILE OPERATING OTHER COMPLEX SYSTEMS**

The risk of performance failure during operation of any complex system is multi-factorial. However, operation of any system requiring good visual acuity, eye-hand coordination, (balance/locomotor skills for surface operations), spatial orientation, and/or cognition could be impaired by physiological adaptations to novel gravitational environments. The risk of impairment is generally greatest during and soon after G-transitions, but the amplitude and duration of the increased risk would need to be evaluated on a system-by-system basis.

## **7. RISKS DURING NEAR-TERM MISSIONS**

The current sensorimotor risk is now high in priority given the prospect of long-duration Mars missions and planned water landings following long-duration ISS missions. There is significant evidence showing sensorimotor alteration after as little as a few weeks of exposure to space flight environments, and that severity increases with increasing exposure time. While these issues may be more severe for Mars missions without artificial gravity, significant risks remain quite real even for more standard ISS and Lunar operations. As the Columbia Accident Investigation Board report warned us repeatedly, a small number of successes without catastrophic failure (*e.g.*, a little over 100 Shuttle landings and 6 lunar landings) does not mean that risk, including human sensorimotor adaptation risks, can be ignored. The near misses reported above provide evidence in this regard. Given the reentry profiles and cross-coupled Coriolis effects induced by the drogue parachutes under nominal Orion re-entry/descent/landing scenarios it will likely be a much more difficult sensorimotor environment than for Shuttle landings. Finally, the proper resolution of automation-human control authority decisions requires an objective and quantitative understanding of sensorimotor compromises. The risk of sub-optimal decisions in this regard has important ramifications for overall mission safety/reliability calculations. Thus, we recommend that this risk be considered high priority for all space flight mission scenarios.

## VIII. GAPS

The authors, representing the sensorimotor (SM) discipline team, have identified the series of knowledge and mitigation gaps listed below. Each of them must be filled before this risk can be fully assessed and/or mitigated.

**SM2.1** – Determine the changes in sensorimotor function over the course of a mission and during recovery after landing.

**SM2.2** – Determine the effects of long-duration space flight on sensorimotor function over a crewmember’s lifetime.

**SM6.1** – Determine if sensorimotor dysfunction during and after long-duration space flight affects ability to control spacecraft and associated system.

**SM7.1** – Determine if there are decrements in performance on functional tasks after long-duration space flight. Determine how changes in physiological function, exercise activity, and/or clinical data account for these decrements.

**SM24** – Determine if the individual capacity to produce adaptive change (rate and extent) in sensorimotor function to transitions in gravitational environments can be predicted with preflight tests of sensorimotor adaptability.

**SM26** – Determine if exposure to long-duration space flight leads to neural structural alterations and if this remodeling impacts cognitive and functional performance.

**SM27** – Determine the most optimal pharmacological and sensorimotor countermeasure combination that reduces Space Motion Sickness (SMS) while minimizing side effect.

**SM28** – Develop a sensorimotor countermeasure system integrated with current exercise modalities to mitigate performance decrements during and after space flight.

## IX. CONCLUSION

A large body of sensorimotor research data obtained from space flight experiments over the past half-century demonstrates significant decrements in oculomotor control, eye-hand coordination, spatial orientation, posture\locomotor control and cognition during space flight missions. While these changes are most severe during and after G-transitions, the most crucial time for many critical operational tasks (e.g., landing and egress), only limited information is available to assess the operational impacts of these changes. Some of the operational observations are compelling, but are confounded by unknown environmental and engineering influences. Others appear to raise little concern, but the safety margins are difficult to estimate. During exploration missions, we can expect that most performance circa G-transitions will be degraded further by the influence of extended time in flight (Mars missions), but the potential influence of extended time in hypogravity (Mars and Lunar missions) is unknown.

The true operational risks associated with the impacts of adaptive sensorimotor (and other) changes on crew mobility and abilities to control vehicles and other complex systems will only be estimable after the gaps (identified above) have been filled and we have been able to accurately assess integrated performance in off-nominal operational settings. While exclusive crew selection procedures, intensive crew training, and highly reliable hardware/software systems have likely minimized the operational impacts of these sensorimotor changes to date, the impacts of new mission and vehicle designs may offset some of benefits.

Forward work in this area must account for the multi-factorial nature of the problem. While sensorimotor and behavioral (cognitive) disciplines clearly have roles to play, muscle (strength and endurance) and cardiovascular (orthostatic tolerance) disciplines also must be involved, as should human factors experts, training experts, vehicle designers, mission designers, and crewmembers. Mechanisms for facilitating cross-disciplinary investigations are only beginning to be established. Future success will clearly require more progress in these approaches.

## X. REFERENCES

- Aihara, T., Kitajo, K., Nozaki, D., and Yamamoto, Y. (2010). How does stochastic resonance work within the human brain? – Psychophysics of internal and external noise. *Chemical Physics* 375, 616-624. doi: 10.1016/j.chemphys.2010.04.027
- Aizikov GS, Kreidich YuV, Grigoryan RA (1991) Sensory interaction and methods of non-medicinal prophylaxis of space motion sickness. *Physiologist* 34: S220-223
- Albery W, Repperger D (1992) Time and mass perception in non-terrestrial environments. *Acta Astronautica* 26: 119-126
- Anderson DJ, Reschke MF, Homick JL, *et al.* (1986) Dynamic posture analysis of Spacelab crewmembers. *Exp Brain Res* 64: 380-391
- André-Deshays C, Israel I, Charade O, Berthoz A, Popov K, Lipshits M (1993) Gaze control in microgravity. 1. Saccades, pursuit, eye-head coordination. *J Vestib Res* 3: 331-343
- Angelaki DE, McHenry MQ, Dickman JD, Newlands SD, Hess BJ (1999) Computation of inertial motion: neural strategies to resolve ambiguous otolith information. *J Neurosci* 19: 316-327
- Antunano MJ, Hernandez JM (1989) Incidence of airsickness among military parachutists. *Aviat Space Environ Med* 60: 792-797
- Aoki H, Oman CM, Natapoff A (2007) Virtual-reality-Based 3D navigation training for emergency egress from spacecraft. *Aviat Space Environ Med* 78: 774-783, 2007
- Aoki H, Oman CM, Natapoff A, Liu A (2006) The effect of the configuration, frame of reference, and spatial ability on spatial orientation during virtual 3-dimensional navigation training. In: ESTEC (ed) *7th Symposium on the Role of the Vestibular Organs in Space Exploration*. ESA: Noordwijk, NL
- Armstrong PA, Wood SJ, Shimizu N, Kuster K, Perachio A, Makishima T (2015) Preserved otolith organ function in caspase-3-deficient mice with impaired horizontal semicircular canal function. *Exp Brain Res* 233: 1825-1835
- Arrott AP, Young LR (1986) M.I.T./Canadian vestibular experiments on the Spacelab-1 mission: 6. Vestibular reactions to lateral acceleration following ten days of weightlessness. *Exp Brain Res* 64: 347-357.
- Arrott AP, Young LR, Merfeld DM (1990) Perception of linear acceleration in weightlessness. *Aviat Space Environ Med* 61: 319-326
- Arya M, Paloski WH, Young LR (2007) Centrifugation protocol for the NASA artificial gravity-bed rest pilot study. *J Gravit Physiol* 14: P5-P8
- Arzeno NM, Stenger MB, Bloomberg JJ, Platts SH (2013) Spaceflight-induced cardiovascular changes and recovery during NASA's Functional Task Test. *Acta Astronautica* 92: 10-14
- Ashkenas IL, Ho RH, Teper GL (1983) Analyses of Shuttle Orbiter Approach and Landing. *J Guid Control Dyn* 6: 448-455
- Assaiante C, Amblard B (1993) Ontogenesis of head stabilization in space during locomotion in children: influence of visual cues. *Exp Brain Res* 93: 499-515
- Attias J, Gordon C, Ribak J, *et al.* (1987) Efficacy of transdermal scopolamine against seasickness: a 3-day study at sea. *Aviat Space Environ Med* 58: 60-62
- Bacal K, Billica R, Bishop S (2003) Neurovestibular symptoms following space flight. *J Vestib Res* 13: 93-102

- Bagian JP (1991) First intramuscular administration in the U.S. space program. *J Clin Pharmacol* 31: 920
- Bagian JP, Ward DF (1994) A retrospective study of promethazine and its failure to produce the expected incidence of sedation during space flight. *J Clin Pharmacol* 34: 649-651
- Baker JT, Nicogossian AE, Hoffler GW, *et al.* (1977) Changes in the Achilles tendon reflexes following Skylab missions. In: Johnston RS, Dietlein LF (eds) *Biomedical Results from Skylab*. NASA: Washington DC, NASA SP-377, pp 131-135
- Balaban C (1996) The role of the cerebellum in vestibular autonomic regulation. In: Yates BJ, Miller AD (eds) *Vestibular Autonomic Regulation*. CRC Press: Boca Raton, pp 127-144
- Baron S (1983) An Optimal Control Model Analysis of Data from a Simulated Hover Task.
- Baron S, Kleinman DL (1969) The human as an optimal controller and information processor. *Man Mach Sys* 10: 9-17
- Barrett R, Lokhandwala M (1981) Presynaptic dopamine receptor stimulation in the cardiovascular actions of lergotriole. *J Pharmacol Exp Ther* 217: 660-665
- Barron KD, Chokroverty S (1993) Anatomy of the autonomic nervous system: brain and brainstem. In: Low PA (ed) *Clinical Autonomic Disorders: Evaluation and Management*. Little Brown: Boston, pp 3-15
- Basner M, Savitt A, Moore TM, Port AM, McGuire S, Ecker AJ, Nasrini J, Mollicone DJ, Mott CM, McCann T, Dinges DF, Gur RC (2015) Development and validation of the Cognition test battery for spaceflight. *Aerosp Med Hum Perform* 86: 942–952.
- Batson CD, Brady RA, Peters B, Ploutz-Snyder R, Mulavara A, Cohen HS, Bloomberg JJ (2011) Gait training improves performance in healthy adults exposed to novel sensory discordant conditions. *Exp Brain Res* 209: 515-524
- Beaton KH, Huffman WC, Schubert MC (2015) Binocular misalignments elicited by altered gravity provide evidence for nonlinear central compensation. *Front Syst Neurosci* 9: 81. Doi 10.3389/fnsys.2015.00081
- Becker G, Goossens H, Seemann K, *et al.* (1984) Prevention of motion sickness with a transdermal therapeutic system containing scopolamine: a randomized, comparative double-blind study in the German Federal Navy. *Dtsch Med Wochenschr* 109: 1881-1885
- Benson AJ, Barnes GR (1978) Vision during angular oscillation: the dynamic interaction of visual and vestibular mechanisms. *Aviat Space Environ Med* 49: 340-345
- Benson AJ, Kass JR, Vogel H (1986) European vestibular experiments on the Spacelab-1 mission: 4. Thresholds of perception of whole-body linear oscillation. *Exp Brain Res* 64: 264-271
- Benson AJ, Vieville T (1986) European vestibular experiments on the Spacelab-1 mission: 6. Yaw axis vestibulo-ocular reflex. *Exp Brain Res* 64: 279-283
- Benson AJ, Guedry FE, Parker DE, Reschke MF (1997) Microgravity vestibular investigations: perception of self-orientation and self-motion. *J Vestib Res* 7: 453-457
- Berger M, Ferstenbrand F, Kozlovskaya IB, *et al.* (1992) Eye head and arm coordination and spinal reflexes in weightlessness — Monimir experiment. In: ASFA Medicine (ed) *Health from Space Research. Austrian Accomplishments*. Springer-Verlag: Wien, pp 119-135
- Berger M, Mescheriakov S, Molokanova E, *et al.* (1997) Pointing arm movements in short- and long-term space flights. *Aviat Space Environ Med* 68: 781-787
- Berrey MM, Shea T, Corey L (1998) Cotton-wool spots in primary HIV infection (letter). *J Acquir Immune Defic Syndr Hum Retrovirol* 19:197-198

- Berry CA, Homick JL (1973) Findings on American astronauts bearing on the issue of artificial gravity for future manned space vehicles. *Aerospace Med* 44: 163-168
- Berthoz A, Brandt T, Dichgans J, Probst T, Bruzek W, Vieville T (1986) European vestibular experiments on the Spacelab-1 mission: 5. Contribution of the otoliths to the vertical vestibulo-ocular reflex. *Exp Brain Res* 64: 272-278
- Berthoz A, Melvill Jones G (1985) *Adaptive Mechanisms in Gaze Control*. Elsevier Science Publishers: Amsterdam
- Bloomberg JJ, Mulavara AP (2003) Changes in walking strategies after space flight. *IEEE Eng Med Biol Mag* 22: 58-62
- Bloomberg JJ, Reschke MF, Huebner WP, Peters BT (1992) The effects of target distance on head movement during locomotion. *Ann NY Acad Sci* 656: 699-707
- Bloomberg JJ, Peters BT, Smith SL, Huebner WP, Reschke MF (1997) Locomotor head-trunk coordination strategies following space flight. *J Vestib Res* 7: 161-177
- Bloomberg JJ, Feiveson AH, Laurie S, Lee SMC, Mulavara AP, Peters BT, Platts SH, Ploutz-Snyder LL, Reschke MF, Ryder JW, Stenger MB, Taylor LC, Wood SJ (2015a) NASA's Functional Task Test: Providing information for an integrated countermeasure system. *Proceedings of the 20<sup>th</sup> IAA Humans in Space Symposium*, Prague, 28 June-3 July 2015, IAA-HIS-15-1A1-2
- Bloomberg JJ, Peters BT, Cohen HS and Mulavara AP (2015b) Enhancing astronaut performance using sensorimotor adaptability training. *Front. Syst. Neurosci.* 9:129.doi: 10.3389/fnsys.2015.00129.
- Bock O (1996) Grasping of virtual objects in changed gravity. *Aviat Space Environ Med* 67: 1185-1189
- Bock O, Cheung BSK (1998) Control of isometric force in hypergravity. *Aviat Space Environ Med* 69: 27-31
- Bock O, Howard IP, Money KE, *et al.* (1992) Accuracy of aimed arm movements in changed gravity. *Aviat Space Environ Med* 63: 994-998
- Bock O, Arnold KE, Cheung BSK (1996a) Performance of a simple aiming task in hypergravity: I. Overall accuracy. *Aviat Space Environ Med* 67: 127-132
- Bock O, Arnold KE, Cheung BSK (1996b) Performance of a simple aiming task in hypergravity: II. Detailed response characteristics. *Aviat Space Environ Med* 67: 133-138
- Bock O, Abeele S, *et al.* (2003) Human adaptation to rotated vision: interplay of a continuous and a discrete process. *Exp Brain Res* 152: 528-532
- Bock O, Weigelt C, Bloomberg JJ (2010) Cognitive demand of human sensorimotor performance during an extended space mission: A dual-task study. *Aviat Space Environ Med* 81: 819-824
- Boeing (2007) *Statistical Summary of Commercial Jet Airplane Accidents in Worldwide Operations 1959-2006*. Boeing Commercial Airplanes: Seattle WA
- Boff K, Lincoln J (1988) *Engineering data compendium: human perception and performance* Wright-Patterson, AFB Ohio: Harry G. Armstrong Aerospace Medical Research Laboratory
- Bonan IV, Colle FM, Guichard JP, Vicaut E, Eisenfisz M, Tran Ba Huy P, Yelnik, AP (2004) Reliance on visual information after stroke. Part I: Balance on dynamic posturography. *Arch. Phys. Med. Rehabil* 85: 268-273
- Borah J, Young LR, Curry RE (1979) Optimal estimator model for human spatial orientation. *IEEE Trans Systems*

- Bortolami SB, Rocca S, Daros S, DiZio P, Lackner JR (2006a) Mechanisms of human static spatial orientation. *Exp Brain Res* 173: 374-388
- Bortolami SB, Pierobon A, DiZio P, Lackner JR (2006b) Localization of the subjective vertical during roll, pitch, and recumbent yaw body tilt. *Exp Brain Res* 173: 364-73
- Bortolami SB, Pigeon P, DiZio P, Lackner JR (2008) Dynamics model for analyzing reaching movements during active and passive torso rotation. *Exp Brain Res* 187: 525-534
- Bokhov BB, Danilov KY, Makarov SI (1969) Investigation of human perception of the coordinates of a two-dimensional space using the VERTIKAL device. *Dev Metho Athl Tra in Exp* 176-177
- Bokhov BB, Yakovleva IY, Kornilova LN, *et al.* (1973) The effect of vestibular tests on the orientation of the subjective vertical in intact and deaf-mute subjects. *Kosm Biol Aviakosm Med* 6: 51-56
- Boyle R, Mesinger AF, Yoshida K, Usui S, Intravaia A, Tricas T, Highstein SM (2001) Neural readaptation to earth's gravity following return from space. *J Neurosci* 86: 2118-2122
- Boyle R, Popova Y, Varelis J (2015) Influence of Gravity on Mouse Inner Ear Otoconia. *Proceedings of the 2014 Human Research Program Investigators' Worksop*, Galveston, 13-15 January 2015
- Brady RA, Peters BT, Bloomberg JJ (2009) Strategies of healthy adults walking on a laterally oscillating treadmill. *J Gait Posture* 29: 645-649
- Brady RA, Peters BT, Batson CD, Ploutz-Snyder R, Mulavara AP, Bloomberg JJ (2012) Gait adaptability training is affected by visual dependency. *Exp Brain Res* 220: 1-9
- Braithwaite MG, Beal KG, Alvarez EA, Jones HD, Estrada A (1998) The optokinetic cervico reflex during simulated helicopter flight. *Aviat Space Environ Med* 69: 1166-1173
- Braun DA, Mehring C, Wolpert DM (2010) Structure learning in action. *Behav Brain Res* 20; 206(2):157-65
- Bryan AS, Bortolami SB, Ventura J, DiZio P, Lackner JR (2007) Influence of gravitoinertial force level on the subjective vertical during recumbent yaw axis body tilt. *Exp Brain Res* 183: 389-397
- Bryanov II, Gorgiladze GI, Kornilova LN, *et al.* (1986) Vestibular function. In: Gazenko OG (ed) *Results of Medical Research Performed on the Salyut 6-Soyuz Orbital Scientific Research Complex*. Meditsina: Moscow, pp 169-185, 248-256
- Bryanov II, Yemel'yanov MD, Matveyev AD, *et al.* (1976) *Space Flights in the Soyuz Spacecraft: Biomedical Research*. Leo Kanner Associates: Redwood City
- Buccello-Stout, RR, Bloomberg, JJ, Cohen, HS, Whorton, EB, Weaver, GD, & Cromwell, RL (2008) Effects of sensorimotor adaptation training on functional mobility in older adults. *J Gerontol B Psychol Sci Soc Sci.* 63(5): 295-300. 2008
- Buckey JC, Lane LD, Levine BD, *et al.* (1996) Orthostatic intolerance after space flight. *J Appl Physiol* 81: 7-18
- Buckey JC, Lane LD, Plath G, *et al.* (1992) Effects of head-down tilt for 10 days on the compliance of the leg. *Acta Physiol Scand Suppl* 604: 53-59
- Bungo MW, Charles JB, Johnson PC (1985) Cardiovascular deconditioning during space flight and the use of saline as a countermeasure to orthostatic intolerance. *Aviat Space Environ Med* 56: 985-990
- Burrough B (1988) *Dragonfly: NASA and the Crisis Aboard Mir*. Harper-Collins: New York, NY

- Campbell MR, Williams DR, Buckey JC, Jr., Kirkpatrick AW (2005) Animal surgery during space flight on the Neurolab Shuttle mission. *Aviat Space Environ Med* 76: 589-593
- Carpenter RD, Lang TF, Bloomfield SA, Bloomberg JJ, Judex S, Keyak JH, Midura RJ, Pajevic PD, Spatz JM (2010) Effects of long-duration spaceflight, microgravity, and radiation on the neuromuscular, sensorimotor, and skeletal systems. *J Cosmology* 12: 3778- 3780
- Carr CE, Newman DJ (2007a) Space suit bioenergetics: framework and analysis of unsuited and suited activity. *Aviat Space Environ Med* 78: 1013-1022
- Carr CE, Newman DJ (2007b) Space suit bioenergetics: cost of transport during walking and running. *Aviat Space Environ Med* 78: 1093-1102.
- Carriot J, Jamali M, Cullen KE (2015) Rapid adaptation of multisensory integration in vestibular pathways. *Front Syst Neurosci* 9: 59
- Charles JB, Lathers CM (1991) Cardiovascular adaptation to space flight. *J Clin Pharmacol* 31: 1010-1023
- Chekirda IF, Bogdashevskiy AV, Yeremin AV, Kolosov IA (1971) Coordination structure of walking of Soyuz-9 crewmembers before and after flight. *Kosm Biol Aviakosm Med* 5: 48-52
- Chekirda IF, Yeremin AV (1977) Dynamics of cyclic and acyclic locomotion of the Soyuz-18 crew after a 63-day space mission. *Kosm Biol Aviakosm Med* 4: 9-13
- Chess GF, Tam RMK, Calarescu FR (1975) Influence of cardiac neural inputs on rhythmic variations of heart period in the cat. *Am J Physiol* 228: 775-780
- Clark JB, Rupert AH (1992) Spatial disorientation and dysfunction of orientation/equilibrium reflexes: aeromedical evaluation and considerations. *Aviat Space Environ Med* 63: 914-918
- Clark JB, Bacal K (2008) Neurologic concerns. In: *Principles of Clinical Medicine for Space Flight*. Barratt MR, Pool SL (eds) Springer, New York, pp 361-380
- Clark TK, Stimpson AJ, Young LR, Oman CM, Duda KR, Natapoff A (2014) Human spatial orientation perception during simulated lunar landing motions. *J Spacecrafts Rockets* 51: 267-280
- Clark TK, Newman MC, Oman CM, Merfeld DM, Young LR (2015) Modeling human perception of orientation in altered gravity. *Front Syst Neurosci* 9: 68
- Clarke AH, Kornilova LN (2007) Ocular torsion response to active head-roll movement under one-g and zero-g conditions. *J Vestib Res* 17: 99-111
- Clarke AH, Teiwes W, Scherer H (1993a) Evaluation of the torsional VOR in weightlessness. *J Vestib Res* 3: 207-218
- Clarke AH, Teiwes W, Scherer H (1993b) Vestibulo-oculomotor testing during the course of a space flight mission. *Clin Investig* 71: 740-748
- Clarke AH, Grigull J, Mueller R, Scherer H (2000) The three-dimensional vestibulo-ocular reflex during prolonged microgravity. *Exp Brain Res* 134: 322-334
- Clarke AH, Schonfeld U, Wood SJ (2010) The OTOLITH experiment – Assessment of unilateral otolith function during postflight re-adaptation. *J Gravit Physiol* 17: 11-12
- Clarke AH, Just K, Krzok W, Schonfeld U (2013) Listing's plane and the 3D-VOR in microgravity – The role of the otolith afferences. *J Vestib Res* 23: 61-70
- Clément G (2011) *Fundamentals of Space Medicine*. Second Edition. Springer: New York
- Clément G, Berthoz A (1988) Vestibulo-ocular reflex and optokinetic nystagmus in microgravity. *Adv Oto-Rhino-Laryngol* 42: 1-4

- Clément G, Lestienne F (1988) Adaptive modifications of postural attitude in conditions of weightlessness. *Exp Brain Res* 72: 381-389
- Clément G, Berthoz A (1990) Cross-coupling between horizontal and vertical eye movements during optokinetic nystagmus and optokinetic afternystagmus elicited in microgravity. *Acta Otolaryngol* (Stockh) 109: 179-187
- Clément G, Reschke MF (1996) Neurosensory and sensory-motor functions. In: Moore D, Bie P, Oser H (eds) *Gravitational and Space Biology: An Overview of Life Sciences Research in Microgravity*. Springer-Verlag: Heidelberg, Chapter 4, pp 178-258
- Clément G, Eckardt J (2005) Influence of the gravitational vertical on geometric visual illusions. *Acta Astronautica* 56: 911-917
- Clément G, Denise P, Reschke MF, Wood SJ (2007) Human ocular counter-rolling and roll tilt perception during off-vertical axis rotation after spaceflight. *J Vestib Res* 17(5-6):209-215
- Clément G, Reschke MF (2008) *Neuroscience in Space*. Springer, New York
- Clément G, Wood SJ (2013) Eye movements and motion perception during off-vertical axis rotation after space flight. *J Vestib Res* 23: 13-22
- Clément G, Wood SJ (2014) Rocking or rolling – Perception of ambiguous motion after returning from space. *PLoS One* 9(10): e111107. doi:10.1371/journal.pone.0111107
- Clément G, Gurfinkel VS, Lestienne F, et al. (1984) Adaptation of posture control to weightlessness. *Exp Brain Res* 57: 61-72
- Clément G, Gurfinkel VS, Lestienne F, et al. (1985) Changes of posture during transient perturbations in microgravity. *Aviat Space Environ Med* 56: 666-671
- Clément G, Viéville T, Lestienne F, Berthoz A (1986) Modifications of gain asymmetry and beating field of vertical optokinetic nystagmus in microgravity. *Neurosci Lett* 63: 271-274
- Clément G, Berthoz A, Lestienne F (1987) Adaptive changes in perception of body orientation and mental-image rotation in microgravity. *Aviat Space Environ Med* 58 (Suppl 9): A159-A163
- Clément G, Wood S, Reschke M, Berthoz A, Igarashi M (1999) Yaw and pitch visual-vestibular interaction in weightlessness. *J Vestib Res* 9: 207-220
- Clément G, Moore ST, Raphan T, Cohen B (2001) Perception of tilt (somatogravic illusion) in response to sustained linear acceleration during space flight. *Exp Brain Res* 138: 410-418
- Clément G, Frayss MJ, Deguine O (2009) Mental representation of space in vestibular patients with otolithic or rotatory vertigo. *NeuroReport* 20: 457-461
- Clément G, Skinner A, Richard G, Lathan C (2012) Geometric illusions in astronauts during long-duration space flight. *NeuroReport* 23: 894-899
- Clément G, Skinner A, Lathan C (2013) Distance and size perception in astronauts during long-duration space flight. *Life* 3: 524-537
- Clément G, Allaway HCM, Demel M, Golemis A, Kindrat AN, Melinyshyn AN, Merali T, Thirsk R (2015a) Long-duration space flight increases depth ambiguity of reversible perspective figures. *PLoS One* 10(7): e0132317. doi:10.1371/journal.pone.0132317
- Clément G, Buckley AP and Paloski WH (2015b) Artificial gravity as a countermeasure for mitigating physiological deconditioning during long-duration space missions. *Front. Syst. Neurosci.* 9:92. doi: 10.3389/fnsys.2015.00092
- Cohen B, Matsuo V, Raphan T (1977) Quantitative analysis of the velocity characteristics of optokinetic nystagmus and optokinetic after-nystagmus. *J Physiol* 270: 321-344

- Cohen HS, Wells J, Kimball KT, Owsley C (2003) Driving disability and dizziness. *J Safety Res* 34: 361-369
- Cohen HS, Bloomberg JJ, Mulavara AP (2005) Obstacle avoidance in novel visual environments improved by variable practice training. *Percept Mot Skills* 101: 853-861
- Cohen HS, Kimball KT, Wood SJ, Paloski WP, Bloomberg JJ, Mulavara AP (2012a) Posturography and locomotor tests of dynamic balance after long-duration spaceflight. *J Vestib Res* 22: 191-196
- Cohen HS, Mulavara AP, Peters BT, Sangi-Haghpeykar H, Bloomberg JJ (2012b) Tests of walking balance for screening vestibular disorders. *J Vestib Res* 22: 95-104
- Cohen HS, Mulavara AP, Peters BT, Sangi-Haghpeykar H, Kung DH, Mosier DR, Bloomberg JJ (2013) Sharpening the tandem walking test for screening peripheral neuropathy. *South Med J* 106: 565-569
- Cohen HS, Mulavara AP, Peters BT, Sangi-Haghpeykar H, Bloomberg JJ (2014) Standing balance tests for screening people with vestibular impairments. *Laryngoscope* 124:545-550
- Cohen MM. Perception of facial features and face-to-face communications in space (2000) *Aviat Space Environ Med.* 2000 Sep;71(9 Suppl):A51-7
- Collins JJ, Priplata AA, Gravelle DC, Niemi J, Harry J, Lipsitz LA (2003) Noise enhanced human sensorimotor function. *IEEE Engineering in Medicine and Biology Magazine.* 2003;22(2):76-83
- Collins, J., Imhoff, T., and Grigg, P. (1996a). Noise-enhanced information transmission in rat SA1 cutaneous mechanoreceptors via aperiodic stochastic resonance. *J Neurophysiol* 76, 642-645
- Collins, J., Imhoff, T., and Grigg, P. (1996b). Noise-enhanced tactile sensation. *Nature* 383, 770.
- Collins, J.J., Imhoff, T.T., and Grigg, P. (1997). Noise-mediated enhancements and decrements in human tactile sensation. *Phys Rev E* 56, 923-926
- Collins, J.J., Priplata, A.A., Gravelle, D.C., Niemi, J., Harry, J., and Lipsitz, L.A. (2003). Noise enhanced human sensorimotor function. *IEEE Engineering in Medicine and Biology Magazine* 22, 76-83
- Convertino VA, Previc FH, Ludwig DA, Engelken EJ (1997) Effects of vestibular and oculomotor stimulation on responsiveness of the carotid-cardiac reflex. *Am J Physiol* 273: R615-R622
- Cooper HSFJ (1976) *A House in Space*. Holt, Rhinehart & Winston: New York
- Corballis MC, Zbrodoff NJ, Shetzer LI, Butler PB (1978) Decisions about identity and orientation of rotated letters and digits. *Memory and Cognition* 6: 98-107
- Costa VP, Arcieri ES (2007) Hypotony maculopathy. *Acta Ophthalmol Scand* 85: 586-597
- Courtine G, Papaxanthis C, Pozzo T (2002) Prolonged exposure to microgravity modifies limb endpoint kinematics during the swing phase of human walking. *Neurosci Lett* 332: 70-74
- Courtine G, Pozzo T (2004) Recovery of the locomotor function after prolonged microgravity exposure. I. Head-trunk movement and locomotor equilibrium during various tasks. *Exp Brain Res* 158: 86-99
- Cowings PS (1990) Autogenic-feedback training: a treatment for motion and space sickness. In: Crampton GH (ed) *Motion and Space Sickness*. CRC Press: Boca Raton, pp 353-372
- Cowings PS, Suter S, Toscano WB, et al. (1986) General autonomic components of motion sickness. *Psychophysiology* 23: 542-551

- Cramer SC, Nelles G, Benson RR, Kaplan JD, Parker RA, Kwong KK, Kennedy DN, Finklestein SP, Rosen BR (1997) A functional MRI study of subjects with motor recovery after stroke. *Stroke* 28: 2518-2527
- Cucinotta FA, Alp M, Sulzman FM, Wang M (2014) Space radiation risks to the central nervous system. *Life Sci Space Res* 2: 54-69
- Currie NJ, Peacock B (2001) International Space Station robotic systems operations - A human factors perspective. In: HFES (ed) *Human Factors and Ergonomics Society Annual Meeting*. Baltimore, MD
- Curry IP, McGhee JS (2007) Spatial disorientation as a cause of accidents in combat helicopter operations. *Aviat Space Environ Med* 78: 307-311
- Curry RE, Young LR, Hoffman WC, Kugel DL (1976). A pilot model with visual and motion cues. *Proceedings of the 12th Annual Conference on Manual Control*, Urbana, Illinois, May 1976, NASA TM X-73170: 25-27
- Dai M, McGarvie L, Kozlovskaya I, Raphan T, Cohen B (1994) Effects of space flight on ocular counterrolling and the spatial orientation of the vestibular system. *Exp Brain Res* 102: 45-56
- Dai M, Raphan T, Kozlovskaya I, Cohen B (1996) Modulation of vergence by off-vertical yaw axis rotation in the monkey: Normal characteristics and effects of space flight. *Exp Brain Res* 111: 21-29
- Dai M, Sofroniou S, Kunin M, Raphan T, Cohen B (2010) Motion sickness induced by off-vertical axis rotation (OVAR). *Exp Brain Res* 204: 207-222
- D'Amelio F, Fox RA, Wu LC, Daunton NG, Corcoran ML (1998) Effects of microgravity on muscle and cerebral cortex: a suggested interaction. *Adv Space Res* 22: 235-244
- Daum SO, Hecht H (2009) Distance estimation in vista space. *Att Percept Psychophys* 71: 1127-1137
- Davids K, Glazier P, Araujo D, Bartlett R (2003) Movement systems as dynamical systems: the functional role of variability and its implications for sports medicine. *Sports Med* 33: 245-260
- Davis JR, Jennings RT, Beck BG (1993a) Comparison of treatment strategies for space motion sickness. *Acta Astronautica* 29: 587-591
- Davis JR, Jennings RT, Beck BG, et al. (1993b) Treatment efficacy of intramuscular promethazine for space motion sickness. *Aviat Space Environ Med* 64: 230-233
- Davis JR, Vanderploeg JM, Santy PA, Jennings RT, Stewart DF (1988) Space motion sickness during 24 flights of the Space Shuttle. *Aviat Space Environ Med* 59: 1185-1189
- de Haart M, Geurts AC, Huidekoper SC, Fasotti L, van Limbeek J (2004) Recovery of standing balance in postacute stroke patients: a rehabilitation cohort study. *Arch Phys Med Rehabil* 85: 886-895
- Demer JL, Amjadi F (1993) Dynamic visual acuity of normal subjects during vertical optotype and head motion. *Invest Ophthalmol Vis Sci* 34: 1894-1906
- Deshpande N, Tourtillott BM, Peters BT, Bloomberg JJ (2013) Dynamic visual acuity (DVA) during locomotion for targets at near and far distances: Effects of aging, walking speed and head-trunk coupling. *J Vestib Res* 23: 195-201
- Diamond SG, Markham CH (1991) Prediction of space motion sickness susceptibility by disconjugate eye torsion in parabolic flight. *Aviat Space Environ Med* 62: 201-205

- Diamond SG, Markham CH (1998) The effect of space missions on gravity-responsive torsional eye movements. *J Vestib Res* 8: 217-231
- Dilda V, Morris TR, Yungher DA, MacDougall HG, Moore ST (2014) Central adaptation to repeated galvanic vestibular stimulation: Implications for pre-flight astronaut training. *PLoS One* 9(11): e112131
- Ditzinger, T., Stadler, M., Struber, D., and Kelso, J.A. (2000). Noise improves three-dimensional perception: stochastic resonance and other impacts of noise to the perception of autostereograms. *Phys Rev E Stat Phys Plasmas Fluids Relat Interdiscip Topics* 62, 2566-2575.
- DiZio P, Lackner JR (1988) The effects of gravito-inertial force level and head movements on post-rotational nystagmus and illusory after-rotation. *Exp Brain Res* 70: 485-495
- DiZio P, Lackner JR (1991) Motion sickness susceptibility in parabolic flight and velocity storage activity. *Aviat Space Environ Med* 62: 300-307
- Doba N, Reis DJ (1974) Role of the cerebellum and the vestibular apparatus in regulation of orthostatic reflexes in the cat. *Circadian Res* 40: 9-18
- Draeger J, Wirt H, Schwartz R (1987) Tonometry under microgravity conditions. In: Sahm PR, Jansen R, Keller MH (eds) *Proceedings of the Norderney Symposium on Scientific Results of the German Spacelab Mission D1*. DFVLR: Norderney, Germany, pp 503-509
- Droulez J, Darlot C (1990) The geometric and dynamic implications of the coherence constraints in three-dimensional sensorimotor interactions. In: Jeannerod M (ed) *Attention and Performance XIII: Motor Representation and Control*. Lawrence Erlbaum Assoc: Hillsdale, NJ, pp 495-526.
- Duda KR, Vasquez RA, Middleton AJ, Hansberry ML, Newman DJ, Jacobs SE, West JJ (2015) The Variable Vector Countermeasure Suit (V2Suit) for space habitation and exploration. *Front Syst Neurosci* 9: 55
- Durnford SJ, Crowley JS, Rosado NR, Harper J, DeRoche S (1995) *Spatial disorientation: A Survey of US Army Helicopter Accidents 1987-1992*.
- Einstein A (1908) Über das Relativitätsprinzip und die aus demselben gezogenen Folgerungen. *Jahrb Radioakt* 4: 411-462
- Eikema DJ, Chien JH, Stergiou N, Myers S, Scott-Pandorf M, Bloomberg J, Mukherjee M. Optic flow improves adaptability of spatiotemporal characteristics during split-belt locomotor adaptation with tactile stimulation. Submitted to *Exp Brain Res* June 2014
- Ellis SR (2000) Collision in space: human factors elements of the Mir-Progress 234 collision. *Ergonomics in Design* 8: 4-9
- Endsley M (1995) Toward a theory of situation awareness in dynamic systems. *Human Factors* 37: 32-64
- Erdeniz B, Rohe T, Done J, Seidler RD (2013) A simple solution for model comparison in bold imaging: the special case of reward prediction error and reward outcomes. *Front Neurosci* 7: 116. Doi 10.3389/fnins.2013.00116
- Essandoh LK, Duprez DA, Shepard JT (1988) Reflex constriction of human limb resistance vessels to head-down neck flexion. *J Appl Physiol* 64: 767-770
- Fernandez C, Goldberg JM (1976) Physiology of peripheral neurons innervating otolith organs of the squirrel monkey. III. Response dynamics. *J Neurophysiol* 39: 996-1008

- Fernandez C, Goldberg JM, Abend WK (1972) Response to static tilts of peripheral neurons innervating otolith organs of the squirrel monkey. *J Neurophysiol* 35: 978-987
- Finke RA, Shepard RN (1986) Visual functions of mental imagery. *Representations* 37: 8
- Fischer CL, Johnson PC, Berry CA (1967) Red blood cell mass and plasma volume changes in manned space flight. *JAMA* 200: 579-583
- Fong KJ, Arya M, Paloski WH (2007) Gender differences in cardiovascular tolerance to short radius centrifugation. *J Gravit Physiol* 14: P15-P19
- Forth KE, Fiedler MJ, Paloski WH (2011) Estimating functional stability boundaries for bipedal stance. *J Gait Posture* 33: 715-717
- Friederici AD, Levelt WJM (1987) Resolving perceptual conflicts: the cognitive mechanism of spatial orientation. *Aviat Space Environ Med* 58: A164-169
- Friederici AD, Levelt WJM (1990) Spatial reference in weightlessness: perceptual factors and mental representations. *Percept Psychophys* 47: 253-266
- Fritsch JM, Charles JB, Bennett BS, Jones MM, Eckberg DL (1992) Short-duration space flight impairs human carotid baroreceptor-cardiac reflex responses. *J Appl Physiol* 73: 664-671
- Fritsch-Yelle JM, Charles JB, Jones MM, Beightol LA, Eckberg DL (1994) Spaceflight alters autonomic regulation of arterial pressure in humans. *J Appl Physiol* 77: 1776-1783
- Fritsch-Yelle JM, Whitson PA, Bondar RL, Brown TE (1996) Subnormal epinephrine release relates to presyncope in astronauts after space flight. *J Appl Physiol* 81: 2134-2141
- Galica, A.M., Kang, H.G., Priplata, A.A., D'andrea, S.E., Starobinets, O.V., Sorond, F.A., Cupples, L.A., and Lipsitz, L.A. (2009). Subsensory vibrations to the feet reduce gait variability in elderly fallers. *Gait Posture* 30, 383-387. doi: 10.1016/j.gaitpost.2009.07.005.
- Gardner CJ, Twissell DJ, Dale TJ, Gale JD, *et al.* (1995) The broad spectrum anti-emetic activity of the novel non-peptide tachykinin NK1 receptor antagonist GR203040. *Br J Pharmacol* 116: 3158-3163
- Gazenko OG, Gernin AM, and Egorov AD (1981) *Major Medical Results of the Salyut-6 Soyuz 18 5-day Space Flight*
- Ghazaleh Samoudi, Maria Jivegård, Ajitkumar P. Mulavara, Filip Bergquist, Effects of Stochastic Vestibular Galvanic Stimulation and LDOPA on Balance and Motor Symptoms in Patients With Parkinson's Disease, *Brain Stimulation*, Volume 8, Issue 3, May–June 2015, Pages 474-480, <http://dx.doi.org/10.1016/j.brs.2014.11.019>.
- Gillingham KK, Wolfe JW (1986) Spatial orientation in flight. In: DeHart R (ed) *Fundamentals of Aerospace Medicine*. Lea and Febiger: Philadelphia, PA, pp 299-381
- Gillingham KK, Previc FH (1996) Spatial orientation in flight. In: DeHart R (ed) *Fundamentals of Aerospace Medicine*. Williams and Wilkins: Baltimore, MD, pp 309-397
- Gillingham KK, Freeman JJ, McNee RC (1977) Transfer functions for eye-level blood pressure during +Gz stress. *Aviat Space Environ Med* 48: 1026-1034
- Ginsberg AP, Vanderploeg JM (1987) Vision in space: near vision acuity and contrast sensitivity. In: Bungo MW, Bagian TM, Bowman MA, Levitan BM (eds) *Results of the Life Sciences DSO Conducted Aboard the Space Shuttle, 1981-1986*. NASA: Houston, TX
- Glasauer S, Amorim MA, Bloomberg JJ, Reschke MF, Peters BT, Smith SL, Berthoz A (1995) Spatial orientation during locomotion following space flight. *Acta Astronautica* 36, 423-431

- Goel R, Kofman I, Jeevarajan J, De Dios Y, Cohen HS, Bloomberg JJ, Mulavara AP (2015) Using low levels of stochastic vestibular stimulation to improve balance function. *PLoS One*. 2015 Aug 21;10(8)
- Golding JF, Bles W, Bos JE, Haynes T, Gresty MA (2003) Motion sickness and tilts of the inertial force environment: active suspension systems vs. active passengers. *Aviat Space Environ Med* 74: 220-227
- Gravelle, D.C., Laughton, C.A., Dhruv, N.T., Katdare, K.D., Niemi, J.B., Lipsitz, L.A., and Collins, J.J. (2002). Noise-enhanced balance control in older adults. *Neuroreport* 13, 1853-1856.
- Graybiel A (1980) Space motion sickness: Skylab revisited. *Aviat Space Environ Med* 51: 814-822
- Graybiel A, Kellogg RS (1967) Inversion illusion in parabolic flight: its probable dependence on otolith function. *Aerospace Med* 38: 1099-1103
- Graybiel A, Knepton J (1977) Evaluation of a new antinauseant drug for the prevention of motion sickness. *Aviat Space Environ Med* 48: 867-871
- Graybiel A, Lackner JR (1987) Treatment of severe motion sickness with antimotion sickness drug injections. *Aviat Space Environ Med* 58: 773-776
- Graybiel A, Miller EF, Billingham J, Waite R, Berry C, Dietlein LF (1967) Vestibular experiments in Gemini flights V and VII. *Aerospace Med* 38: 360-370
- Graybiel A, Knepton J, Shaw J (1976) Prevention of experimental motion sickness by scopolamine absorbed through the skin. *Aviat Space Environ Med* 47: 1096-1100
- Graybiel A, Miller EF, Homick JL (1977) Experiment M131. Human vestibular function. In: Johnston RS, Dietlein LF (eds) *Biomedical Results from Skylab*. NASA: Washington DC, pp 74-103
- Graybiel A, Cramer DB, Wood CD (1981) Experimental motion sickness: efficacy of transdermal scopolamine plus ephedrine. *Aviat Space Environ Med* 52: 337-339
- Green AM, Angelaki DE (2004). An integrative neural network for detecting inertial motion and head orientation. *J Neurophysiol* 92: 905-925
- Gresty M, Leech J (1977) Coordination of the head and eyes in pursuit of predictable and random target motion. *Aviat Space Environ Med* 48: 741-744
- Griffin MJ (1990) Handbook of Human Vibration. *Academic Press Inc*: San Diego
- Grigoriev AI, Bugrov SA, Bogomolov VV, Egorov AD, Kozlovskaya IB, Pestov ID, Polyakov VV, Tarasov IK (1991) Preliminary medical results of the Mir year-long mission. *Acta Astronautica* 23: 1-8
- Grigoriev AI, Nichiporuk IA, Yasnetsov VV, Shashkov VS (1986) Peculiarities of horizontal status and fluid-electrolyte metabolism during motion sickness. *Proceedings of the 7th IAA Man in Space Symposium: Physiological Adaptation of Man in Space*, Houston, TX, 10-13 Feb 1986
- Grigoryan GA, Gazenko OG, Kozlovskaya IB, Barmin VA, Kreidich YV (1986) The vestibulo-cerebellar regulation of oculomotor reactions in microgravitational conditions. In: Keller EL, Zee DS (eds) *Adaptive Processes in Visual and Oculomotor Systems*. Pergamon Press: New York, pp 111-120
- Grigoryeva LS, Kozlovskaya IB (1987) Effect of weightlessness and hypokinesia on velocity and strength properties of human muscles. *Kosm Biol Aviakosm Med* 21: 27-30
- Groen JJ (1957) The semicircular canal system of the organs of equilibrium. *Physics Med Biol.* 1: 225-242

- Gualtierotti TF, Bracchi F, Rocca E (1972) The orbiting frog otolith experiment. In: *Final report of the Data Reduction and Control Experimentation*. NASA: Houston, TX
- Guedry FE (1964) Visual control of habituation to complex vestibular stimulation in man. *Acta Otolaryngol (Stockh)* 58: 377-389
- Guedry FE (1974) Psychophysics of vestibular sensation. In: Kornhuber HH (ed) *Handbook of Sensory Physiology*. Springer Verlag: Berlin: pp 3-154.
- Guedry FE, Rupert AR, Reschke MF (1998) Motion sickness and development of synergy within the spatial orientation system. A hypothetical unifying concept. *Brain Res Bull* 47: 475-480
- Gurfinkel VS, Lestienne F, Levik Yu S, Popov KE, Lefort L (1993) Egocentric references and human spatial orientation in microgravity II. Body-centered coordinates in the task of drawing ellipses with prescribed orientation. *Exp Brain Res* 95: 343-348
- Gurovskii NN, Eremin AV, Gazenko OG, Egorov AD, Brianov II (1975) Medical investigations during flights of the spaceships 'Soyuz-12', 'Soyuz-13', 'Soyuz-14' and the 'Salyut-3' orbital station. *Kosm Biol Aviakosm Med* 9: 48-54
- Hadfield C (2013) *An Astronaut's Guide to Life*. Macmillan: London
- Hanes DA, McCollum G (2006) Cognitive-vestibular interactions: a review of patient difficulties and possible mechanisms. *J Vestib Res* 95: 343-348
- Harm DL, Parker DE (1993) Perceived self-orientation and self-motion in microgravity, after landing and during preflight adaptation training. *J Vestib Res* 3: 297-305
- Harm DL, Parker DE (1994) Preflight adaptation training for spatial orientation and space motion sickness. *J Clin Pharmacol* 34: 618-627
- Harm DL, Bloomberg JJ, Reschke MF, et al. (1991) Adaptive modification of gaze control following tilt reinterpretation. *Aviat Space Environ Med* 62: 477-481
- Harm DL, Parker DE, Reschke MF (1994) DSO 468. Preflight adaptation trainer. In: *Results of Life Sciences DSOs Conducted Aboard the Space Shuttle, 1991-1993*. NASA: Houston, TX, pp 27-43
- Harm DL, Parker DE, Reschke MF, Skinner NC (1998) Relationship between selected orientation rest frame, circular vection, and space motion sickness. *Brain Res Bull* 47: 497-501
- Harm DL, Reschke MF, Parker DE (1999) Visual-vestibular integration motion perception reporting (DSO 604 OI-1). In: Sawin CF, Taylor GR, Smith WL (eds) *Extended Duration Orbiter Project Final Report 1989-1995*, NASA: Houston, TX, pp 5.2-1 - 5.2-12
- Harm DL, Taylor LC, Bloomberg JJ (2007) Adaptive responses in eye-head-hand coordination following exposures to a virtual environment as a possible space flight analog. *J Gravit Physiol* 14: 83-84
- Harm DL, Reschke MF, Wood SJ (2008) Sensorimotor coordination aftereffects of exposure to a virtual environment. *Vis Comput* 24: 995-999
- Harm DL, Reschke MF, Wood SJ (2015) Spatial orientation and motion perception in microgravity. In: *Cambridge Handbook of Applied Perception Research. Cambridge Handbooks in Psychology*. Cambridge University Press: New York, pp 912-929
- Harris BA Jr, Billica RD, Bishop SL, et al. (1997) Physical examination during space flight. *Mayo Clin Proc* 72: 301-308
- Hayes JL, McBrine JJ, Roper ML, Stricklin MD, Siconolfi SF, Greenisen MC (1992) Effects of Space Shuttle flights on skeletal muscle performance. *FASEB J* 6: A1770
- Herdman SJ (1994) *Vestibular Rehabilitation*. FA Davis Company: Philadelphia

- Hernandez-Korwo R, Kozlovskaia IB, Kreidich IuV, Martinez Fernandez S, Rakhamanov AS (1983) Effect of 9-day space flight on structure and function of human locomotor system. *Kosm Biol Aviakosm Med* 17: 37-44
- Hillman EJ, Bloomberg JJ, McDonald PV, Cohen HS (1999) Dynamic visual acuity while walking in normals and labyrinthine-deficient patients. *J Vestib Res* 9: 49-57
- Hirasaki E, Moore ST, Raphan T, Cohen B (1999) Effects of walking velocity on vertical head and body movements during locomotion. *Exp Brain Res* 127: 117-130
- Hofstetter-Degen K, Weizig J, von Baumgarten R (1993) Oculovestibular interactions under microgravity. *Clin Invest* 10: 749-756
- Holly JE, Wood SJ, McCollum G (2010) Phase-linking and the perceived motion during off-vertical axis rotation. *Biol Cybern* 102: 9-29
- Holmes SR, Bunting A, Brown DL, Hiatt KL, Braithwaite MG, Harrigan MJ (2003) Survey of spatial disorientation in military pilots and navigators. *Aviat Space Environ Med* 74: 957-965
- Holstein GR, Kukielka E, Martinelli GP (1999) Anatomical observations of the rat cerebellar nodulus after 24 hr of space flight. *J Gravit Physiol* 6: 47-50
- Holstein GR, Martinelli GP (2003) The effect of space flight on the ultrastructure of the cerebellum. In: Buckley JC, Homick JL (eds) *The Neurolab Spacelab Mission Neuroscience Research in Space*. NASA: Houston, TX
- Holt KG, Jeng SF, Ratcliffe R, Hamill J (1995) Energetic cost and stability during human walking at the preferred stride frequency. *J Motor Behav* 27: 164-178
- Homick JL (1985) Motion sickness: general backgrounds and methods. In: Kohl RL (ed) *Proceedings of the Space Adaptation Syndrome Drug Workshop*, NASA Space Biomedical Research Institute: Houston, TX, pp 9-19
- Homick JL, Miller EF (1975) Apollo flight crew vestibular assessment. In: Johnston RS, Dietlein LF, Berry CA (eds) *Biomedical Results of Apollo*. NASA: Washington DC, NASA SP-368, pp 323-340
- Homick JL, Reschke MF (1977) Postural equilibrium following exposure to weightless space flight. *Acta Otolaryngol* 83: 45 45-64
- Homick JL, Reschke MF, Miller EF (1977) Effects of prolonged exposure to weightlessness on postural equilibrium. In: Johnston RS, Dietlein LF (eds) *Biomedical Results from Skylab*. NASA: Washington DC, NASA SP-377, pp 104-112
- Homick JL, Kohl RL, Reschke MF, Degioanni J, Cintron-Trevino NM (1983) Transdermal scopolamine in the prevention of motion sickness: evaluation of the time course of efficacy. *Aviat Space Environ Med* 54: 994-1000
- Homick JL, Reschke MF, Vanderploeg JM (1987) Prediction of susceptibility to space motion sickness, in: Graham MD, Kemink JL (eds) *The Vestibular System: Neurophysiologic and Clinical Research*. Raven Press: New York, pp 39-49
- Hopp JJ, Fuchs AF (2006) Amplitude adaptation occurs where a saccade is represented as a vector and not as its components. *Vision Res* 46: 3121-3128
- How J, Lee PS, Seet LC, Tan PK (1988) The Republic of Singapore Navy's Scopoderm TTS study: results after 2,200 man-days at sea. *Aviat Space Environ Med* 59: 646-650
- Howard IP (1982) *Human Visual Orientation*. John Wiley & Sons: Chichester
- Howard IP, Templeton WB (1966) *Human Spatial Orientation*. John Wiley & Sons: London
- Howard IP, Hu G (2001) Visually induced reorientation illusions. *Perception* 30: 583-600

- Hu S, Willoughby LM, Lagomarsino JJ, *et al.* (1996) Optokinetic induced taste aversions correlate with over-all symptoms of motion sickness in humans. *Percept Mot Skills* 82: 859-864
- Hu S, McChesney KA, Player KA, Bahl AM, Buchanan JB, Scozzafava JE (1999) Systematic investigation of physiological correlates of motion sickness induced by viewing an optokinetic rotating drum. *Aviat Space Environ Med* 70: 759-765
- Hudson TE, Lackner JR, DiZio P (2005) Rapid adaptation of torso pointing movements to perturbations of the base of support. *Exp Brain Res* 165: 283-293
- Ito J, Honjo I (1990) Central fiber connections of the vestibulo-autonomic reflex arc in cats. *Acta Otolaryngol* 110: 379-385
- Ivey, C., Apkarian, AV, and Chialvo, DR. (1998). Noise-Induced Tuning Curve Changes in Mechanoreceptors. *J Neurophysiol* 79, 1879–1890
- Jacob RG, Furman JM (2001) Psychiatric consequences of vestibular dysfunction. *Curr Opin Neurol* 14: 41-46
- Jacob RG, Furman JM, Durrant JD, Turner SM (1996) Panic, agoraphobia, and vestibular dysfunction. *Am J Psychiatry* 153: 503-512
- Jaramillo, F., and Wiesenfeld, K. (1998). Mechano-electrical transduction assisted by Brownian motion - a role for noise in the auditory system. *Nature Neuroscience* 1, 384-388
- Jenkin HL, Zacher JE, Jenkin MR, Oman CM, and Harris LR (2007) Effect of field of view on the Levitation Illusion. *J Vestib Res* 2007;17(5-6):271-7
- Jennings RT (1998) Managing space motion sickness. *J Vestib Res* 8: 67-70
- Johnson RS, Dietlein LF, Berry CA (1975) *Biomedical Results of Apollo*. NASA: Washington DC
- Johnson RL, Driscoll TB, LeBlanc AD (1977) Blood volume changes. In: Johnston RS, Dietlein LF (eds) *Biomedical Results from Skylab*. NASA: Washington DC, NASA SP-377, pp 235–241
- Johnson RL, Nicogossian AE, Bergman SA Jr, Hoffler GW (1976) Lower body negative pressure: the second manned Skylab mission. *Aviat Space Environ Med* 47: 347-353
- Jones EM, Glover K (2014) *Apollo Lunar Surface Journal*.  
<http://www.history.nasa.gov/alsj/frame.html>.
- Jones TD (2006) *Sky Walking: An Astronaut's Memoir*. Harper-Collins: New York
- Kalina RE, Mills RP (1980) Acquired hyperopia with choroidal folds. *Ophthalmol* 87: 44-50
- Kanas N, Manzey D (2003) *Space Psychology and Psychiatry*. Microcosm Press: El Segundo, CA
- Karkishchenko NN (1989) Specification for an “ideal” drug to prevent space motion sickness. *Kosm Biol Aviakosm Med* 23: 33-36
- Karni A, Meyer G, Jezard P, Adams MM, Turner R, Ungerleider LG (1995) Functional MRI evidence for adult motor cortex plasticity during motor skill learning. *Nature* 377: 155-158
- Kennedy PM, Cressman EK, Carlsen AN, Chua R (2005) Assessing vestibular contributions during changes in gait trajectory. *Neuroreport* 16: 1097–1100
- Kenyon RV, Young LR (1986) MIT/Canadian vestibular experiments on Spacelab-1 mission: 5. Postural responses following exposure to weightlessness. *Exp Brain Res* 64: 335-346
- Keshner EA (1994) Postural abnormalities in vestibular disorders. In: *Vestibular Rehabilitation*. FA Davis Co: Philadelphia, pp 47-67
- Keshner EA, Cromwell RL, Peterson BW (1995) Mechanisms controlling human head stabilization. II. Head-neck characteristics during random rotations in the vertical plane. *J Neurophysiol* 73: 2302-2312

- Keshner EA, Kenyon RV, Langston J (2004) Postural responses exhibit multisensory dependencies with discordant visual and support surface motion. *J Vestib Res Equilib Orientat* 14: 307–319
- King WM, Zhou W, Tang B, Newlands S (1999) Neural processing of otolith signals in macaque vestibular nuclei [abstract]. In: *Vestibular Influences on Spatial Orientation. A Satellite of the Society for the Neural Control of Movement*. Ninth Annual Meeting. Princeville: Hawaii
- Knierim JJ, McNaughton BL, Poe GR (2000) Three-dimensional spatial selectivity of hippocampal neurons during saccade flight. *Nature Neurosci* 3: 209-210
- Koch KL, Xu L, Bingaman S, et al. (1994) Effect of ondansetron on motion sickness, gastric dysrhythmias, and plasma vasopressin. *Gastroenterology* 106: A525
- Kolev O, Reschke MF (2014) Internally versus externally mediated triggers in the acquisition of visual targets in the horizontal plane. *Behav Brain Res* 266: 131-136
- Koppelmans V, Erdeniz B, De Dios YE, Wood SJ, Reuter-Lorenz PA, Kofman I, Bloomberg JJ, Mulavara AP, Seidler RD (2013) Study protocol to examine the effects of spaceflight and a spaceflight analog on neurocognitive performance: Extent, longevity, and neural bases. *BMC Neurology* 13: 205. Doi: 10.1186/1471-2377-13-20
- Kornilova LN (1997) Vestibular function and sensory interaction in altered gravity. *Adv Space Biol Med* 6: 275-313
- Kornilova LN, Kaspransky RR (1994) Vestibular function and sensory interaction after space flight. *Proceedings of the Fifth European Symposium on Life Sciences Research in Space*, Arcachon, France 26 Sept-1 Oct 1993, ESA SP-366, pp 345-350
- Kornilova LN, Goncharenko AM, Bodó G, Elkán K, Grigorova V, Manev A (1991a) Pathogenesis of sensory disorders in microgravity. *Physiologist* 34: S36-S39
- Kornilova LN, Goncharenko AM, Korsunskii SB, Tarasov IK, Alekseev VN (1991b) Vestibular function and interanalyzer interaction following space flight. *Kosmiches. Biol Aviakosmiches Med* 25: 12-17
- Kornilova LN, Yakovleva IY, Tarasov IK, Gorgiladze GI (1983) Vestibular dysfunction in cosmonauts during adaptation to zero-g and readaptation to one-g. *Physiologist* 26: S35-S40
- Kornilova LN, Muller KH, Chernobylskiy LM (1995) Phenomenology of illusory reactions in weightlessness. *Fiziologiya Cheloveka* 21: 50-62
- Kornilova LN, Naumov IA, Makarova SM (2011) Static Torsional Otolith-Cervical-Ocular Reflex After Prolonged Exposure to Weightlessness and a 7-day Immersion. *Acta Astronautica* 68: 1462-1468
- Kozlovskaya IB, Kreidich YuV, Oganov VS, Koserenko OP (1981) Pathophysiology of motor functions in prolonged manned space flights. *Acta Astronaut* 8: 1059–1072, 1981.
- Kozlovskaya IB, Babaev BM, Barmin VA, Beloozerova II, Kreidich YuV, Sirota MG (1984) The effect of weightlessness on motor and vestibulo-motor reactions. *Physiologist* 27: S111-S114
- Kozlovskaya IB, Barmin VA, Kreidich Y, Repin AA (1985) The effects of real and simulated microgravity on vestibulo-oculomotor interaction. *Physiologist* 28: S51-56
- Kozlovskaya I, Sayenko I, Vinogradova O, Miller T, Khusnutdinova D, Melnik K, Yarmanova E (2006) New approaches to countermeasures of the negative effects of microgravity in long-term space flights. *Acta Astronaut* 59: 13–19, 2006.

- Kozlovskaya IB, Sayenko IV, Sayenko DG, Miller TF, Khusnutdinova DR, Melnik KA (2007) Role of support afferentation in control of the tonic muscle activity. *Acta Astronaut* 60: 285–294, 2007.
- Krasnov IB (1994) Gravitational neuromorphology. *Adv Space Biol Med* 4: 85-110
- Krakauer JW (2006) Motor learning: its relevance to stroke recovery and neurorehabilitation. *Curr Opin Neurol*. 2006 Feb;19(1):84-90.
- Kubis JF, Elrod JT, Rusnak R, Barnes JE (1972a) Apollo 15 Time and Motion Study. Final Report M72-4. NASA Manned Spacecraft Center, Houston Texas
- Kubis JF, Elrod JT, Rusnak R, Barnes JE, Saxon SC (1972b) Apollo 16 Time and Motion Study. Final Mission Report M72-6. NASA Manned Spacecraft Center, Houston Texas
- Kubis J, McLaughlin E, Jackson J, Rusnak D, McBride G, Saxon S (1977) Task and work performance on Skylab missions 2, 3, and 4: Time and motion study - Experiment M151. In: Johnston RS, Dietlein LF, Berry CA (eds) Biomedical Results from Skylab. NASA: Washington DC, pp 136-154
- Kurtzer I, DiZio PA, Lackner JR (2005) Adaptation to a novel multi-force environment. *Exp Brain Res* 164: 120-132
- Kyparos A, Feedback DL, Layne CS, Martinez DA, Clarke MSF (1985) Mechanical stimulation of the plantar foot surface attenuates soleus muscle atrophy induced by hindlimb unloading in rats. *J Appl Physiol Bethesda Md* 1985 99: 739–746, 2005.
- Lackner JR, DiZio P (1992) Gravitoinertial-force level affects the appreciation of limb position during muscle vibration. *Brain Res* 592, 175-180
- Lackner JR, DiZio P (1993) Multisensory cognitive, and motor influences on humans spatial orientation in weightlessness. *J Vestib Res* 3: 361-372
- Lackner JR DiZio P (2003) Adaptation to rotating artificial gravity environments. *J Vestib Res* 13: 321-330
- Lackner JR, DiZio P (2005a) Motor control and learning in altered dynamic environments. *Curr Opin Neurobiol* 15: 653-659
- Lackner JR, DiZio P (2005b) Vestibular, proprioceptive, and haptic contributions to spatial orientation. *Ann Rev Psychol* 56: 115-147
- Lackner JR, DiZio P (2006) Space motion sickness. *Exp Brain Res* 175: 377-399
- Lackner JR, DiZio P (2000) Human orientation and movement control in weightless and artificial gravity environments. *Exp Brain Res*. 2000 Jan;130(1):2-26.
- Laitinen LA, Tokola O, Gothoni G, Vapaatalo H (1981) Scopolamine alone or combined with ephedrine in seasickness: a double-blind, placebo-controlled study. *Aviat Space Environ Med* 52: 6-10
- Lapayev EV, Vorobyev OA (1986) The problem of vestibular physiology in aerospace medicine and prospects for its solution. In: Gzenko OG (ed) *Eighth All-Union Conference on Space Biology and Aerospace Medicine*. Kaluga, USSR, 25-27 June 1986. Nauka: Moscow, pp 85-86
- Lathan CE, Tracey M (2002) The effects of operator spatial perception and sensory feedback on human-robot teleoperation performance. *Presence: Teleop Virtual Environ* 11: 368-377
- Lathan CE, Wang Z, Clément G (2000) Changes in the vertical size of a three-dimensional object drawn in weightlessness by astronauts. *Neurosci Lett* 295: 37-40
- Layne CS, Forth KE (2008) Plantar stimulation as a possible countermeasure to microgravity-induced neuromotor degradation. *Aviat Space Environ Med* 79: 787-794

- Layne CS, McDonald PV, Bloomberg JJ (1997) Neuromuscular activation patterns during locomotion after space flight. *Exp Brain Res* 113: 104-116
- Layne, C.S., Mulavara, A.P., Pruett, C.J., McDonald, P.V., Kozlovskaya, I.B., Bloomberg, J.J (1998a) The use of inflight foot pressure as a countermeasure to neuromuscular degradation. *Acta Astronautica*, 42: 231-246, 1998
- Layne CS, Lange GW, Pruett CJ, *et al.* (1998b) Adaptation of neuromuscular activation patterns during locomotion after long-duration space flight. *Acta Astronautica* 43: 107-119
- Layne CS, Mulavara A, McDonald PV, Pruett CJ, Kozlovskaya IB, Bloomberg JJ (2004) Alterations in human neuromuscular activation during overground locomotion after long-duration spaceflight. *J Gravit Physiol* 11: 1-15
- Lee MH, Durnford SJ, Crowley JS, Rupert AH (1997) Visual vestibular interaction in the dynamic visual acuity test during voluntary head rotation. *Aviat Space Environ Med* 68: 111-117
- Léone G, Lipshits M, Gurfinkel V, Berthoz A (1995) Is there an effect of weightlessness on mental rotation of three-dimensional objects? *Cogn Brain Res* 2: 255-267
- Leonov AA, Lebedev VI (1968) *Perception of Space and Time in Outer Space*. Nauka Press: Moskow. NASA Technical Translation, NASA: Washington DC, NASA TT F-545, 1969
- Levy GD, Rapaport MH (1985) Transdermal scopolamine efficacy related to time of application prior to the onset of motion. *Aviat Space Environ Med* 56: 591-593
- Liao K, Schneider RM, Yaniglos SS, Bertolini G, Glendinning P, Sawyer Jr RN, Reschke MF, Leigh RJ (2011) Visual and vestibular determinants of the translational vestibulo-ocular reflex. *Ann NY Acad Sci* 1233: 263-270
- Linenger JM (2000) *Off the Planet: Surviving Five Perilous Months Aboard the Space Station Mir*. McGraw-Hill: New York
- Lipshits MI, Gurfinkel EV, Matsakis F, Lestienne F (1993) Microgravity effects on sensorimotor interaction during operational activity: visual feedback. Latent period of motor response. *Aerospace Environ Med* 27: 22-25
- Lowrey CR, Perry SD, Strzalkowski NDJ, Williams DR, Wood SJ, Bent LR. Selective skin sensitivity changes and sensory reweighting following short-duration space flight. *J Appl Physiol* Bethesda Md 1985 116: 683–692, 2014.
- Lu E (2003) Ed Lu's Journal Entry #8, Which way is up? SpaceRef. Available at: <http://www.spaceref.com/news/viewsr.html?pid=10441>
- Lyons TJ, Ercoline WR, Freeman JE, Gillingham KK (1994) Classification problems of U.S. Air Force spatial disorientation accidents, 1989-91. *Aviat Space Environ Med* 65: 147-152
- MacDougall HG, Moore ST (2005a) Functional assessment of head-eye coordination during vehicle operation. *Optom Vis Sci* 82: 706-715
- MacDougall HG, Moore ST (2005b) Marching to the beat of the same drummer: the spontaneous tempo of human locomotion. *J Appl Physiol* 99: 1164-1173
- Madansingh S, Bloomberg JJ (2015) Understanding the effects of spaceflight on head–trunk coordination during walking and obstacle avoidance. *Acta Astronaut* 115: 165-172
- Mader TH, Gibson CR, Pass AF, *et al.* (2011) Optic disc edema, globe flattening, choroidal folds, and hyperopic shifts observed in astronauts after long-duration space flight. *Ophthalmol* 118: 2058-2069

- Mader TH, Gibson CR, Pass AF, Lee AG, Killer HE, Hansen HC, Dervay JP, Barratt MR, Tarver WJ, Sargsyan AE, Kramer LA, Riascos R, Bedi DG, Pettit DR (2013) Optic disc edema in an astronaut after repeat long-duration space flight. *J Neuro Ophthalmol* 33:249-255
- Magalhaes, F.H., and Kohn, A.F. (2011). Vibratory noise to the fingertip enhances balance improvement associated with light touch. *Exp Brain Res* 209, 139-151. doi: 10.1007/s00221-010-2529-3
- Magalhaes, F.H., and Kohn, A.F. (2012). Imperceptible electrical noise attenuates isometric plantar flexion force fluctuations with correlated reductions in postural sway. *Exp Brain Res* 217, 175-186. doi: 10.1007/s00221-011-2983-6
- Malcik V (1968) Performance decrement in a flight simulator due to galvanic stimulation of the vestibular organ and its validity for success in flight training. *Aerospace Med* 39: 941-943
- Manzey D, Lorenz B (1998) Mental performance during short-term and long-term space flight. *Brain Res Rev* 28: 215-221
- Manzey D, Lorenz B, Schiewe A, Finell G, Thiele G (1995) Dual-task performance in space: results from a single-case study during a short-term space mission. *Human Factors* 37: 667-681
- Manzey D, Lorenz B, Polyakov VV (1998) Mental performance in extreme environments: Results from a performance monitoring study during a 438-day space flight. *Ergonomics* 41: 537-559
- Massion J, Popov K, Fabrel JC, *et al.* (1997) Is the erect posture in microgravity based on the control of trunk orientation or center of mass position? *Exp Brain Res* 114: 384-389
- Masterton BA, Biederman GB (1983) Proprioceptive versus visual control in autistic children. *J Autism Dev Disord* 13: 141-152
- Mayne R (1974) A systems concept of the vestibular organs. In: Kornhuber HH (ed) *Handbook of Sensory Physiology*. Springer Verlag: Berlin, pp 493-580
- McCauley ME, Royal JW, Shaw JE, Schmitt LG (1979) Effect of transdermally administered scopolamine in preventing motion sickness. *Aviat Space Environ Med* 50: 1108-1111
- McCluskey R, Clark J, Stepaniak P (2001) Correlation of Space Shuttle landing performance with cardiovascular and neurovestibular dysfunction resulting from space flight. In: *Human Systems 2001*, Clear Lake City, TX
- McDonald PV, Bloomberg JJ, Layne CS (1997) A review of adaptive change in musculoskeletal impedance during space flight and associated implications for post-flight head movement control. *J Vestib Res* 7: 239-250
- McDonnell MD, Abbott D. What is stochastic resonance? Definitions, misconceptions, debates, and its relevance to biology. *PLoS computational biology*. 2009;5(5):e1000348. doi: 10.1371/journal.pcbi.1000348.
- McRuer DT (1972) The development of pilot-in-the-loop analysis. *J Aircraft* 10: 515-524
- Meddick RDL, Griffin MJ (1976) The effect of two-axis vibration on the legibility of reading material. *Ergonomics* 19: 21-33
- Melvill-Jones, G (1974) Adaptive neurobiology in space flight. In: *Skylab Life Sciences Symposium*, Houston, TX
- Menchaca-Brandan MA, Liu AM, Oman C, Natapoff A (2007) Influence of perspective-taking and mental rotation abilities in space teleoperation. In: *ACM/IEEE International Conference on Human-Robot Interaction*. Washington, DC.

- Merfeld DM (1996a) Effect of space flight on ability to sense and control roll tilt: human neurovestibular studies on SLS-2. *J Appl Physiol* 81: 50-57
- Merfeld DM (2003) Rotation otolith tilt-translation reinterpretation (ROTTR) hypothesis: a new hypothesis to explain neurovestibular space flight adaptation. *J Vestib Res* 13: 309-320
- Merfeld DM, Polutchko KA, Schultz K (1996b) Perceptual responses to linear acceleration after space flight: human neurovestibular studies on SLS-2. *J Appl Physiol* 81: 58-68
- Merfeld DM, Young LR, Oman CM, Shelhamer MJ (1993) A multidimensional model of the effect of gravity on the spatial orientation of the monkey. *J Vestib Res* 3: 141-161
- Merfeld DM, Zupan LH (2002) Neural processing of gravito-inertial cues in humans. III. Modeling tilt and translation responses. *J Neurophysiol* 87: 819-833
- Merfeld DM, Park S, Gianna-Poulin C, Black FO, Wood S (2005a) Vestibular perception and action employ qualitatively different mechanisms. I. Frequency response of VOR and perceptual responses during translation and tilt. *J Neurophysiol* 94: 186-198
- Merfeld DM, Park S, Gianna-Poulin C, Black FO, Wood S (2005b) Vestibular perception and action employ qualitatively different mechanisms. II. VOR and perceptual responses during combined tilt and translation. *J Neurophysiol* 94: 199-205
- Michaud L, Parker DE, Harm DL (1989) Changes in the vestibulo-ocular reflex associated with simulated stimulus conditions of space flight. *Soc Neurosci Abstr* 15: 514
- Miller CA, Peters BT, Brady RR, Richards JR, Ploutz-Snyder RJ, Mulavara AP, Bloomberg JJ (2010) Changes in toe clearance during treadmill walking after long-duration space flight. *Aviat Space Environ Med* 81: 919-928
- Mindell DA (2008) Digital Apollo: Human, Machine and Space Flight. MIT Press: Cambridge, MA
- Mittelstaedt H (1986a) In search of the cause of the 'inversion illusion.' In: *Physiological Adaptation of Man in Space: 7th International "Man in Space" Symposium*, Houston, TX
- Mittelstaedt H (1986b) In-flight and postflight results on the cessation of inversion illusions and space sickness. *Proceedings of the Symposium on Scientific Results of the German Spacelab Mission D-1*, Norderney, Germany
- Mittelstaedt H (1986c) The subjective vertical as a function of visual and extraretinal cues. *Acta Psychol*. 63: 63-85
- Mittelstaedt H (1999) The role of the otoliths in perception of the vertical and in path integration. *Ann. NY Acad. Sci.* 871: 334-344
- Mittelstaedt H (1988) Determinants of space perception in space flight. *Adv Otorhinolaryngol* 42: 18-23
- Mittelstaedt H (1992) Somatic versus vestibular-gravity reception in man. *Annals NY Acad Sci* 656: 124-139
- Mittelstaedt H, Glasauer S (1993) Illusions of verticality in weightlessness. *Clin Investig* 71: 732-739
- Mittelstaedt ML, Mittelstaedt H (1996) The influence of otoliths and somatic graviceptors on angular velocity estimation. *J Vestib Res* 6: 355-366
- Money KE (1970) Motion sickness. *Physiol Rev* 50: 1-39
- Money KE, Lackner J, Cheung R (1996) The autonomic nervous system and space motion sickness. In: Yates BJ, Miller AD (eds) *Vestibular Autonomic Regulation*. CRC Press: Boca Raton, pp 147-173

- Moore ST, Clément G, Raphan T, Cohen B (2001) Ocular counterrolling induced by centrifugation during orbital space flight. *Exp Brain Res* 137: 323-335
- Moore ST, Cohen B, Raphan T, Berthoz A, Clément G (2005a) Spatial orientation of optokinetic nystagmus and ocular pursuit during orbital space flight. *Exp Brain Res* 160: 38-59
- Moore ST, Hirasaki E, Raphan T, Cohen B (2005b) Instantaneous rotation axes during active head movements. *J Vestib Res* 15: 73-80
- Moore ST, MacDougall, Peters BT, Bloomberg JJ, Curthoys IS, Cohen HS (2006) Modeling locomotor dysfunction following space flight with galvanic vestibular stimulation. *Exp Brain Res* 174: 647-659
- Moore ST, MacDougall HG, Gracies JM, Cohen HS, Ondo WG (2007) Long-term monitoring of gait in Parkinson's disease. *J Gait Posture* 26: 200-207
- Moore ST, MacDougall HG, Lescue X, Speyer JJ, Wuyts F, Clark JB (2008a) Head-eye coordination during simulated orbiter landings. *Aviat Space Environ Med* 79: 888-898
- Moore ST, Dilda V, MacDougall HG (2011) Galvanic vestibular stimulation as an analog of spatial disorientation after spaceflight. *Aviat Space Environ Med* 82: 535-542
- Moore ST, Dilda V, Morris T, Yungher D, MacDougall H (2015) Pre-adaptation to noisy galvanic vestibular stimulation is associated with enhanced sensorimotor performance in novel vestibular environments. *Front Syst Neurosci* 8; 9. Doi 10.3389/fnsys.2015.00088
- Morse, R.P., and Evans, E.F. (1996). Enhancement of vowel coding for cochlear implants by addition of noise. *Nature Medicine* 2, 928-932
- Moseley MJ, Griffin MJ (1986) Effects of display vibration and whole-body vibration on visual performance. *Ergonomics* 29: 977-983
- Moss F, Ward LM, Sannita WG. Stochastic resonance and sensory information processing: a tutorial and review of application. *Clinical Neurophysiology*. 2004;115(2):267-81. doi: 10.1016/j.clinph.2003.09.014
- Mueller C, Kornilova L, Wiest G, Steinhoff N, Deecke L (1993) Results from vertical vection experiments in short and long-term space flights. *Proceedings of the 5th European Symposium on Life Sciences Research in Space*, Arcachon, France
- Muir J, Judex S, Qin Y-X, Rubin C (2011) Postural instability caused by extended bed rest is alleviated by brief daily exposure to low magnitude mechanical signals. *J Gait Posture* 33: 429-435
- Mulavara AP, Bloomberg JJ (2003) Identifying head-trunk and lower limb contributions to gaze stabilization during locomotion. *J Vestib Res* 12: 255-269
- Mulavara AP, Houser J, Miller C, Bloomberg JJ (2005a) Full-body gaze control mechanisms elicited during locomotion: effects of VOR adaptation. *J Vestib Res* 15: 279-289
- Mulavara AP, Richards JT, Ruttley T, Marshburn A, Nomura Y, Bloomberg JJ (2005b) Exposure to a rotating virtual environment during treadmill locomotion causes adaptation in heading direction. *Exp Brain Res* 166: 210-219
- Mulavara AP, Cohen HS, Bloomberg JJ (2009) Critical features of training that facilitates adaptive generalization of over ground locomotion. *J Gait Posture* 29: 242-248
- Mulavara AP, Feiveson A, Feidler J, Cohen HS, Peters BT, Miller CA, Brady R, Bloomberg JJ (2010) Locomotor function after long-duration space flight: Effects and motor learning during recovery. *Exp Brain Res* 202: 649-659

- Mulavara AP, Fiedler MJ, Kofman IS, Wood SJ, Serrador JM, Peters B, Bloomberg JJ (2011) Improving balance function using vestibular stochastic resonance: optimizing stimulus characteristics. *Exp Brain Res* 210: 303-312
- Mulavara AP, Ruttley T, Cohen HS, Peters BT, Miller C, Brady R, Merkle L, Bloomberg JJ (2012) Vestibular-somatosensory convergence in head movement control during locomotion after long-duration space flight. *J Vestib Res* 22: 153-166
- Mulavara, AP Cohen HS, Peters BT, Sangi-Haghpeykar H, Bloomberg, JJ (2013) New analyses of the sensory organization test compared to the clinical test of sensory integration and balance. *Laryngoscope* 123: 2276-2280
- Mulavara AP, Kofman IS, De Dios YE, Miller C, Peters BT, Goel R, Galvan-Garza R, Bloomberg JJ (2015) Using low levels of stochastic vestibular stimulation to improve locomotor stability. *Front Syst Neurosci* 2015 Aug 24;9:117
- Nachum Z, Shupak A, Letichevsky V, Ben-David J, Tal D, Tamir A, Talmon Y, Gordon CR, Luntz M (2004) Mal de débarquement and posture: reduced reliance on vestibular and visual cues. *The Laryngoscope* 114: 581–586
- Nelson GA (2009) Neurological effects of space radiation. *Gravit Space Biol* 22: 33-38
- Newberg A (1994) Changes in the central nervous system and their clinical correlates during long-term space flight. *Aviat Space Environ Med* 65: 562-572
- Newman DJ, Jackson DK, Bloomberg JJ (1997) Altered astronaut lower-limb and mass center kinematics in downward jumping following space flight. *Exp Brain Res* 117: 30-42
- Nicogossian AW, Leach-Huntoon C, Pool SL (1989) *Space Physiology and Medicine*. Lea & Febiger: Philadelphia
- Nomura Y, Mulavara AP, Richards JT, Brady R, Bloomberg JJ (2005) Optic flow dominates visual scene polarity in causing adaptive modification of locomotor trajectory. *Cogn Brain Res* 3: 624-631
- Normand H, Etard O, Denise P (1997) Otolithic and tonic neck receptors control of limb blood flow in humans. *J Appl Physiol* 82: 1734-1738
- Noy S, Shapira S, Zilbiger A, Ribak J (1984) Transdermal therapeutic system scopolamine (TTSS), dimenhydrinate, and placebo - a comparative study at sea. *Aviat Space Environ Med* 55: 1051-1054
- Offenloch K, Zahner G, Dietlein G, Franz I (1986) Comparative in-flight study of a scopolamine-containing membrane plaster versus dimenhydrinate under defined acceleration conditions. *Arzneimittelforschung* 36: 1401-1406
- Olesen PJ, Westerberg H, Klingberg T (2004) Increased prefrontal and parietal activity after training of working memory. *Nat Neurosci* 7: 75-79
- Oman CM (1982a) A heuristic mathematical model for the dynamics of sensory conflict and motion sickness. *Acta Otolaryngol* 392: 1-44
- Oman CM (1982b) Space motion sickness and vestibular experiments in Spacelab. *Proceedings of the 12th AIAA, SAE, ASME, AChE, and ASMA, Intersociety Conference on Environmental Systems*, San Diego, 19-21 July 1982
- Oman, CM (1988) The role of static visual-orientation cues in the etiology of space motion sickness. *Proceedings of the Symposium on Vestibular Organs and Altered Environments*, Houston, TX

- Oman CM (1998) Sensory conflict theory and space sickness: Our changing perspective. *J Vestib Res* 8: 51-56
- Oman CM (2003) Human Visual Orientation in Weightlessness. In: Jenkin LH (ed) *Levels of Perception*. Springer Verlag: New York, pp 375-398
- Oman CM (2007) Spatial Orientation and Navigation in Microgravity. In: Janeke FW (ed) *Spatial Processing in Navigation, Imagery and Perception*. Springer Verlag: New York
- Oman CM, Kulbaski MJ (1988) Space flight affects the 1-g post-rotatory vestibulo-ocular reflex. *Adv Oto Rhino Laryngol* 42: 5-8
- Oman CM, Balkwill MD (1993) Horizontal angular VOR, nystagmus dumping, and sensation duration in Spacelab SLS-1 crewmembers. *J Vestib Res* 3: 315-330
- Oman CM, Lichtenberg BK, Money KE, McCoy RK (1986) M.I.T./Canadian vestibular experiments on the Spacelab-1 mission: 4. Space motion sickness: symptoms, stimuli, and predictability. *Exp. Brain Res.* 64: 316-334
- Oman CM, Marcus EN, Curthoys IS (1987) The influence of semicircular canal morphology on endolymph flow dynamics. An anatomically descriptive mathematical model. *Acta Otolaryngol* 103: 1-13
- Oman CM, Lichtenberg BK, Money KE (1990) Space motion sickness monitoring experiment: Spacelab 1. In: Crampton GH (ed) *Motion and Space Sickness*. CRC Press: Boca Raton, FL, pp 217-246
- Oman CM, Pouliot CF, Natapoff A (1996) Horizontal angular VOR changes in orbital and parabolic flight: human neurovestibular studies on SLS-2. *J Appl Physiol* 81: 69-81
- Oman CM, Benveniste D, Buckland DA, Aoki H, Liu A, Natapoff A (2006) Spacecraft module visual verticals and training affect spatial task performance. *Habitation* 10: 202-203
- Onai T, Takayama K, Miura M (1987) Projections to areas of the nucleus tractus solitarii related to circulatory and respiratory responses in cats. *J Auton Nerv Syst* 18: 163-175
- Ortega HJ, Harm DL (2008) Space and entry motion sickness. In: Barratt M, Pool SL (eds) *Principles of Clinical Medicine for Space Flight*. Springer: New York, chapter 10, pp 211-222.
- Page NG, Gresty MA (1985) Motorist's vestibular disorientation syndrome. *J Neurol Neurosurg Psychiatry* 48: 729-735
- Paige GD, Barnes GR, Telford L, Seidman SH (1996) Influence of sensorimotor context on the linear vestibulo-ocular reflex. *Ann NY Acad Sci* 781: 322-331
- Paige GD, Seidman SH (1999) Characteristics of the VOR in response to linear acceleration. *Ann NY Acad Sci* 871: 123-135
- Paloski WH, Harm DL, Reschke MF, *et al.* (1990) Postural changes following sensory reinterpretation as an analog to space flight. *Proceedings of the Fourth European Symposium on Life Sciences Research in Space*, Paris, France, 28 May-1 June 1990. ESA SP-307, pp 175-178
- Paloski WH, Black FO, Reschke MF, Calkins DS, Shupert C (1993) Vestibular ataxia following shuttle flights: effects of microgravity on otolith-mediated sensorimotor control of posture. *Am J Otolaryngol* 14: 9-17
- Paloski WH, Bloomberg JJ, Reschke MF, Harm DL (1994) Space flight-induced changes in posture and locomotion. *J Biomech* 27: 812
- Paloski WH, Oman CM, Bloomberg JJ, Reschke MF, Wood SJ, Harm DH, Peters BT, Mulavara AP, Locke JP, Stone LS (2008) Risk of sensory-motor performance failures during exploration-

- class space missions: A review of the evidence and recommendations for future research. *J Gravit Physiol* 15: 1-29
- Panait L, Merrell RC, Rafiq A, Dudrick S, Broderick T (2006) Virtual reality laparoscopic skill assessment in microgravity. *J Surg Res* 136: 198-203
- Papaxanthis C, Pozzo T, Popov KE, McIntyre J (1998) Hand trajectories of vertical arm movements in one-G and zero-G environments. *Exp Brain Res* 120: 496-502
- Parihar VK, Pasha J, Tran KK, Craver BM, Acharya MM, Limoli CL (2014) Persistent changes in neuronal structure and synaptic plasticity caused by proton irradiation. *Brain Struct Funct* 1-11
- Park S, Gianna-Poulin C, Black FO, Wood S, Merfeld DM (2006) Roll rotation cues influence roll tilt perception assessed using a somatosensory technique. *J Neurophysiol* 96: 486-491
- Parker DE, Parker KL (1990) Adaptation to the simulated stimulus rearrangement of weightlessness. In: Crampton GH (ed) *Motion and Space Sickness*. CRC Press: Boca Raton, FL, pp 247-262
- Parker DE, Harm DL (1992) Mental rotation: A key to mitigation of motion sickness in the virtual environment? *Presence* 1: 329-333
- Parker DE, Tjernstrom O, Ivarsson A, Gullledge WL, Poston RL (1983) Physiological and behavioral effects of tilt-induced body fluid shifts. *Aviat Space Environ Med* 54: 402-409
- Parker DE, Reschke MF, Arrott AP, Homick JL, and Lichtenberg BK (1985) Otolith tilt-translation reinterpretation following prolonged weightlessness: implications for preflight training. *Aviat Space Environ Med* 56: 601-606
- Parker DE, Reschke MF, Ouyang L, Arrott AP, Lichtenberg BK (1986) Vestibulo-ocular reflex changes following weightlessness and preflight adaptation training. In: Keller EL, Zee DS (eds) *Adaptive Processes in Visual and Oculomotor Systems*. Pergamon Press: Oxford, pp 103-109
- Persterer A, Berger M, Koppensteiner C, et al. (1992) AudiMir–Directional hearing at microgravity. In: Society for Aerospace Medicine (ed) *Health from Space Research. Austrian Accomplishments*. Springer-Verlag: Wien, pp 21-38
- Peters BT, Bloomberg JJ (2005) Dynamic visual acuity using FAR and NEAR targets. *Acta Otolaryngol* 125: 353-357
- Peters BT, van Emmerik RA, Bloomberg JJ (2006) Stride cycle influences on goal-directed head movements made during walking. *Gait Posture* 24: 70-76
- Peters BT, Miller CA, Richards JT, Brady RA, Mulavara AP, Bloomberg JJ (2011) Dynamic visual acuity during walking after long-duration space flight. *Aviat Space Environ Med* 82: 463-466
- Peters BT, Brady RA, Bloomberg JJ (2012a) Walking on an oscillating treadmill: Strategies of stride-time adaptation. *Ecological Psychology* 24: 265-278
- Peters BT, Mulavara AP, Cohen HS, Sangi-Haghpeykar H, Bloomberg JJ (2012b) Dynamic visual acuity testing for screening patients with vestibular impairments. *J Vestib Res* 22: 145-151
- Peters BT, Cohen HS, Sangi-Haghpeykar H, Bloomberg JJ (2013a) Effects of distance and duration on vertical dynamic visual acuity in screening healthy adults and people with vestibular disorders. *J Vestib Res* 23: 285-291
- Peters BT, Brady RA, Batson CD, Guined JR, Ploutz-Snyder RJ, Mulavara AP, Bloomberg JJ (2013b) Changes in locomotor stability, cognition, and metabolism during adaptation to walking in discordant sensory conditions. *Aviat Space Environ Med* 84: 567-572

- Peterson BW, Chen-Huang C (2002) Otolith neurons distinguish between translations and tilts. *J Vestib Res* 11: 255
- Pettit D (2010) Mars landing on Earth: An astronaut's perspective. *J Cosmology* 12: 3529-3536
- Piana, M., Canfora, M., and Riani, M. (2000). Role of noise in image processing by the human perceptive system. *Phys Rev E Stat Phys Plasmas Fluids Relat Interdiscip Topics* 62, 1104-1109.
- Pompeiano O (2003) Gene expression in the rat brain during space flight. In: Buckley JC Homick JL (eds) *The Neurolab Space Mission Neuroscience Research in Space*. NASA: Houston, TX
- Poon C, Merfeld D (2005) Internal models: the state of the art. *J Neural Eng* 2: S144-S148
- Pozzo T, Berthoz A, Lefort L (1990) Head stabilization during various locomotor tasks in humans. I. Normal subjects. *Exp Brain Res* 82: 97-106
- Pozzo T, Levik Y, Berthoz A (1995) Head and trunk movements in the frontal plane during complex dynamic equilibrium tasks in humans. *Exp Brain Res* 106: 327-338
- Previc FH (1993) Do the organs of the labyrinth differentially influence the sympathetic and parasympathetic system? *Neurosci Biobehav Rev* 17: 397-404
- Previc FH, Ercoline WR (2004) Spatial disorientation in aviation. In: *Progress in Aeronautics and Astronautics*. AIAA: Reston, VA
- Previc FH, Yauch DW, DeVilbiss CA, Ercoline WR, Sipes WE (1995) Defense of traditional views of spatial disorientation and loss of situation awareness. *Aviat Space Environ Med* 66: 1103-1106
- Priplata, A., Niemi, J., Salen, M., Harry, J., Lipsitz, L.A., and Collins, J.J. (2002). Noise-enhanced human balance control. *Phys Rev Lett* 89, 238101
- Priplata, A.A., Niemi, J.B., Harry, J.D., Lipsitz, L.A., and Collins, J.J. (2003). Vibrating insoles and balance control in elderly people. *Lancet* 362, 1123-1124. doi: 10.1016/S0140-6736(03)14470-4
- Priplata, A.A., Prittiti, B.L., Niemi, J.B., Hughes, R., Gravelle, D.C., Lipsitz, L.A., Veves, A., Stein, J., Bonato, P., and Collins, J.J. (2006). Noise-enhanced balance control in patients with diabetes and patients with stroke. *Ann Neurol* 59, 4-12. doi: 10.1002/ana.20670
- Pyykkö I, Padoan S, Schalen L, Lyttkens L, Magnusson M, Henriksson NG (1985) The effects of TTS-scopolamine, dimenhydrinate, lidocaine, and tocainide on motion sickness, vertigo, and nystagmus. *Aviat Space Environ Med* 56: 777-782
- Rabin E, DiZio P, Lackner JR (2006) Time course of haptic stabilization of posture. *Exp Brain Res* 170: 122-126
- Rafiq A, Hummel R, Lavrentyev V, Derry W, Williams D, Merrell RC (2006) Microgravity effects on fine motor skills: tying surgical knots during parabolic flight. *Aviat Space Environ Med* 77: 852-856
- Ramachandran VS (1988) Perceiving shape from shading. *Sci Am* 259: 76-83, 1988
- Ratino D, Repperger D, Goodyear C, Potor G, Rodriguez L (1988) Quantification of reaction time and time perception during Space Shuttle operations. *Aviat Space Environ Med* 59: 220-224
- Ray CA, Hume KM, Shortt TL (1997) Skin sympathetic outflow during head-down neck flexion in humans. *Am J Physiol* 273: R1142-R1146
- Reason JT (1978) Motion sickness adaptation: a neural mismatch model. *J Roy Soc Med* 71: 819-829
- Reason JT, Brand JJ (1975) *Motion Sickness*. Academic Press: London

- Reinertson RC, Nelson VA, Aunon SM, Schlegel TT, Lindgren KN, Kerstman E (2007) Medical monitoring during the NASA artificial gravity-bed rest pilort study. *J Gravity Physiol* 14: P9-P13
- Reschke MF, Anderson DJ, Homick JL (1984) Vestibulospinal reflexes as a function of microgravity. *Science* 225: 212-214
- Reschke MF, Anderson DJ, Homick JL (1986) Vestibulo-spinal response modification as determined with the Hoffman reflex during the Spacelab-1 flight. *Exp Brain Res* 64: 367-379
- Reschke MF, Parker DE (1987) Effects of prolonged weightlessness on self-motion perception and eye movements evoked by roll and pitch. *Aviat Space Environ Med* 58: A153-158
- Reschke MF, Parker DE, Harm DL, Michaud L (1988) Ground-based training for the stimulus rearrangement encountered during space flight. *Acta Otolaryngol Suppl (Stockh)* 460: 87-93
- Reschke MF, Parker DE, Skinner N (1985) Oculer counterrolling. In: Bungo MW, Bagian TM, Bowman MA, Levitan BM (eds) *Results of the Life Sciences DSOs Conducted aboard the Space Shuttle (1981-1986)*. Space Biomedical Research Institute, NASA: Houston, TX, pp 141-144
- Reschke MF, Harm DL, Parker DE, Paloski WH (1991) DSO 459. Otolith tilt-translation reinterpretation. In: *Results of Life Sciences DSOs Conducted Aboard the Space Shuttle, 1988-1990*. NASA: Houston, TX, pp 33-50
- Reschke MF, Harm DL, Parker DE, Sandoz GR, Homick JL, Vanderploeg JM (1994) Neurophysiologic aspects: space motion sickness. In: Nicogossian AE, Huntoon CL, Pool SL (eds) *Space Physiology and Medicine*. Lea & Febiger: Philadelphia, pp 228-260
- Reschke MF, Harm DL, Bloomberg JJ, Paloski WH, Kornilova LN (1996) Neurosensory and sensory-motor function. In: Antipov VV, Grogoriev AI, Leach Huntoon C (eds) *Space Biology and Medicine*. American Institute of Aeronautics and Astronautics: Reston VA, Vol 3, pp 135-193
- Reschke MF, Bloomberg JJ, Harm DL, *et al.* (1998) Posture, locomotion, spatial orientation, and motion sickness as a function of space flight. *Brain Res Rev* 28: 102-117
- Reschke MF, Bloomberg JJ, Harm DL, Huebner WP, Krnavek JM, Paloski WH, Berthoz A (1999) Visual-vestibular integration as a function of adaptation to space flight and return to Earth. In: Sawin CF, Taylor GR, Smith WL (eds) *Extended Duration Orbiter Medical Project Final Report 1989-1995*. NASA: Houston, TX, pp 5.3-1-5.3-41
- Reschke MF, Somers JT, Ford G (2006) Stroboscopic vision as a treatment for motion sickness: strobe lighting vs. shutter glasses. *Aviat Space Environ Med* 77: 2-7
- Reschke MF, Krnavek JM, Somers JT, Ford G (2007a) *A Brief History Of Space Flight With A Comprehensive Compendium Of Vestibular And Sensorimotor Research Conducted Across The Various Flight Programs*. NASA: Houston, TX
- Reschke MF, Krnavek JM, Somers JT, Ford G, Hwang EJ, Leigh RJ, Estrada A (2007b) Stroboscopic vision as a treatment for retinal slip induced motion sickness. *Proceedings of the First International Symposium on Visually Induced Motion Sickness, Fatigue, and Photosensitive Epileptic Seizures VIMS2007* 1: 51-58
- Reschke MF, Bloomberg JJ, Paloski WH, Mulavara AP, Feiveson A, Harm DL (2009) Postural reflexes, balance control, and functional mobility with long-duration head-down bed rest. *Aviat Space Environ Med* 80 (Supp 5): A45-A54

- Reschke MF, Wood SJ, Harm DL, Bloomberg JJ, Paloski WH, Schlegel TT, Clément G (2013) Neurovestibular/Sensorimotor System. In: Risin D, Stepaniak P (eds) *Biomedical Results of the Space Shuttle Program*. NASA: Washington DC, NASA/SP2013-607, Chapter 4.9
- Reschke MF, Cohen HS, Cerisano JM, Clayton JA, Cromwell R, Danielson RW, Hwang EY, Tingen C, Allen JR, Tomko DL (2014) Effects of sex and gender on adaptation to space: neurosensory systems. *Women Health* 23: 959-962
- Reschke MF, Kozlovskaya IB, Kofman IS, Tomilovskaya ES, Cerisano JM, *et al.* (2015) Initial sensorimotor and cardiovascular data acquired from Soyuz landings: Establishing functional performance recovery time constant. *Proceedings of the 20<sup>th</sup> IAA Humans in Space Symposium*, Prague, 28 June-3 July 2015, IAA-HIS-15-1A1-1
- Riani, M., and Simonotto, E. (1994). Stochastic resonance in the perceptual interpretation of ambiguous figures: A neural network model. *Phys Rev Lett* 72, 3120-3123
- Richards JT, Mulavara AP, Bloomberg JJ (2004) Postural stability during treadmill locomotion as a function of the visual polarity and rotation of a three-dimensional environment. *Presence Teleoper Vir Real* 13: 371-384
- Richards JT, Mulavara AP, Bloomberg JJ (2007) The interplay between strategic and adaptive control mechanisms in plastic recalibration of locomotor function. *Exp Brain Res* 78: 326-338
- Richardson, K., Imhoff, T.T., Grigg, P., and Collins, J.J. (1998). Using electrical noise to enhance ability of humans to detect subthreshold mechanical stimuli. *Chaos* 8, 599-603.
- Roberts DR, Ramsey D, Johnson K, Kola J, Ricci R, Hicks C, Borckardt JJ, Bloomberg JJ, Epstein C, George MS (2010) Cerebral cortex plasticity after 90 days of bed rest: data from TMS and fMRI. *Aviat Space Environ Med* 81: 30-40
- Robinson DA (1981) The use of control systems analysis in the neurophysiology of eye movements. *Ann Rev Neurosci* 4: 463-503
- Rogers WF (1972) *Apollo Experience Report-Lunar Module Landing Gear Subsystem*. NASA: Houston, TX
- Roll JP, Popov KE, Gurfinkel VS, Lipshits MI, André-Deshays C, Gilhodes JC, Quoniam C (1993) Sensorimotor and perceptual function of muscle proprioception in microgravity. *J Vestib Res* 3: 259-274
- Roller CA, Cohen HS, Kimball KT, and Bloomberg JJ (2001) Variable practice with lenses improves visuo-motor plasticity. *Cognitive Brain Research*. 12: 341-352, 2001
- Roller CA, Cohen HS, Bloomberg JJ, Mulavara AP (2009) Improvement of obstacle avoidance on a compliant surface during transfer to a novel visual task after variable proactive under unusual visual conditions. *Percept Motor Skills* 108: 173-180
- Ross HE, Brodie EE, Benson AJ (1984) Mass discrimination during prolonged weightlessness. *Science* 225: 219-221
- Ross HE, Brodie EE, Benson AJ (1986a) Mass discrimination in weightlessness and re-adaptation to Earth's gravity. *Exp Brain Res* 64: 358-366
- Ross HE, Schwartz E, Emmerson P (1986b) Mass discrimination in weightlessness improves with arm movements of higher acceleration. *Naturwissenschaften* 73: 453-454
- Ross MD (1993) Morphological changes in rat vestibular system following weightlessness. *J Vestib Res* 3: 241-251

- Ross MD (1994) A space flight study of synaptic plasticity in adult rat vestibular maculas. *Acta Otolaryngol* 516: 1-14
- Ross MD (2000) Changes in ribbon synapses and rough endoplasmic reticulum of rat utricular hair cells in weightlessness. *Acta Otolaryngol* 120: 490-499
- Ross MD, and Varelas J (2003) Ribbon synaptic plasticity in gravity sensors of rats flown on neurolab. In: Buckley JC, Homick JL (eds) *The Neurolab Space Mission: Neuroscience Research in Space*. NASA: Houston, TX
- Ross, S.E. (2007). Noise-enhanced postural stability in subjects with functional ankle instability. *Br J Sports Med* 41, 656-659; discussion 659. doi: 10.1136/bjsm.2006.032912
- Ross, S.E., and Guskiewicz, K.M. (2006). Effect of coordination training with and without stochastic resonance stimulation on dynamic postural stability of subjects with functional ankle instability and subjects with stable ankles. *Clinical journal of sport medicine* 16, 323-328.
- Ross, S.E., Arnold, B.L., Blackburn, J.T., Brown, C.N., and Guskiewicz, K.M. (2007). Enhanced balance associated with coordination training with stochastic resonance stimulation in subjects with functional ankle instability: an experimental trial. *J Neuroeng Rehabil* 4, 47. doi: 10.1186/1743-0003-4-47.
- Ross, S.E., Linens, S.W., Wright, C.J., and Arnold, B.L. (2013). Customized noise-stimulation intensity for bipedal stability and unipedal balance deficits associated with functional ankle instability. *J Athl Train* 48, 463-470. doi: 10.4085/1062-6050-48.3.12.
- Ruttley TM, Harm DL, Evans CA, Robinson JA (2010) International utilization at the threshold of "assembly complete"- science returns from the International Space Station. *Acta Astronautica* 67: 513-519
- Ryder JW, Buxton RE, Goetchius E, Scott-Pandorf M, Hackney KJ, Fiedler J, Ploutz-Snyder RJ, Bloomberg JJ, Ploutz-Snyder LL (2013) Influence of muscle strength to weight ratio on functional task performance. *Eur J Appl Physiol* 113: 911-921
- Sangals J, Heuer H, Manzey D, et al. (1999) Changed visuomotor transformations during and after prolonged microgravity. *Exp Brain Res* 129: 378-390
- Santy PA, Bungo MW (1991) Pharmacologic considerations for Shuttle astronauts. *J Clin Pharmacol* 31: 931-933
- Satake H, Matsunami K, Miyata H (1991) The vestibulo-autonomic function viewed from cardiac responses in centrifuged monkeys. *Acta Otolaryngol Suppl* 481: 543-547
- Schmitt HH, Reid DJ (1985) Anecdotal information on space-adaptation syndrome. In: *Space Adaptation Syndrome Drug Workshop*, Houston, TX
- Schubert MC, Herdman SJ, et al. (2001) Functional measure of gaze stability in patients with vestibular hypofunction. *Ann NY Acad Sci* 942: 490-491
- Schubert MC, Herdman SJ, et al. (2002) Vertical dynamic visual acuity in normal subjects and patients with vestibular hypofunction. *Otol Neurotol* 23: 372-377
- Seaton KA, Slack KJ, Sipes W, et al. (2007) Artificial gravity as a multi-system countermeasure: effects on cognitive function. *J Gravit Physiol* 14: P27-30
- Seidler RD (2010) Neural correlates of motor learning, transfer of learning, and learning to learn. *Exercise Sport Sci Rev* 38: 3-9
- Seidler RD, Noll DC, Chintalapati P (2006) Bilateral basal ganglia activation associated with sensorimotor adaptation. *Exp Brain Res* 175: 544-555

- Seidler, RD, Mulavara AP, Bloomberg JJ, Peters BT (2015) Individual predictors of sensorimotor adaptability. *Front Syst Neurosci*. 2015 July 6;9:100
- Serrador JM, Lipsitz LA, Gopalakrishnan GS, Black FO, Wood SJ (2009) Loss of otolith function with age is associated with increased postural sway measures. *Neurosci Lett* 465: 10-15
- Serrador JM, Schlegel TT, Black FO, Wood SJ (2009) Vestibular effects on cerebral blood flow. *BMC Neurosci* 10: 119
- Shashkov VS, Sabayev VV (1981) Pharmacological agents for the prevention and treatment of motion sickness. *Kosm Biol Aviakosm Med* 15: 9-18
- Shelhamer M, Beaton K (2010) Mission to Mars: training and maintenance of sensorimotor responses: considerations based on context-specific adaptation. *J Cosmology* 12: 3817-3824
- Shelhamer M, Beaton K (2012) Pre-flight sensorimotor adaptation protocols for suborbital flight. *J Vestib Res* 22: 139-144
- Shelhamer, M (2015) Trends in sensorimotor research and countermeasures for exploration-class space flights. *Front Syst Neurosci*. 2015 Aug 11;9:115
- Shepard RN, Metzler J (1971) Mental rotation of three-dimensional objects. *Science* 171: 701-703
- Shirley RS, Young LR (1968) Motion cues in man-vehicle control. *IEEE Trans Mach Sys* 9: 121-128
- Shortt TL, Ray CA (1997) Sympathetic and vascular responses to head-down neck flexion in humans. *Am J Physiol* 272: H1780-H1784
- Shupak A, Gordon CR, Spitzer O, Mendelowitz N, Melamed Y (1989) Three-years' experience of transdermal scopolamine: long-term effectiveness and side-effects. *Pharmatherapeutica* 5: 365-370
- Simonotto, E., Riani, M., Seife, C., and Oberts, M. (1997). Visual perception of stochastic resonance. *Physical Review Letters* 256, 6-9. doi: <http://link.aps.org/doi/10.1103/PhysRevLett.78.1186>
- Sirota MG, Babayev BM, Beloozerova IB, Nyrova AN, Yakushin SB, Kozlovskaya IB (1987) Characteristics of vestibular reactions to canal and otolith stimulation at an early stage of exposure to microgravity. *Physiologist* 30: S82-S84
- Slack KJ, Williams TJ, Schneiderman JS, Picano JJ, Leveton LB, Schmidt LL, Shea C (2016) Evidence Report: Risk of Adverse Cognitive or Behavioral Conditions and Psychiatric Disorders. Human Research Program, NASA Johnson Space Center, Houston, Texas. Available at: <https://humanresearchroadmap.nasa.gov/evidence/reports/BMed.pdf?rnd=0.541643132707869>
- Smith JW, Edwards JW (1980) *Design of a Nonlinear Adaptive Filter for Suppression of Shuttle Pilot-Induced Oscillation Tendencies*. DFRC
- Smith SM, Heer MA, Shackelford LC, Sibonga JD, Ploutz-Snyder L, Zwart SR (2012) Benefits for bone from resistance exercise and nutrition in long-duration spaceflight: Evidence from biochemistry and densitometry. *J Bone Miner Res* 27: 1896-1906
- So CW, Bent LR (2009) Increased vestibular contribution to posture control in individuals with chronic headache. *J Vestib Res Equilib Orientat* 19: 49-58
- Soma R, Nozaki D, Kwak S, Yamamoto Y. 1/f Noise outperforms white noise in sensitizing baroreflex function in human brain. *Physical Review Letters*. 2003;91(7):078101-1-4

- Spiering BA, Lee SMC, Mulavara AP, Bentley JR, Buxton RE, Lawrence EL, Sinka J, Guilliams ME, Ploutz-Snyder LL, Bloomberg JJ (2011) Test battery designed to quickly and safely assess diverse indices of neuromuscular function following unweighting. *J Strength Cond Res* 25: 545-555
- Strangman GE, Sipes W, Beven G (2014) Human cognitive performance in spaceflight and analogue environments. *Aviat Space Environ Med* 85:1033-1048
- Stefanucci JK, Proffitt DR (2009) The roles of altitude and fear in the perception of height. *J Exp Psychol* 35: 424-438
- Stephen, D.G., Wilcox, B.J., Niemi, J.B., Franz, J.R., Kerrigan, D., and D'andrea, S.E. (2012). Baseline-dependent effect of noise-enhanced insoles on gait variability in healthy elderly walkers. *Gait Posture* 36, 537-540. doi: 10.1016/j.gaitpost.2012.05.014
- Stott JR, Barnes GR, Wright RJ, Ruddock CJ (1989) The effect on motion sickness and oculomotor function of GR 38032F, a 5-HT<sub>3</sub>-receptor antagonist with anti-emetic properties. *Br J Clin Pharmacol* 27: 147-157
- Streeper T, Cavanagh PR, Hanson AM, Carpenter RD, Saeed I, Kornak J (2011) Development of an integrated countermeasure device for use in long-duration spaceflight. *Acta Astronautica* 68: 2029-2037
- Streepey JW, Kenyon RV, Keshner EA (2007) Visual motion combined with base of support width reveals variable field dependency in healthy young adults. *Exp Brain Res* 176: 182–187
- Strzalkowski ND, Lowrey CR, Perry SD, Williams DR, Wood SJ, Bent LR (2015) Selective weighting of cutaneous receptor feedback and associated balance impairments following short duration space flight. *Neurosci Lett* 592: 94-98
- Symons TB, Sheffield-Moore M, Chinkes DL, Ferrando AA, Paddon-Jones D (2009) Artificial gravity maintains skeletal muscle protein synthesis during 21 days of simulated microgravity. *J Appl Physiol* 107: 34-38
- Task HL, Genco LV (1987) Effects of short-term flight on several visual functions. In: Bungo MW, Bagian T, Bowman MA, Levitan BM (eds) *Results of the Life Sciences DSO Conducted Aboard the Space Shuttle, 1981-1986*. NASA: Houston, TX
- Taube JS, Stackman RW, Calton JL, Oman CM (2004) Rat head direction cell responses in zero-gravity parabolic flight. *J Neurophysiol* 92: 2887-2997
- Thornton WE, Rummel J (1977) Muscular deconditioning and its prevention in space flight. In: Johnston RS, Dietlein LF (eds) *Biomedical Results of Skylab*. NASA: Washington DC: pp 191-197
- Thornton WE, Biggers WP, Thomas WG, Pool SL, Thagard NE (1985) Electronystagmography and audio potentials in space flight. *Laryngoscope* 95: 924-932
- Thornton WE, Moore TP, Pool SL, Vanderploeg JM (1987a) Clinical characterization and etiology of space motion sickness. *Aviat Space Environ Med* 58: A1-A8
- Thornton WE, Linder BJ, Moore TP, Pool SL (1987b) Gastrointestinal motility in space motion sickness. *Aviat Space Environ Med* 58: A16-A21
- Thornton WE, Moore TP, Uri J., Pool SL (1988) *Studies of the Vestibulo-Ocular Reflex on STS 4, 5, and 6*. NASA: Houston, TX
- Thornton WE, Uri JJ, Moore T, Pool S (1989) Studies of the horizontal vestibulo-ocular reflex in space flight. *Arch Otolaryngol Head Neck Surg* 115: 943-949

- Tian JR, Shubayev I, Demer JL (2001) Dynamic visual acuity during transient and sinusoidal yaw rotation in normal and unilaterally vestibulopathic humans. *Exp Brain Res* 137: 12-25
- Tomilovskaya ES, Berger M, Gerstunbranel F, Kozlovskaya IB (2011) Effects of long-duration space flight on target acquisition. *Acta Astronautica* 68: 1454-1464
- Treisman M (1977) Motion sickness: an evolutionary hypothesis. *Science* 197: 493-495
- Uri JJ, Linder BJ, Moore TD (1989) *Saccadic Eye Movements during Space Flight*. NASA: Houston, TX
- van Hedel HJA, Dietz V (2004) The influence of age on learning a locomotor task. *Clin Neurophysiol Off J Int Fed Clin Neurophysiol* 115: 2134–2143.
- van Marion WF, Bongaerts MC, Christiaanse JC, Hofkamp HG, van Ouwerkerk W (1985) Influence of transdermal scopolamine on motion sickness during 7 days exposure to heavy seas. *Clin Pharmacol Ther* 38: 301-305
- Viaud-Delmon I, Ivanenko YP, Berthoz A, Jouvent R (2000) Adaptation as a sensorial profile in trait anxiety: a study with virtual reality. *J Anxiety Disord* 14: 583–601
- Vieville T, Clément G (1986) Adaptive modifications of the optokinetic vestibulo-ocular reflexes in microgravity. In: Zee DS (ed) *Adaptive Processes in Visual and Oculomotor Systems*. Pergamon Press: New York, pp 111-120
- Vogel H, and Kass JR. European vestibular experiments on the Spacelab-1 mission: 7. Ocular counterrolling measurements pre- and post-flight. *Exp Brain Res* 64: 284-290, 1986.
- von Baumgarten R, Benson AJ, Berthoz A, et al. (1986) European experiments on the vestibular system during the Spacelab D-1 mission. In: Sahm PR, Jansen R, Keller MH (eds) *Proceedings of the Symposium on Scientific Results of the German Spacelab Mission D-1*, Norderney, Germany, 27-29 August 1986, pp 477-490
- von Baumgarten RJ, Kass J, Vogel H, Wetzig J (1989) Influence of proprioceptive information on space orientation on the ground and in orbital weightlessness. *Adv Space Res* 9: 223-230
- von Baumgarten RJ, Thuemler R (1978) A model for vestibular function in altered gravitational states. *Life Sci Space Res* 17: 161-170
- von Baumgarten RJ, Wetzig J, Vogel H, Kass JR (1982) Static and dynamic mechanisms of space vestibular malaise. *Physiologist* 25: S33-S36
- von Holst E, Mittelstaedt H (1973) The reafference principle (interaction between the central nervous system and the periphery). In: Martin R (ed) *The Behavioural Physiology of Animals and Man*. Coral Gables: University of Miami Press, pp 139-173
- Vorob'ev EI, Gazenko OG, Gurovskii NN, Nefedov IuG, Egorov BB (1975) Preliminary results of medical investigations carried out during flight of the second expedition of the 'Salyut-4' orbital station. *Kosm Biol Aviakosm Med* 10: 3-18
- Wagner SA (2006) *The Apollo Experience Lessons Learned for Constellation Lunar Dust Management*. NASA/TP-213726; JSC, Sep 2006.
- Ward LM, Neiman A, and Moss F (2002). Stochastic resonance in psychophysics and in animal behavior. *Biol Cybern* 87, 91-101. doi: 10.1007/s00422-002-0328-z
- Warren LE, Reinerstson R, Camacho ME, Paloski WH (2007) Implementation of the NASA artificial Gravity bed rest pilot study. *J Gravit Physiol* 14: P1-P4
- Warren WH, Kay BA, Yilmaz EH (1996) Visual control of posture during walking: functional specificity. *J Exp Psychol Hum Percept Perform* 22: 818–838

- Watt DG, Money KE, Bondar RL, Thirsk RB, Garneau M, Scully-Power P (1985) Canadian medical experiments on Shuttle flight 41-G. *Can Aeronaut Space J* 31: 215-226
- Watt DG (1987) The vestibulo-ocular reflex and its possible roles in space motion sickness. *Aviat Space Environ Med* 58: A170-174
- Watt DGD, Money KE, Tomi LM (1986) MIT/Canadian vestibular experiments on the Spacelab-1 mission: 3. Effects of prolonged weightlessness on a human otolith-spinal reflex. *Exp Brain Res* 64: 308-379
- Watt DGD, Landolt JP, Young LR (1992) Effects of long-term weightlessness on roll-circularvection. *Proceedings of the 7th CASI Conference on Astronautics*, Ottawa, Canada
- Watt DGD (1997) Pointing at memorized targets during prolonged microgravity. *Aviat Space Environ Med* 68: 99-103
- Welch RB (1986) Adaptation of space perception. In: Boff KR, Kaufman L, Thomas JP (eds) *Handbook of Perception and Human Performance*. John Wiley & Sons: New York, pp 24-145
- Welch R, Hoover M, Southward EF (2009) Cognitive performance during prismatic displacement as a partial analogue of "space fog." *Aviat Space Environ Med* 80: 771-780
- Wertheim AH, Mesland BS, Bles W (2001) Cognitive suppression of tilt sensations during linear horizontal self-motion in the dark. *Perception* 30: 733-741
- Wetzig J, Hofstetter-Degen K, von Baumgarten R (1993) Responses to eccentric rotation in two space-bound subjects. *Clin Investig* 71: 757-760.
- Wiener SI, Taube JS (2005) *Head Direction Cells and the Neural Mechanisms of Spatial Orientation*. MIT Press Bradford Books: Cambridge, MA
- Williamson P (2007) Mobile Servicing Systems Lessons Learned. In: *DX23 ISS Mechanical and Robotics Systems Training*. NASA: Houston, TX
- Wood CD (1979) Antimotion sickness and antiemetic drugs. *Drugs* 17: 471-479
- Wood CD (1990) Pharmacological countermeasures against motion sickness. In: Crampton GH (ed) *Motion and Space Sickness*. CRC Press: Boca Raton, FL, pp 343-351
- Wood CD, Graybiel A (1968) Evaluation of sixteen antimotion sickness drugs under controlled laboratory conditions. *Aerospace Med* 39: 1341-1344
- Wood CD, Graybiel A (1970) Evaluation of antimotion sickness drugs: a new effective remedy revealed. *Aerospace Med* 41: 932-933
- Wood CD, Graybiel A (1972) Theory of antimotion sickness drug mechanisms. *Aerospace Med* 43: 249-252
- Wood CD, Manno JE, Manno BR, Odenheimer RC, Bairnsfather LE (1986) The effect of antimotion sickness drugs on habituation to motion. *Aviat Space Environ Med* 57: 539-542
- Wood CD, Manno JE, Wood MJ, Manno BR, Redetzki HM (1987a) Mechanisms of antimotion sickness drugs. *Aviat Space Environ Med* 58: A262-A265
- Wood CD, Stewart JJ, Wood MJ, et al. (1994) Habituation and motion sickness. *J Clin Pharmacol* 34: 628-634
- Wood MJ, Wood CD, Manno JE, Manno BR, Redetzki HM (1987b) Nuclear medicine evaluation of motion sickness and medications on gastric emptying time. *Aviat Space Environ Med* 58: 1112-1114
- Wood SJ (2002) Human otolith-ocular reflexes during off-vertical axis rotation: Effect of frequency on tilt-translation ambiguity and motion sickness. *Neurosci Letters* 323: 41-44

- Wood SJ, Black FO, Reschke MF, Kaufman GD, Paloski WH (2009) Effects of eccentric rotation on the human pitch vestibulo-ocular reflex. *Acta Oto-Laryngologica* 129: 521-526
- Wood SJ, Loehr JA, Williams M (2011) Sensorimotor reconditioning during and after spaceflight. *NeuroRehabilitation* 29: 185-195
- Wood SJ, Reschke MF, Black FO (2012) Continuous equilibrium scores: Factoring in the time before a fall. *J Gait Posture* 36: 487-489
- Wood SJ, Paloski WH, Clark JB (2015) Assessing sensorimotor function following ISS with computerized dynamic posturography. Spaceflight with Computerized Dynamic Posturography. *Aerosp Med Hum Perform* 86(12 Suppl): A45-53.
- Woodard D, Knox G, Wood S, DiBiase C (2014) Off-vertical axis rotational assessment of transdermal scopolamine for motion sickness prophylaxis. *Gravit Space Res* 2: 13-20
- Woodring SF, Rossiter CD, Yates BJ (1997) Pressor response elicited by nose-up vestibular stimulation in cats. *Exp Brain Res* 113: 165-168
- Wright WG, DiZio P, Lackner JR (2006) Perceived self-motion in two visual contexts: dissociable mechanisms underlie perception. *J Vestib Res* 16: 23-28
- Xerri C, Barthelemy J, Harlay F, Borel L, Lacour M (1987) Neuronal coding of linear motion in the vestibular nuclei of the alert cat. I. Response characteristics to vertical otolith stimulation. *Exp Brain Res* 65: 569-581
- Yates BJ (1992) Vestibular influences on the sympathetic nervous system. *Brain Res Rev* 17: 51-59
- Yates BJ (1998) Autonomic reaction to vestibular damage. *Otolaryngol Head Neck Surg* 119: 106-112
- Yates BJ, Goto T, Bolton PS (1993) Responses of neurons in the rostral ventrolateral medulla of the cat to natural vestibular stimulation. *Brain Res* 601: 255-264
- Yates BJ, Kerman IA (1998) Post-space flight orthostatic intolerance: possible relationship to microgravity-induced plasticity in the vestibular system. *Brain Res Rev* 28: 73-82
- Yates BJ, Miller AD (1996) *Vestibular Autonomic Regulation*. CRC Press, Boca Raton FL
- Yates BJ, Miller AD (1998) Physiological evidence that the vestibular system participates in autonomic and respiratory control. *J Vestib Res* 8: 17-25
- Yates BJ, Miller AD, Lucot JB (1998) Physiological basis and pharmacology of motion sickness: an update. *Brain Res Bull* 47: 395-406
- Yegorov AD (1979) *Results of Medical Studies during Long-Term Manned Flights on the Orbital Salyut-6 and Soyuz Complex*. USSR Academy of Sciences, NASA TM-76014
- Young LR (1967) Some effects of motion cues on manual tracking. *J Spacecraft* 4: 1300-1303
- Young LR (1969) On adaptive manual control. *Ergonomics* 12: 635-674
- Young LR (1974) Perceptions of the body in space: mechanisms. In: Darian-Smith I (ed) *Handbook of Physiology - The Nervous System*. American Physiological Society: Bethesda MD, pp 1023-1066
- Young LR (2003) Spatial Orientation. In: Vidulich PT (ed) *Principles and Practice of Aviation Psychology*. Earlbaum: Mahwah NJ, pp 69-144
- Young LR, Oman CM (1969) Model for vestibular adaptation to horizontal rotation. *Aerospace Med.* 40: 1076-1080
- Young LR, Shelhamer M (1990) Microgravity enhances the relative contribution of visually-induced motion sensation. *Aviat Space Environ Med* 61: 525-530

- Young LR, Sinha P (1998) Space flight influences on ocular counterrolling and other neurovestibular reactions. *Otolaryngol Head Neck Surg* 118: S31-34
- Young LR, Zuber BL, Stark L (1967) *Visual and Control Aspects of Saccadic Eye Movements*. NASA CR-564. NASA Contractor Report
- Young LR, Dichgans J, Murphy R, Brandt T (1973) Interaction of optokinetic and vestibular stimuli in motion perception. *Acta Otolaryngol* 76: 24-31
- Young LR, Oman CM, Watt DG, Money KE, Lichtenberg BK (1984) Spatial orientation in weightlessness and readaptation to earth's gravity. *Science* 225: 205-208
- Young LR, Oman CM, Watt DGD, *et al.* (1986) MIT/Canadian vestibular experiments on the Spacelab-1 mission: 1. Sensory adaptation to weightlessness and re-adaptation to 1 g: An overview. *Exp Brain* 64: 291-298
- Young LR, Oman CM, Merfeld D, *et al.* (1993) Spatial orientation and posture during and following weightlessness: human experiments on Spacelab Life Sciences. *J Vestib Res* 3: 231-239
- Young LR, Paloski WH (2007) Short radius intermittent centrifugation as a countermeasure to bed-rest and 0-G deconditioning: IMAG pilot study summary and recommendations for research. *J Gravit Physiol* 14: P31-P33
- Zangemeister WH, Bulgheroni MV, Pedotti A (1991) Normal gait is differentially influenced by the otoliths. *J Biomed Eng* 13: 451-458
- Zangemeister WH, Schlueter D, Kunze K (1988) Compensatory eye movement gain and head-eye latencies change with verbal feedback: Voluntary adjustment of gaze types. *Adv Otorhinolaryngol* 41: 227-234
- Zangemeister WH, Stark L (1981) Active head rotations and eye-head coordination. *Ann NY Acad Sci* 374: 540-559
- Zangemeister WH, Stark L (1982) Gaze latency: variable interactions of head and eye latency. *Exp Neurol* 75: 389-406
- Zeng, F.G., Fu, Q.J., and Morse, R. (2000). Human hearing enhanced by noise. *Brain Research* 869, 251-255
- Zupan LH, Peterka RJ, Merfeld DM (2000) Neural processing of gravito-inertial cues in humans. I. Influence of the semicircular canals following post-rotatory tilt. *J Neurophysiol* 84: 2001-2015
- Zupan LH, Merfeld DM (2005) An internal model of head kinematics predicts the influence of head orientation on reflexive eye movements. *J Neural Eng* 2: S180-S197.
- Zwart SR, Crawford GE, Gillman PL, Kala G, Rodgers A, Rogers A, *et al.* (2009) Effects of 21 days of bed rest, with or without artificial gravity, on nutritional status of humans. *J Appl Physiol* 107: 54-62

## **XI. TEAM**

Jacob J. Bloomberg, Ph.D., Sensorimotor Lead Scientist, Neurosciences Laboratory, Biomedical Research & Environmental Sciences Division, Johnson Space Center, Houston, TX Johnson Space Center, Houston, TX.

Millard F. Reschke, Ph.D., Chief of Neuroscience, Neurosciences Laboratory, Biomedical Research & Environmental Sciences Division, Johnson Space Center, Houston, TX.

Gilles R. Clément, Ph.D., Artificial Gravity Lead Scientist (Wyle Science Technology & Engineering Group), Human Health Countermeasures Element, Biomedical Research & Environmental Sciences Division, Johnson Space Center, Houston, TX.

Ajitkumar P. Mulavara, Ph.D., Senior Research Scientist (Universities Space Research Association), Neurosciences Laboratory, Biomedical Research & Environmental Sciences Division, Johnson Space Center, Houston, TX.

Laura C. Taylor, M.S., Technical Lead/Biomedical Engineer (Wyle Science Technology & Engineering Group), Neurosciences Laboratory, Biomedical Research & Environmental Sciences Division, Johnson Space Center, Houston, TX.

The authors are indebted to Elizabeth Fisher, Elisa Allen, and Janet Cook for their assistance with the collation and organization of the report references.

## **XII. LIST OF ACRONYMS**

AFT	Autogenic Feedback System
ARED	Advanced Resistive Exercise Device
CDP	Computerized Dynamic Posturography
CEVIS	Cycle Ergometer with Vibration Isolation System
CFIT	Controlled Flight Into Terrain
CMD	Commander
CNS	Central Nervous System
COLBERT	Combined Operational Load Bearing External Resistance Treadmill
Dec	Decrease
DIS	Document Index System
DOF	Degree of Freedom
DOME	Device for Orientation and Motion Environments
DOUG	Dynamic Onboard Ubiquitous Graphics
DVA	Dynamic Visual Acuity
EAFB	Edwards Air Force Base
ECV	Exploration Class Vehicle

EOM	Effective Oculomotor Range
EQ	Equilibrium
EVA	Extra-Vehicular Activity
FMT	Functional Mobility Test
FTT	Functional Task Test
GCA	Ground Control Assist
GCR	Galactic Cosmic Rays
GIA	Gravitoinertial Acceleration
GPWS	Ground Proximity Warning System
GVS	Galvanic Vestibular Stimulation
HAC	Heading Alignment Cone
HUD	Head Up Display
HRP	Human Research Program
IBMP	Russian Institute of Biomedical Problems
IM	Intramuscular
Inc	Increase
iRED	interim Resistive Exercise Device
ISS	International Space Station
KSC	Kennedy Space Center
LCD	Liquid Crystal Display
LEE	Latching End Effector
LM	Apollo Lunar Module
MCC	Mission Control Center
MS	Mission Specialist
MSBLS	Microwave Scanner Beam Landing System
MT	Mobile Transporter
NTS	Nucleus tractus Solitarius
NTSB	National Transportation Safety Board
NVA	Near Visual Acuity
OCR	Ocular Counterrolling
OKN	Optokinetic Nystagmus
ONTO	Oxygen/Nitrogen tank
OTTR	Otolith Tilt-Translation Reinterpretation
OVAR	Off Vertical Axis Rotation
PAPI	Precision Approach Path Indicator
PFMS	Post-Flight Motion Sickness
PIO	Pilot Induced Oscillation
PLT	Pilot
PS	Payload Specialist
PSOI	Post-Spaceflight Orthostatic Intolerance
ROTTR	Rotation Otolith Tilt-Translation Reinterpretation
SA	Sensorimotor Adaptability
SD	Spatial Disorientation

SFRM	Space Flight Resource Management
SM	Sensorimotor
SMS	Space Motion Sickness
SPE	Solar Particle Event
SR	Stochastic Resonance
SSRMS	Space Station Remote Manipulator System
STA	Shuttle Training Aircraft
STS	Space Transportation System (Space Shuttle)
TACAN	Tactical Air Navigation
TAEM	Terminal Area Energy Management
TAWS	Terrain Awareness and Warning System
TCC	Time to Complete the Course
TSAS	Tactile Spatial Awareness System
TTD	Tilt-Translation Device
TVIS	Treadmill Vibration Isolation System
UHF	Ultra High Frequency
VEMP	Vestibular Evoked Myogenic Potentials
VIIP	Vision Impairment and Intercranial Pressure
VOR	Vestibulo-Ocular Reflex
VRI	Visual Reorientation Illusion
VRT	Vestibular Rehabilitation Therapy

### XIII. APPENDIX A: SHUTTLE LANDING PERFORMANCE

**Authors: Millard F. Reschke, Deborah L. Harm, William H. Paloski, Jacob J. Bloomberg, Scott J. Wood**

#### **Shuttle Landing Performance**

Control of vehicles, and other complex mechanical motion systems, is a high-level integrative function of the central nervous system (CNS) that requires good visual acuity, eye-hand coordination, spatial (and, in some cases, geographic) orientation perception, and cognitive function. Existing evidence from space flight research (Paloski *et al.*, 2008; Clément and Reschke 2008; Reschke *et al.*, 2007) demonstrates that the function of each of these systems is altered by removing (and subsequently by reintroducing) a fundamental orientation reference (gravity), which is sensed by vestibular, proprioceptive, and haptic receptors and used by the CNS for spatial orientation, navigation, and coordination of movements. The available evidence also shows that the degree of alteration of each system depends on crew experience and mission-related factors.

There is only limited operational evidence that alterations in the gravito-inertial force environment will cause functional impacts on mission-critical vehicle (or complex system) control capabilities. Furthermore, while much of the operational performance data collected during space flight has not been available for independent analysis, those that have been reviewed are somewhat equivocal owing to uncontrolled (and/or unmeasured) environmental and/or engineering factors.

Thus, our current understanding of the ***Risk of Impaired Control of Spacecraft/Associated Systems and Decreased Mobility Due to Vestibular/Sensorimotor Alterations Associated with Space Flight*** is limited primarily to extrapolation of scientific research findings, and, since there are no robust ground-based analogs of the sensory-motor changes associated with space flight, observation of their functional impacts is limited to studies performed in the space flight environment. Fortunately, more than 500 sensorimotor experiments have been performed during and/or after space flight missions since 1959 (reviewed by Reschke *et al.* 2007). While not all of these experiments were directly relevant to the question of vehicle/complex system control, many provide insight into changes in aspects of sensory-motor control that might have a bearing on the physiological subsystems underlying this high-level integrated function.

#### **Evidence Obtained from Observation, Analysis, and/or First Hand Reports of Space Flight Operations**

An accurate assessment of the risks posed by the impacts of physiological and psychological adaptations to space flight on control of vehicles and other complex systems must account for the potentially off setting influences of training/recency and engineering aids to task performance. Thus, it behooves us to review performance data obtained from space flight crews engaged in true mission operations. Evidence of operational performance decrements during space flight missions has been obtained from several sources; however, to our knowledge no well-designed scientific studies have been performed on critical operational

task performance, so interpretation is frequently confounded by small numbers of observations, inconsistent data collection techniques, and/or uncontrolled engineering and environmental factors. Much of the relevant, existing operational data has been previously inaccessible to life sciences researchers. Recent programmatic changes have putatively improved access to both data and experts to help with interpretation. However, a real understanding of the factors involved in manual control will come with dedicated investigations that specifically address the problem.

## Landing the Space Shuttle

The space shuttle landing process begins on the opposite side of the world from the primary landing strip with a deorbit burn approximately one hour before touchdown. If the primary runway is located at the Kennedy Space Center (KSC) reentry begins over the Indian Ocean off of the western coast of Australia. Thirty minutes before landing the orbiter enters the Earth's atmosphere at an altitude of 121,920 m and flies across the Pacific Ocean, the Baja Peninsula, Mexico, southern Texas, and out over the Gulf of Mexico to the west coast of Florida. Adapted to a weightless environment, the entry process places an inertial load on the shuttle crew that lasts for a minimum of 5 min and ranges from 1.2 to 1.6-g with the summed gravitoinertial acceleration (GIA) vectors (gravity and centripetal) located primarily through the Z-axis.

Within approximately 5.5 min prior to landing, the shuttle is traveling at 2,735 km/hr (1,700 mph) at an altitude of 25,338 m, and the crew is conserving the falling spacecraft's energy needed to reach the landing strip (Terminal Area Energy Management or TAEM). The final approach into KSC begins with runway alignment (first waypoint in the final landing process). As illustrated in Figure 1, runway alignment is accomplished by flying the shuttle around one of two imaginary cones (heading alignment cones or HAC) about 11 km from the end of the runway (threshold). Depending on the runway, approach and available energy, flight around the HAC (from 0° to 360°) can be low energy (shallow bank angle) or high energy (higher bank angle). A high energy HAC can peak at close to 2.0-g and present potential manual control problems to the weightless adapted commander and pilot.




Figure 1. Path of the space shuttle into KSC.

The shuttle leaves the HAC at what is called the runway entry point, descending on a 20-22° glide slope within 12 km (7.5 mi) to the runway threshold (Figure 2) at 682 km/h (424 mph). In addition to the head-up display (HUD), which most commanders we have interviewed

indicate is used primarily on roll out to the final approach, there are other landing aids that can help guide the shuttle to a safe landing. These include: (1) the Tactical Air Navigation (TACAN) system that provides range and bearing, (2) the Microwave Scanner Beam Landing System (MSBLS) that provides precise close-in information on slant range, azimuth and elevation, (3) the Precision Approach Path Indicator (PAPI) lights that show the commanders if they are on the correct outer glide slope, and (4) the Ball-Bar Light system that provides a visual reference to the inner glide slope and height off of the runway surface (e.g. high or low).




Figure 2. Final approach of the shuttle.

At 17 sec to touchdown with the spacecraft 1,079 m from the runway threshold and traveling at approximately 496 km/h (308 mph) the commander completes a pre-flare maneuver bringing the shuttle 1.5° nose-up on a 1.5° glide slope. At 14 sec to touchdown the wheels are lowered (335 m altitude - 100 ft) at 496 km/h (270 mph). Touchdown occurs 689 m (2,270 ft) down range from the runway threshold with an air speed of 346 km/h (215 mph). The KSC runway is 4,572 m (15,000 ft) long and 91.4 m (300 ft) wide while the alternate orbiter landing site at Edwards Air Force Base (EAFB) has several dry lake bed runways and one hard surface runway. The longest strip at EAFB is 12 km long (7.5 miles). There are a number of parameters that can be traded off to achieve a safe landing, some specific values that are highly desirable and others that are mandatory to avoid potential loss of the spacecraft during the landing process. The preferred touchdown should occur downrange on the runway from 2200-2700 ft. Less than 1000 ft is considered dangerous and more than 5000 ft risks brake and tire problems. Crossrange the spacecraft should touch down within  $\pm 12$  ft (optimal) of the runway center, but keeping the vehicle between  $\pm 50$  ft is acceptable. Optimal touchdown airspeed ranges between 210 kts and 185 kts depending on vehicle weight. Optimal sink rate is between 2 and 3 fps, however anything lower than 4 fps is acceptable. A pitch attitude between 8-10 deg is optimal, not to exceed 14 deg (14.6 deg will result in tail-scrape) at touchdown.


### Shuttle Entry and Landing Spatial Disorientation

Despite recent (relative to launch), intensive training for all shuttle commanders and pilots, all shuttle landings have not been performed as well as desired, perhaps in part as a result the loss of situational awareness and out right spatial disorientation. Shuttle entry and landing spatial disorientation (SD) differs from aviation SD, at least in terms of prevalence. Almost every instrument rated aircraft pilot has experienced SD, but episodes occur relatively infrequently in ordinary flying. In contrast, stimuli capable of producing SD are present during every Shuttle



data from nine missions, and noted trends, such as an apparent correlation between down range touchdown, sink rate and difficulty arising from a chair without using the arms. Generally scores indicative of neurovestibular dysfunction correlated with flying a lower approach and landing shorter or longer, faster, and harder. When plotted as a function of generalized gaze problems observed by the flight surgeons, there were only four landings within the optimal range on the runway. When gaze was judged as severely affected the landings were clearly short of the desired touchdown point (Figure 3). These observations suggest that further analysis of Shuttle landing performance is warranted.

Both optimal and less than optimal landing are charted in Figure 4. This figure shows the landing of 58 shuttle flights for which we could obtain accurate speed of the vehicle over the threshold. It is clear that only a few of the landing occur within the optimal touchdown area, with many landing short of this area, but more importantly, plotting the landing in this fashion shows the trade-offs that can be used to safely put the shuttle on the ground.



**Figure 4.** 3-D plot of 58 shuttle landings, superimposed on the landing strip, showing the touchdown point (crossrange = Y, downrange = X and altitude over the runway threshold = Z) associated with each landing. Sink rates are indicated by specific colors (yellow = < 2 fps, green = 2 to 3 fps, blue = 3.1 – 4.0 fps and red indicates sink rates greater than 4.0 fps). Airspeed velocities over the runway threshold are shown as symbols (Circle indicates no data is available, left facing triangle = < 223 kts, square = 223 – 231 kts, diamond = 232– 238 kts and the star = velocities > 238 kts). The dashed and solid lines connecting the pinheads to the runway show altitude (solid = optimal height off of the runway surface and dashed = altitudes either higher or lower than optimal). The blue square represents the optimal runway touchdown area, and the red line on the Z-axis shows touchdown clearly short of the optimal downrange distance required for a safe landing.

When additional flights are added (where we do not have threshold velocity, but do have touchdown velocity) it can be seen that several flights have been extremely close to the runway threshold, and at least one flight landed (at EAFB) short of the runway threshold (Figure 5). All of the yellow symbols have less than the optimal sink rates. However, touchdown velocity for these flights suggests that there was (in most cases) energy available for successful downrange landings. The data we have from some of these short landing has indicated that there may have been some difficulty while flying the HAC, including significant spatial disorientation. Furthermore, several commanders have indicated that they believed they were higher off the runway surface than they actually were, and that the spatial disorientation may have involved issues with their perception of speed (motion parallax).

Most commanders interviewed have indicated that perceptually it is not possible to accurately judge sink rates. This could be related to not only motion parallax, but to the difficulty most have in accurately determining vertical position when translated in the Z-axis parallel to the gravitational vector. In any case, of all the landings between STS-1 and STS-108, the Shuttle crossed the runway threshold abnormally low 20 times, and vehicle speeds at touchdown were fast or slow of the targeted value 23 times (10 exceeded flight rules and two exceeded the tire speed limit). Seven landings touched down abnormally long or short, and 13 had high touchdown sink rates, with three exceeding the 5 ft/s structural limit.

Regardless of the commanders ability to land the shuttle following stays of various length in a weightless environment, there are other critical aspects of vehicular control that relate to Lunar landing, docking while on orbit and control of the remote arm found on both the shuttle and International Space Station. In all of these situations eye-head-hand coordination, perception of speed (sink rates and closing velocities), motion parallax in a visually barren environment, and vection or perception of self/surround motion could be critical components in the control of a spacecraft, lunar rover or remote 3-D manipulation system.




Figure 5. Sink rate, roll angle, and y-position at touchdown are a result of atmospheric conditions and how well the commander flies the vehicle.

The following graphs summarize shuttle landing performance data across several landing parameters including:

**Touchdown Downrange Position (Distance from the Threshold)**

Nominal range from 2200-2700 ft. NOTE: Pilots have mentioned that 1000 ft should be the minimum as it is dangerously close.

**Touchdown Sink Rate (Altitude rate)**

Vertical Speed at touchdown; Nominal = -2 to -3 fps, not to exceed -4 fps.

**Altitude over Threshold**

Height over runway start; Nominal = 23-29 ft.

**Touchdown Groundspeed**

High groundspeed at touchdown results in excessive tire wear and possibly tire failure. Tires are approved up to 225 kts.

**Touchdown Pitch Attitude**

Nose-up pitch at landing. Nominal = 8-10°, not to exceed 14°.

**Touchdown Crossrange**


Distance from centerline (Y position) on runway position; nominal range is  $0 \pm 12$ .

**Touchdown Equivalent Airspeed**

Lightweight vehicles: 195 KEAS (+5, -10)

Heavyweight vehicles: 205 KEAS (+5, -10)

### Touchdown Position on Runway



## Downrange Position

The approach and landing trajectory is designed to nominally touchdown the orbiter 2500 ft downrange at a safe speed and small altitude rate. Most runways also have painted markers at the nominal 2500-ft touchdown location to give the pilot's a downrange point of reference.

Nominal Range: 2200-2700 ft.


Pilot-Recommended Minimum: 1000 ft.

(*Landing Procedures, 4-21, p.84*)

62 shuttles landed short of the target minimum of 2200 feet.


7 landed less than 1000 feet from the threshold.

MISSION	DOWNRANGE
STS-37	-623
STS-39	169
STS-33	740
STS-2	780
STS-79	807
STS-4	948
41-G	962



### Cross range (Y-Position)

Nominal:  $0 \pm 12$ .




## Altitude over Threshold

Height over start of runway.

Nominal: 23-29 ft

NOTE: Due to the limited availability of landing this chart reflects data from 58 shuttle landings from STS-65 to STS-124.




## Sink Rate

The sink rate, roll angle, delta heading, and y-position at touchdown are a result of atmospheric conditions and how well the commander flies the vehicle through them. The Final Flare phase is used to reduce the sink rate from about 12 fps on the IGS to less than -3 fps for touchdown. It is also used to control the time and point of touchdown so that the target touchdown airspeed is achieved.

Nominal: -2 to -3 fps, not to exceed -4 fps

13 missions had sink rates higher than -3 fps, out of which 5 missions landed with higher sink rates than the adequate -4.0 fps cutoff. STS-90 had the highest sink rate. The landing touch down appeared harder than normal. A sink rate analysis of the main landing gear was performed. The maximum allowable main gear sink rate values are 9.6 ft/sec for a 212,000 lb. vehicle and 6.0 ft/sec for a 240,000 lb. vehicle. The landing weight of the STS-90 vehicle was estimated to be 232,965 lb.


Mission	Sink Rate
90	-6.0
3	-5.7
114	-5.5
120	-5.4
77	-4.6



## Groundspeed

Two potential consequences of off-nominal touchdown energies are high touchdown ground speed or excessive touchdown altitude rates. High ground speed at touchdown causes excessive tire wear during the initial spin up, which may result in a tire failure.


Nominal: The shuttle tires are certified to a ground speed up to 225 kts ground speed.  
(*Landing Procedures*, 4-25, 89)



## Touchdown Pitch Attitude

Nominal: Targeted touchdown pitch altitude is 8-10°, not to exceed 14°.

(Landing procedures, B-16, p.182; USA Entry, TAEM, and Approach/Landing Guidance, 4-13, p. 121)




## Equivalent Airspeed

Nominal:

Lightweight vehicles: 195 KEAS (+5, -10)

Heavyweight vehicles: 205 KEAS (+5, -10)

(*Landing Procedures*, 4-13, p. 77; 4-21, p.85; 4-24, p.88; *USA Entry TAEM*, and *Approach/Landing Guidance*, 4-13, p.122)



## **XIV. APPENDIX B: APOLLO LUNAR LANDING PERFORMANCE**

**Authors: Millard F. Reschke, Deborah L. Harm, William H. Paloski, Jacob J. Bloomberg, Scott J. Wood**

Surface obscuration during descent varied for each LM landing. For Apollo 11, visibility was degraded. During Apollo 12 and 15 the surface was completely obscured. For Apollo 14 and 17 LM descents, the surface was readily visible. Apollo 16 descent visibility was moderate ([Wagner 2006](#)).

Contributing factors that may be considered in mission and engineering design include:

- Landing site selection may be a factor in mission design, however, mission reports indicated that landing sites were similar, both in terms of characteristics and behavior of lunar dust.
- Improved hovering procedures may have contributed to improved visibility due to reduced blowing dust.
- Solar elevation angle may affect visibility.
- Landing radar output was affected by blowing dust and debris.

### **Apollo 11 – Crew: Armstrong, Collins, Aldrin; LM: Eagle; Launch: 16-Jul-69**

First manned landing on the Moon, July 20.

Ten meters above the surface, the lander started slipping to the left and rear. Armstrong, working with the controls, had apparently tilted the lander so the engine was firing against the flight path. With the velocity as low as it was at the time, the lander began to move backward. With no rear window to help him avoid obstacles behind the lander, he could not set the vehicle down and risk landing on the rim of a crater. He was able to shift the angle of the lunar module and stop the backward movement, but he could not eliminate the drift to the left. He was reluctant to slow the descent rate any further, but the figures Aldrin kept ticking off told him they were almost out of fuel. Armstrong was concentrating so hard on flying the lunar module that he was unable to perceive the first touch on the moon nor did he hear Aldrin call out "contact light," when the probes below the footpads brushed the surface. The lander settled gently down, like a helicopter, and Armstrong cut off the engine. Armstrong was not pleased with his piloting, but landing on the moon was much trickier than on the earth. He related the maneuver to his past experience in touching down during a ground fog, except that the moon dust had movement and that had interfered with his ability to judge the direction in which his craft was moving. Aldrin thought it "a very smooth touchdown," and said so at the time. They were tilted at an angle of 4.5 degrees from the vertical and turned 13 degrees to the left of the flight path trajectory.

## **Apollo 12 – Crew: Conrad, Gordon, Bean; LM: Intrepid, Launch: 14-Nov-69**

First precise manned landing on the Moon. Recovered part of Surveyor 3 probe.

As the lander passed north of Mare Nectaris, Conrad turned it on its back with the descent engine pointed along the flight path and switched the engine on to begin the final approach. Everything went exactly as expected, and after two minutes Conrad commented that it "feels good to be standing up in a g-field again." Three minutes later the module's attitude-control thrusters began firing busily - more than Conrad thought they should - but Houston assured him that all was well. After seven minutes Intrepid nosed over into a near-upright position and for the first time Conrad could see the lunar surface. The principal landmark identifying his landing point was a pattern of craters the astronauts called "Snowman"; Surveyor III lay halfway up the eastern wall of the crater that was the Snowman's torso, and Intrepid was targeted for the center of the crater. As soon as he could see out the window, Conrad cried delightedly, "Hey, there it is [Snowman]! There it is! Son of a gun! Right down the middle of the road!" Then, as Bean called out altitude, velocity, and quantity of fuel remaining; Conrad maneuvered the craft with his hand controller to pick a smooth spot to land on. The engine exhaust began kicking up dust about a hundred feet (30 meters) above the surface and by the time Intrepid reached 50 feet (15 meters) the cloud obscured the surface completely. At 1:54:36 a.m. EST on November 20, Pete Conrad made a blind landing - exactly where, he could not tell, but certainly close to the intended spot.

Conrad was naturally anxious to determine where he had set Intrepid down, and while he and Bean went through the post-landing check list they occasionally looked out the windows for landmarks that would allow Houston to pinpoint their location, but without success. After changing his mind a time or two, Conrad finally concluded, "I'm not sure that I'm not sitting right smack on the other side of the Surveyor crater, just a little bit past it." Two hours later, Dick Gordon in Yankee Clipper confirmed Conrad's guess when he sighted both Intrepid and Surveyor III through his sextant as he passed over the site. He told Houston that the lunar module was "on the left shoulder of the Snowman, about a third of the way from the Surveyor crater to the head. Post mission calculations placed Intrepid on the northwest rim of the Surveyor crater, 535 feet (163 meters) from the inert spacecraft. Had there been windows in the back of the lunar module, Conrad could have spotted the Surveyor as soon as the dust settled.

After postlanding checks of systems, Conrad and Bean described what they could see from their spacecraft. Intrepid had landed in undulating terrain pocked with craters ranging from a few feet to several hundred feet across, the larger ones rimmed by large blocks of rock. Numerous boulders, up to 20 feet (6 meters) in size, were scattered around the site, most of them angular rather than rounded, many showing fillets of dust around the base. Immediately in front of the landing craft Bean saw an area of "patterned ground" - parallel cracks in the surface soil perhaps an eighth of an inch (3 millimeters) deep. From the lunar module the crew could distinguish no color differences in the rocks or soil; everything seemed the same bright white.

### **Apollo 14 – Crew: Shepard, Roosa, Mitchell; LM: Antares; Launch: 31-Jan-71**

Shepard piloted the spacecraft to a routine landing 175 feet (53 meters) from its targeted landing site. During the final approach, they recognized Cone Crater right at pitch over and, soon thereafter, picked up the familiar pattern of smaller craters near their aim-point another mile or so to the west. There wasn't any really flat ground close at hand; there were either craters or sloping ground wherever he looked; but he had no trouble finding a crater-free, LM-sized patch that was only 30 meters from his target. The only problem with the landing spot was that it was on an eight degree slope; and, for 24 of the next 33 hours the astronauts had to contend with a tilting floor that threatened to dump Shepard over onto Mitchell's side of the spacecraft. In addition, the tilt contributed to a sleepless night between the two EVA's; but, otherwise, the LM attitude had no effect on the mission.

### **Apollo 15 – Crew: Scott, Worden, Irwin; LM: Falcon; Launch: 26-Jul-71**

First mission with the Lunar Rover vehicle.

The Hadley landing area is littered with craters but, as it turned out, few are large enough or deep enough to have early morning shadows. Toward the south edge of the landing area, virtually at the foot of Hadley Delta, Scott could see a grouping called the South Cluster and, of course, the rille was out in front of him. But, out in the middle, a couple of kilometers NNW of where Scott wanted to set down, landmarks are few and far between. There are a few moderate-sized craters, which, from the pre-flight analysis, looked as though they would have shadows in them at landing time; and Scott had spent time in the simulators learning to recognize them. However, as with the Apollo 14 site, the map and model makers had missed the subtle undulations that make this a rolling countryside and make the identification of small craters difficult at best. Part of the reason was that the Apollo 15 site is well north of the Moon's equator and the photo coverage didn't have the resolution that had been available for the earlier sites; and, the net effect was that, as Scott looked out the window, he couldn't find the patterns he had hoped to see. Just before pitchover when Scott was about 6 kilometers east of the target, Houston warned him that he was probably about a kilometer south of the planned track. A quick look at his position relative to the South Cluster and to the point on the rille where it makes a sharp bend at the foot of Hadley Delta gave Scott enough information to show him that Houston's call was a good one. So, he nudged Falcon's line-of-flight toward the north. Without clear local landmarks, he couldn't be certain of setting down right on target, but he'd be close enough that the difference wouldn't matter. He was about the right distance short of the rille (now trending northwest out in front of him); he was just about due north of South Cluster; and he was well out into the middle of the desired landing area. He was certainly within a few hundred meters of the target and, with the mobility that the Rover would give them, a miss of a few hundred meters would be a matter of only a few minutes' drive. The worst that could happen would be that they would spend a few extra minutes during their initial traverse getting their bearings.

**Apollo 16 – Crew: Young, Mattingly, Duke; LM: Orion; Launch: 16-Apr-72**

First landing in the lunar highlands.

This was referred to as a ‘near perfect landing.’ Young's greatest concern was the fact that there were few shadows to show him where the level spots were in this rolling terrain; and it was only in the last few moments that he got some clues from the LM shadow. It was Duke who saw the shadow first as they came down through 250 feet. Seconds later, as they were coming through 200 feet, Young rotated (yawed) the LM to the right and could see the shadow out his own window and use it to estimate his altitude and the sizes of the craters ahead. Through a combination of skill and luck, he set down on a remarkably level spot. As he and Duke discovered once they got outside, had they landed 25 meters in any direction from the actual landing site, [the LM] would have been on a local slope of six to ten degrees. In particular, in the final seconds, Young had to hover so that he could fly forward and to the right, just beyond a small crater fifteen meters in diameter. Only Armstrong and Conrad landed more upright, and both had the advantage of young, lightly-cratered, mare sites.

**Apollo 17 – Crew: Cernan, Evans, Schmitt; LM: Challenger; Launch: 7-Dec-72**

Final Apollo lunar mission.

Both crewmembers saw dust by 60-70 feet, although visibility through the final phase was excellent. “The tendency, once you redefine your landing area, is to become less concerned with your peripheral landmarks... you get more tunnel vision, and you are concerned with finding these specific touchdown points within that landing area. ... I can't say enough for what I consider the accuracy of the guidance. Manual control of the spacecraft was hard and firm ... exactly what I expected it to be.”

## Sources for Apollo Guidance/Navigation Information:

Apollo Lunar Surface Journal (mission reports, transcripts, etc.):

<http://www.hq.nasa.gov/alsj/frame.html>

NASA Document Index System (DIS)

<http://www6.isc.nasa.gov/dis/>

Manned Spaceflight PDF Documents

<http://www.geocities.com/bobandrepoint/spacepdf.htm>

JSC Engineering Directorate

[http://ea.isc.nasa.gov/Ea\\_web/HTML/index.asp](http://ea.isc.nasa.gov/Ea_web/HTML/index.asp)

Working on the Moon (Lessons from Apollo)

<http://www.workingonthemoon.com/index.html>

NASA Human Space Flight

<http://spaceflight.nasa.gov/history/shuttle-mir/people/oral-histories.htm>

## People contacted:

Eric Jones (editor, Apollo Lunar Surface Journal)

David Woods (Author: "How Apollo Flew to the Moon")

Floyd Bennett (designed the approach trajectories)

Ken Young (Apollo flight veteran)

Kim So (Library Technician)

Michele Hendershot (MOD Chief Curator)

Thomas Jones (astronaut)

## Documents:

Lunar Module Timelines (Apollo 12, 14, 16)

Aerodynamic Stability Characteristics of the Apollo Command Module

Postflight Entry Analysis

Mission Reports

Technical Reports

GNC Performance Analysis (Apollo 8, 9, 14, 15, 16)

Apollo Experience Report: Mission Planning for Lunar Module Descent and Ascent

Lunar Descent and Ascent Trajectories

A Study of Lunar module Navigation Systems Accuracies for Powered Descent, Ascents and Aborts

Study of Powered Descent Trajectories for Manned Lunar Landings

Downloaded several videos on Apollo Falls from various missions

Analysis of Apollo Navigation Accuracy

## **XV. APPENDIX C: INCIDENTS DUE TO SITUATIONAL AWARENESS DURING EVA**

**Authors: Millard F. Reschke, Deborah L. Harm, William H. Paloski, Jacob J. Bloomberg, Scott J. Wood**

### **MSS Lessons Learned**

#### **DX23 / ISS Mechanical & Robotics Systems Training**

April 18, 2007

#### **Scope of Lesson:**

This lesson is intended to inform crewmembers assigned to Station Robotics about the historical issues that have been overcome and why the system is operated the way it is today. The Robotics training flow is designed to gain generic understanding for operating the system as well as flight specific task familiarization. The operational habits discussed in this lesson apply to operating in both familiar and unfamiliar circumstances. These habits reinforce Space Flight Resource Management (SFRM) habits of Leadership, Communication and Situational Awareness. Many of the issues stem from failures in one or more of these areas.

#### **MSS Close Call Flight Incidents:**

##### **1. PDGF “Stiction” Problem**

#### ***Situation:***

Crew was releasing and backing off a grapple fixture. The normal method for this is to send an auto release command and the system relieves tension, opens the snares and extends the carriage forward. After the snares are opened, the operator starts backing off the grapple fixture so the carriage does not bump the fixture and pin. During this release the Latching End Effector (LEE) became stuck in the grapple fixture. Operator observed no motion so input higher hand controller command (building up forces in the arm). As the carriage moved forward it pushed the LEE off of the grapple fixture. The arm sprang off the fixture due to the force buildup and recoiled back striking the grapple fixture and lab (actual contact made and heard by crew inside).

#### ***Contributing factors:***

Situational Awareness - Loss of Situational Awareness of overlays and video. Not sure if crew noticed the overlay data indicating high command inputs but no actual motion. Video also showed no motion. If the crew noticed they did not react in time to prevent hitting.

#### ***Operational Changes:***

Situational Awareness - Education of crew on stiction problem - Train the crew on the signatures of a sticky grapple fixture. Overlay will indicate high level of commanded input but little to no actual FOR motion. Video will not show motion. Input should be removed immediately to limit arm force build up.

Changes to release technique - After this incident the ground team investigated causes of the stiction. No root cause has been determined but for expected grapple fixtures and scenarios a “push-off” technique is used to release. This technique allows the carriage to extend and “push” the LEE off the grapple fixture.

## **2. ISS-7A / STS-104 Close Call**

### **Space Station Remote Manipulator System (SSRMS) to Ultra High Frequency (UHF) Antenna Close Call**

#### **Situation:**

- The crew got ahead in the timeline and decision was made to install a third oxygen/nitrogen tank (ONTO). Tank was grappled by the SSRMS and handed off to EV crew at the airlock. At the end of the handoff the crew was asked to park the arm at a known location but the trajectory to get there was unfamiliar since the crew had not trained flying to a park position from the third tank install location.
- The big picture camera (Camera C on the Orbiter) was zoomed in on EV crew installing the ONTO tank while the arm was being maneuvered to park. This stage of assembly utilized station crewmembers for robotics operations and all three crewmembers were trained on the tasks. This maneuver was very late in the crew day and the “least” trained operator was performing this “simple” maneuver. The arm was maneuvered well away from structure and hit a six-degree of freedom (6 DOF) singularity causing the elbow to swing toward the UHF antenna. Eventually the camera C view was zoomed out to see whole arm, but elbow to UHF clearance not discernable. Good orthogonal camera views did not exist due to station configuration (no outboard structure yet to house a camera). Ground made a call to stop the arm and the Lab window was utilized to view actual clearance between arm and antenna. The ground tools showed 2 inches of clearance between arm and antenna when the arm stopped.
- The arm was maneuvered using a joint held algorithm. If a singularity was encountered, the arm would release the held joint and manage the singularity. This was identified by sluggish FOR motion, moving joint indication and pitch plane change (elbow joint swing) of the arm.

#### **Contributing Factors**

- Situational Awareness - Poor camera views - Crew had no big picture view. What had been a big picture view earlier, Cam C was left zoomed in on the ONTO. Crew was reluctant to ask shuttle crew to adjust cam C. Also the lack of cameras at this stage of assembly limited visibility into some clearances.
- Long Day, Tired Crew - The nominal plan for the day was to do two tanks, but they were ahead in the timeline and crew agreed to press with the third tank. After the mission, the crew reported that the arm ops were both physically and mentally draining.

- Unscripted Task/Unscripted Operator - There was never a plan or procedure to go from the 3rd ONTO to Lab FRGF. Therefore, this trajectory was never evaluated on the ground for clearance or singularities. This took them into the SY or WY singularity. Although the SSMRS Operator was fully certified on SSRMS, they had no experience operating in this area of the ISS. The operator that flew this last maneuver of the day was not the same operator that flew the EVA task just prior (the crew switched off operators).
- No Full-Time M2 - Because the day was longer than expected, the M2 was trying to do two jobs: M2 role and helping with IV Ops as EVA crew were ingressing the Airlock. M2 was not fully engaged in the final task.
- Crew was unaware of singularity indications - Crew was never trained to identify a singularity condition based on "sluggish" FOR motion, joint release and elbow swing. Generic Robotics Training and Spec Skills flow did not exist in this timeframe.

**Operational Changes:**

- Improvements in Communication - "All stop on the arm" voice protocol implemented - The call to crew was not very clean nor quick. A formalized method for declaring and implementing an All Stop was developed for both the Station and Shuttle Arm. Caution was taken to make the call and response similar for each system so cross trained crews would not be confused. New method has the person identifying the emergency issue declare all stop three times (All Stop, All Stop, All Stop). If heard the operator releases the hand controllers and safes the system. If called during a docked mission both the Shuttle and Station arms react to the same call so there is no reason to discern between the systems in trouble.
- Tight clearance scenarios implemented in Generic simulations & MSS lessons - The training team developed a series of scenarios and lessons to exercise tight clearance situations and unscheduled trajectory flying.
- Improvements in Situational Awareness - "Joint release" computation developed for the Mission Control Center (MCC) - The ground created an indication that the held joint had released. This would alert the ground team to possible elbow swing and they could make the call to crew.

**3. ISS-8A / STS-110**

**Operational Changes:**

No issues occurred during this mission that were SFRM errors. The arm did have a joint failure prior to flight, which was worked around using procedures and software changes. Operational changes came during training. The areas the arm was operating for the flight had numerous 6 DOF singularity zones. To avoid the earlier issues the community decided to start locking joints, which disabled the singularity management. If a singularity was encountered, the arm would simply stop and the team could decide how to continue with the task. Also in this timeframe the software was updated to include crew annunciations of

6 DOF singularities and enhancements to the Dynamic Onboard Ubiquitous Graphics (DOUG) tool provided better Situational awareness to the crews.

#### **4. ISS-UF2 / STS-111 Close Call SRMS to SSRMS Close Call**

##### **Situation:**

During a scheduled EVA activity, both the Shuttle and Station robotic arms were being used in close proximity to each other. The EV crews were directing the motion of both arms (called Ground Control Assist or GCA). At one point the two arms were moving at the same time toward each other. The ground called to have motion stopped (new All stop protocol was still being worked out so not utilized). Actual arm-to-arm distance is unknown but inside of desirable 5 feet.

##### **Contributing Factors:**

- Communication - Shuttle and Station arm operators were not coordinating motion. Communication between Shuttle and Station operators and ground teams not ideal in this scenario. There was no documented rules stating how to handle this situation.
- Situational Awareness - M1 not familiar with procedure. At this stage in assembly the Station crew was still utilized for arm operations and often times paired with a shuttle crewmember. Since Station crews could launch months ahead of the Shuttle mission, training the M1 and M2 together for tasks was very difficult to schedule. Many times the procedure would change after the Station crew had launched and they would have to adapt to those changes in real time.
- Communication and Situational Awareness - EV providing GCA to both arms. EVA tasks were happening in parallel and neither EV crew was very aware of the arm locations.

##### **Operational Changes:**

- Improvements in Situational Awareness - Ground developed a flight rule to minimize dual arm motion when in close proximity to each other. The flight rule (FR C12-7) states that simultaneous motion is allowed if 5 feet between the arms is maintained.
- Camera conflicts are minimized pre-flight since the Station and Shuttle arm operators can and do share cameras.
- Improvements in Communication - Shuttle and Station ground teams work together to develop a choreographed plan for arm motion. Procedures now contain callouts and sync points between the arms to minimized unnecessary dual motion. An example would be a callout in the Station procedure that says “give a go to the SRMS to maneuver to Install viewing position”. The Shuttle procedure would then say “Notify Station when at Install viewing position”.

## 5. ISS-9A / STS-112 Close Call SSRMS Elbow to PLB Door Close Call

### **Situation:**

During an EVA task the arm was in a closed configuration (not ideal). The EV crew was giving GCA calls to maneuver the arm and was helping some with helmet video views of clearances. This particular maneuver required rotation and translation commands in a particular sequence to limit the elbow swing of the arm. This swing was due to arm configuration (closed up so a very small angle between the tip and base booms) and not singularities. The crew was aware of the elbow swing from training. They had notes indicating how to avoid this effect. During the maneuver the prime camera for the elbow (Camera C from the Orbiter) got washed out with sunlight. Crew switched to a backup camera but mistook a box on the arm for the elbow so lost sight of closest object. The maneuver was flown as trained but response was not as expected and the elbow started to swing toward the Orbiter payload bay door. The All Stop was called by the ground and the proper response was issued. When the arm came to stop the ground tools indicated contact but no evidence of this was seen onboard (i.e. no damage to arm or door). Actual distance is unknown.

### **Contributing Factors:**

- Situational Awareness - Poor camera views. Crew lost the big picture view when Camera C was washed out. The backup view was misused when they thought they had the elbow in one of their scenes but the object turned out to be a box on the arm.
- Maneuver known to potentially cause large Elbow Pitch joint motion due to proximity of the tip of the SSRMS to its base. Crew was aware of the issue through training and had details in their crew notes concerning this maneuver.
- The technique used for the maneuver was not performing as expected. Crew made simultaneous translation and rotation inputs. Performing the rotation corrections first, minimizes the unwanted elbow joint motion. Crew commented that the on-orbit rotation response seemed less than what they had experienced during training.

### **Operational Changes:**

- Improvements to Situational Awareness - Ground now has camera pan/tilt data which can be sent to their clearance monitoring tool. This will help identify what the crew is watching and the ground team can direct them if they are watching something incorrectly. Crew is now provided with pan and tilt data in procedures so they can know they are looking at the proper things.
- Limit unscripted manual mode use. The arm is now maneuvered primarily by auto sequences or single joint mode. The manual operations are reserved for small EVA

maneuvers to final worksite and grapples. This allows the ground to evaluate issues and contingencies from known positions. The ground also designs better arm configurations for tasks to avoid having the arm closed up on itself.

- Improvements to Communication - Established sync points for communicating with EV crew and ground.

## 6. ISS-11A / STS-113 Incident

### Mobile Transporter hit the stowed P1 UHF Antenna

#### Situation:

During this mission the Mobile Transporter (MT) was timelined to translate after the deployment of the antenna. The EVA to deploy the antenna got behind and the deploy was delayed. The decision to translate was not altered and during the maneuver the MT ran over the antenna. When it hit, the ground didn't know what was going wrong. An EVA Crew inspection determined a box on the MT was hitting the UHF antenna on P1. The EV crew deployed the UHF antenna so the MT translation path was clear. Analysis failed to realize the antenna in its stowed state was in the translation corridor for the MT. There was some damage to the MT box but the antenna was fine.

#### Contributing Factors:

- Situational Awareness - Camera views were not required for the maneuver. Translation analysis had been done with P1 UHF antenna deployed since this was the expected configuration. During the fit test at Cape Canaveral, the UHF antenna was deployed, and they didn't check to see if the MT would fit past the antenna if it was stowed.
- P1 UHF antenna deployment was delayed because the EVA activity ran long. Failure of the ground team to determine the impact this might have on the translation.

#### Operational Changes:

Improvements to Situational Awareness - A flight rule now exists (FR B12-5) that states a CCTV survey of MT translation path is required before motion.

## 7. Other Onboard Events

#### Situation:

There have been several events where operators have made input errors in typing auto sequence numbers. All but one was caught prior to motion. The event where motion occurred was minor since only the wrist roll joint moved. Typically there are several checks in place to assure proper entry of numbers. Breakdowns in these checks have led to the mistakes. Breakdowns include having solo operators (no second set of eyes to verify what was entered or what should be entered), Crew fatigue (performing operations late in a crew's day) and improper mismatch or use of the M1 and M2.

**Contributing Factors:**

- Situational Awareness - Crew fatigue can lead to omitting or adding a sign change in the typed number. Can also lead to moving a decimal place or transposing numbers.
- Communication - Solo operations (without an M2) has led to issues.
- Leadership - Improper use of an M2 can lead to problems.

**Operational Changes:**

- Improvements in Situational Awareness - Educating the crew on when they are fatigued and steps they should take when they are in this state. Training touches on this by putting crews in timed situations adding pressure and trying to overload the operator. Educating the ground to be aware of what they are asking the crew to perform and the timing of the request (i.e. don't schedule tasks very late in the day without taking extra precautions).
- Improvements in Communication - New Ground rules and constraints are being generated to indicate the types of operations that can be done solo and which ones require an M2. There are also rules being generated to detail what steps should be taken if an M2 is not available. This includes more sync points with the ground and possible communication with station requirements for the operation to proceed.
- Improvements in Leadership - Steps are being taken to properly match skill levels of M1 and M2 operators. Keeping skill levels the same minimizes the chances that one operator will get ahead or behind the other. Training has developed a lesson to educate operators on proper M1 and M2 habits and roles. All arm operators take this class.

**Summary**

Robotics community has gained knowledge & experience. Operators' prevention goes back to SFRM basics of Command/Leadership, Communication/Workload Management, Situational Awareness and Decision Making. Ground has better mission design tools and real-time data. Today the arm is flown with limited Manual mode use, locking joints to limit singularity management, and rules are in place for coordinating and carrying out tasks. Procedures now contain Clearances, task unique item notes, coordination calls, sync points. Training has been updated to include the lessons learned in this class and throughout the training flow. There is no guarantee there will not be other incidents but the hope is the measures implemented will greatly reduce or eliminate the occurrences in the future.